

**Polarization transfer in the  $^{16}\text{O} (\vec{e}, e'\vec{p}) ^{15}\text{N}$  reaction.**

S. Malov, K. Wijesooriya, F.T. Baker, L. Bimbot, E.J. Brash, C.C. Chang,  
J.M. Finn, K.G. Fissum, J. Gao, R. Gilman, et al.

► **To cite this version:**

S. Malov, K. Wijesooriya, F.T. Baker, L. Bimbot, E.J. Brash, et al.. Polarization transfer in the  $^{16}\text{O} (\vec{e}, e'\vec{p}) ^{15}\text{N}$  reaction.. Physical Review C, American Physical Society, 2000, 62, pp.057302-057307. in2p3-00006477

**HAL Id: in2p3-00006477**

**<http://hal.in2p3.fr/in2p3-00006477>**

Submitted on 10 Oct 2000

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Polarization transfer in the  $^{16}\text{O}(\vec{e}, e'\vec{p})^{15}\text{N}$  reaction

S. Malov,<sup>25</sup> K. Wijesooriya,<sup>33</sup> F. T. Baker,<sup>7</sup> L. Bimbot,<sup>22</sup> E. J. Brash,<sup>24</sup> C. C. Chang,<sup>16</sup> J. M. Finn,<sup>33</sup> K. G. Fissum,<sup>17</sup> J. Gao,<sup>17</sup> R. Gilman,<sup>25,30</sup> C. Glashauser,<sup>25</sup> M. K. Jones,<sup>33</sup> J. J. Kelly,<sup>16</sup> G. Kumbartzki,<sup>25</sup> N. Liyanage,<sup>17</sup> J. McIntyre,<sup>25</sup> S. Nanda,<sup>30</sup> C. F. Perdrisat,<sup>33</sup> V. A. Punjabi,<sup>19</sup> G. Quémener,<sup>33</sup> R. D. Ransome,<sup>25</sup> P. M. Rutt,<sup>30</sup> D. G. Zainea,<sup>24</sup> B. D. Anderson,<sup>13</sup> K. A. Aniol,<sup>2</sup> L. Auerbach,<sup>29</sup> J. Berthot,<sup>1</sup> W. Bertozzi,<sup>17</sup> P. -Y. Bertin,<sup>1</sup> W. U. Boeglin,<sup>5</sup> V. Breton,<sup>1</sup> H. Breuer,<sup>16</sup> E. Burtin,<sup>26</sup> J. R. Calarco,<sup>18</sup> L. Cardman,<sup>30</sup> G. D. Cates,<sup>23</sup> C. Cavata,<sup>26</sup> J. -P. Chen,<sup>30</sup> E. Cisbani,<sup>12</sup> D. S. Dale,<sup>14</sup> R. De Leo,<sup>10</sup> A. Deur,<sup>1</sup> B. Diederich,<sup>21</sup> P. Djawotho,<sup>33</sup> J. Domingo,<sup>30</sup> B. Doyle,<sup>14</sup> J. -E. Ducret,<sup>26</sup> M. B. Epstein,<sup>2</sup> L. A. Ewell,<sup>16</sup> J. Fleniken,<sup>7</sup> H. Fonvielle,<sup>1</sup> B. Frois,<sup>26</sup> S. Frullani,<sup>12</sup> F. Garibaldi,<sup>12</sup> A. Gasparian,<sup>8,14</sup> S. Gilad,<sup>17</sup> A. Glamazdin,<sup>15</sup> J. Gomez,<sup>30</sup> V. Gorbenko,<sup>15</sup> T. Gorringer,<sup>14</sup> K. Griffioen,<sup>33</sup> F. W. Hersman,<sup>18</sup> J. Hines,<sup>7</sup> R. Holmes,<sup>28</sup> M. Holtrop,<sup>18</sup> N. d'Hose,<sup>26</sup> C. Howell,<sup>4</sup> G. M. Huber,<sup>24</sup> C. E. Hyde-Wright,<sup>21</sup> M. Iodice,<sup>12</sup> C. W. de Jager,<sup>30</sup> S. Jaminion,<sup>1</sup> K. Joo,<sup>32</sup> C. Jutier,<sup>1,21</sup> W. Kahl,<sup>28</sup> S. Kato,<sup>34</sup> S. Kerhoas,<sup>26</sup> M. Khandaker,<sup>19</sup> M. Khayat,<sup>13</sup> K. Kino,<sup>31</sup> W. Korsch,<sup>14</sup> L. Kramer,<sup>5</sup> K. S. Kumar,<sup>23</sup> G. Laveissière,<sup>1</sup> A. Leone,<sup>11</sup> J. J. LeRose,<sup>30</sup> L. Levchuk,<sup>15</sup> M. Liang,<sup>30</sup> R. A. Lindgren,<sup>32</sup> G. J. Lolos,<sup>24</sup> R. W. Lourie,<sup>27</sup> R. Madey,<sup>8,13,30</sup> K. Maeda,<sup>31</sup> D. M. Manley,<sup>13</sup> D. J. Margaziotis,<sup>2</sup> P. Markowitz,<sup>5</sup> J. Marroncle,<sup>26</sup> J. Martino,<sup>26</sup> J. S. McCarthy,<sup>32</sup> K. McCormick,<sup>21</sup> R. L. J. van der Meer,<sup>24</sup> Z. -E. Meziani,<sup>29</sup> R. Michaels,<sup>30</sup> J. Mougey,<sup>3</sup> D. Neyret,<sup>26</sup> E. A. J. M. Offermann,<sup>30</sup> Z. Papandreou,<sup>24</sup> R. Perrino,<sup>11</sup> G. G. Petratos,<sup>13</sup> S. Platchkov,<sup>26</sup> R. Pomatsalyuk,<sup>15</sup> D. L. Prout,<sup>13</sup> T. Pussieux,<sup>26</sup> O. Ravel,<sup>1</sup> Y. Roblin,<sup>1</sup> R. Roche,<sup>6</sup> D. Rowntree,<sup>17</sup> G.A. Rutledge,<sup>33</sup> A. Saha,<sup>30</sup> T. Saito,<sup>31</sup> A. J. Sarty,<sup>6</sup> A. Serdarevic-Offermann,<sup>24</sup> T. P. Smith,<sup>18</sup> A. Soldi,<sup>20</sup> P. Sorokin,<sup>15</sup> P. Souder,<sup>28</sup> R. Suleiman,<sup>13</sup> J. A. Templon,<sup>7</sup> T. Terasawa,<sup>31</sup> L. Todor,<sup>21</sup> H. Tsubota,<sup>31</sup> H. Ueno,<sup>34</sup> P. E. Ulmer,<sup>21</sup> G.M. Urciuoli,<sup>12</sup> P. Vernin,<sup>26</sup> S. van Verst,<sup>17</sup> B. Vlahovic,<sup>20</sup> H. Voskanyan,<sup>35</sup> J. W. Watson,<sup>13</sup> L. B. Weinstein,<sup>21</sup> R. Wilson,<sup>9</sup> B. Wojtsekhowski,<sup>30</sup> V. Zeps,<sup>14</sup> J. Zhao,<sup>17</sup> Z. -L. Zhou<sup>17</sup>

(The Jefferson Lab Hall A Collaboration)

<sup>1</sup> *Université Blaise Pascal/IN2P3, F-63177 Aubière, France*

<sup>2</sup> *California State University, Los Angeles, California, 90032*

<sup>3</sup> *Institut des Sciences Nucléaires, F-38026 Grenoble, France*

<sup>4</sup> *Duke University, Durham, North Carolina, 27706*

<sup>5</sup> *Florida International University, Miami, Florida, 33199*

<sup>6</sup> *Florida State University, Tallahassee, Florida, 32306*

<sup>7</sup> *University of Georgia, Athens, Georgia, 30602*

<sup>8</sup> *Hampton University, Hampton, Virginia, 23668*

<sup>9</sup> *Harvard University, Cambridge, Massachusetts, 02138*

<sup>10</sup> *INFN, Sezione di Bari and University of Bari, I-70126 Bari, Italy*

<sup>11</sup> *INFN, Sezione di Lecce, I-73100 Lecce, Italy*

<sup>12</sup> *INFN, Sezione Sanità and Istituto Superiore di Sanità, Laboratorio di Fisica, I-00161 Rome, Italy*

<sup>13</sup> *Kent State University, Kent, Ohio, 44242*

<sup>14</sup> *University of Kentucky, Lexington, Kentucky, 40506*

<sup>15</sup> *Kharkov Institute of Physics and Technology, Kharkov 310108, Ukraine*

<sup>16</sup> *University of Maryland, College Park, Maryland, 20742*

<sup>17</sup> *Massachusetts Institute of Technology, Cambridge, Massachusetts, 02139*

<sup>18</sup> *University of New Hampshire, Durham, New Hampshire, 03824*

<sup>19</sup> *Norfolk State University, Norfolk, Virginia, 23504*

<sup>20</sup> *North Carolina Central University, Durham, North Carolina, 27707*

<sup>21</sup> *Old Dominion University, Norfolk, Virginia, 23529*

<sup>22</sup> *Institut de Physique Nucléaire, F-91406 Orsay, France*

<sup>23</sup> *Princeton University, Princeton, New Jersey, 08544*

<sup>24</sup> *University of Regina, Regina, Saskatchewan, Canada, S4S 0A2*

<sup>25</sup> *Rutgers, The State University of New Jersey, Piscataway, New Jersey, 08854*

<sup>26</sup> *CEA Saclay, F-91191 Gif-sur-Yvette, France*

<sup>27</sup> *State University of New York at Stony Brook, Stony Brook, New York, 11794*

<sup>28</sup> *Syracuse University, Syracuse, New York, 13244*

<sup>29</sup> *Temple University, Philadelphia, Pennsylvania, 19122*

<sup>30</sup> *Thomas Jefferson National Accelerator Facility, Newport News, Virginia, 23606*

<sup>31</sup> *Tohoku University, Sendai 980, Japan*

<sup>32</sup> *University of Virginia, Charlottesville, Virginia, 22901*

<sup>33</sup> *College of William and Mary, Williamsburg, Virginia, 23187*

<sup>34</sup> *Yamagata University, Yamagata 990, Japan*

<sup>35</sup> *Yerevan Physics Institute, Yerevan 375036, Armenia*

(October 10, 2000)

The first ( $\vec{e}, e'\vec{p}$ ) polarization transfer measurements on a nucleus heavier than deuterium have been carried out at Jefferson Laboratory. Transverse and longitudinal components of the polarization of protons ejected in the reaction  $^{16}\text{O}(\vec{e}, e'\vec{p})$  were measured in quasielastic perpendicular kinematics at a  $Q^2$  of  $0.8 \text{ (GeV/c)}^2$ . The data are in good agreement with state of the art calculations.

PACS numbers: 13.40.Gp, 24.70.+s, 25.30.Fj, 27.20.+n

Polarization transfer in the ( $\vec{e}, e'\vec{p}$ ) reaction on a proton target is a direct measure of the ratio of the electric and magnetic form factors of the proton,  $G_E^p/G_M^p$ . When such measurements are carried out on a nuclear target, the polarization transfer observables are sensitive to the form factor ratio of the proton embedded in the nuclear medium. Because such experiments involve the measurement of ratios of polarizations at a single kinematic setting, the systematic errors are small, and different from those in standard Rosenbluth separations.

We report here measurements of polarization transfer in the  $^{16}\text{O}(\vec{e}, e'\vec{p})^{15}\text{N}$  reaction, the first such measurement on a nucleus heavier than deuterium [1]. The experiment, E89-033 at the Thomas Jefferson National Accelerator Facility (JLab), was part of the commissioning effort for Hall A [2]. It was the first experiment to use polarized beam at JLab, and the first to use the focal plane polarimeter (FPP) mounted on the high-resolution hadron spectrometer. Comparison of the results with state of the art calculations lays the groundwork for high precision tests of changes of the form factors in the nuclear medium. The distorted-wave impulse approximation (DWIA) provides a good description of the reaction, and the predictions are shown to be insensitive to various theoretical corrections.

The issue of possible modification of the properties of hadrons in the nucleus has been attracting experimental and theoretical attention for some years. It remains unsettled. Interpretations of inclusive ( $e, e'$ ) cross-section measurements in the y-scaling regime suggest that the radius of the nucleon is changed by less than a few percent at least for values of the four-momentum transfer squared ( $Q^2$ ) up to about  $1 \text{ (GeV/c)}^2$  [3]. These measurements are primarily sensitive to the magnetic form factor. Measurements of the Coulomb sum rule over a similar region in  $Q^2$  indicate that the electric form factor in  $^3\text{He}$  is close to its free value [4], and some studies suggest that this is true in  $^{12}\text{C}$  and  $^{56}\text{Fe}$  as well [5], but recent work disputes this conclusion [6]. Attempts to measure the ratio of electric to magnetic form factors of the nucleon in nuclei by Rosenbluth separations of cross sections in ( $e, e'p$ ) reactions indicated changes of about 25% at low  $Q^2$  [7], but some other experiments and theoretical analyses disagree [8]. Recent theoretical work by the Adelaide group based on the quark-meson coupling model predicted changes in the ratio of the two form factors for

$^{16}\text{O}$  of roughly 10% at  $Q^2$  around  $1 \text{ (GeV/c)}^2$  and about 20% or larger at about  $2.5 \text{ (GeV/c)}^2$  [9]. Changes of this magnitude have also been suggested previously [10].

For the free nucleon, the polarization transfer can be written in terms of the form factors as [11]

$$I_0 P_l' = \frac{E + E'}{m_p} \sqrt{\tau(1 + \tau)} G_M^2 \tan^2(\theta/2)$$

$$I_0 P_t' = -2\sqrt{\tau(1 + \tau)} G_M G_E \tan(\theta/2)$$

$$I_0 = G_E^2 + \tau G_M^2 [1 + 2(1 + \tau) \tan^2(\theta/2)]$$

$$\tau = Q^2/4m_p^2,$$

where  $E$  and  $E'$  are the energies of the incident and scattered electron,  $\theta$  is the electron scattering angle, and  $m_p$  is the proton mass.  $P_l'$  and  $P_t'$  are the longitudinal and transverse polarization transfer observables, respectively. The two components of the actual polarization in the scattering plane [12] are  $hP_l'$ , parallel to the proton momentum, and  $hP_t'$ , perpendicular to the proton momentum;  $h$  is the electron beam polarization. The measured polarizations change sign when the electron helicity changes sign, so these polarization transfer quantities are insensitive to instrumental asymmetries in the detectors.

The ratio of the transferred polarizations is then

$$\frac{P_t'}{P_l'} = \frac{-2m_p}{(E + E') \tan(\theta/2)} \frac{G_E}{G_M}.$$

For a free proton target, the ratio of polarizations can be used to determine the ratio of the form factors with small systematic errors; systematic problems associated with Rosenbluth separations are eliminated. The ratio is independent of the beam polarization (assuming it is not zero) and of the analyzing power of the proton polarimeter. One experimental datum requires a coincidence measurement at a single kinematic setting. The systematic error on the ratio of polarizations in the present experiment is about  $\pm 0.022$ , due almost entirely to uncertainty in the precession of the proton's spin in the hadron spectrometer. Even smaller errors have been achieved in the subsequent E93-027 measurements of the free form-factor ratio on a liquid hydrogen target at a  $Q^2$  of  $0.79 \text{ (GeV/c)}^2$  [13].

For nuclear targets, the polarization transfer observables depend sensitively on the nucleon form factors, but they depend also on the nuclear wave functions. In addition, they may be affected by final-state interactions of the outgoing proton, meson exchange and isobar currents, off-shell effects and the distortion of spinors by strong Lorentz scalar and vector potentials [14–19].

Electrons from the CEBAF accelerator of energy 2.45 GeV and longitudinal polarization about 30% were focussed on a waterfall target with three foils whose total thickness was about  $0.39 \text{ g/cm}^2$ . Scattered electrons

were detected in the focal plane array of the high resolution electron spectrometer in Hall A at a fixed laboratory angle of  $23.4^\circ$  and a fixed central energy of 2.00 GeV, the quasielastic peak. Protons with a fixed central momentum of 973 MeV/c were detected in coincidence with electrons in the focal plane array of the hadron spectrometer. Measurements in quasi-perpendicular kinematics were made at proton angles of  $53.3^\circ$  (for hydrogen data only),  $55.7^\circ$ , and  $60.5^\circ$ , corresponding to central missing momenta  $p_m$  of 0, 85, and 140 MeV/c. Elastic scattering from hydrogen dominates the spectrum at  $53.3^\circ$  and is visible also at  $55.7^\circ$ . The missing-mass resolution of about 1 MeV was sufficient to easily distinguish the  $p_{1/2}$  ground state of  $^{15}\text{N}$  from the strongly excited  $p_{3/2}$  state at 6.32 MeV, but small contributions from nearby weakly excited states could not be entirely excluded. In the continuum, a peak corresponding primarily to knockout of nucleons from the  $s_{1/2}$  shell rises weakly above a (physics) background presumably related to multi-particle knockout.

The JLab focal plane polarimeter (FPP) was designed and built by a collaboration of Rutgers, William & Mary, Georgia, Norfolk State, and Regina [20–22]. The polarimeter, consisting of four tracking straw chambers and a graphite analyzer set to a thickness of 22.5 cm for this experiment, is mounted in the hadron spectrometer behind vertical drift chambers and scintillators in the focal plane. The analyzing power  $A_c$  of the FPP was taken from the parametrization by McNaughton *et al.* [23]. Measured values of  $A_c$ , obtained by analyzing data from scattering on hydrogen, have been shown to agree well with this parameterization [13]. The beam polarization was measured at varying intervals with a Mott polarimeter in the injector beam line. For the 85 MeV/c data point on  $^{16}\text{O}$ , values of the beam polarization determined from the Mott polarimeter and from the FPP results for hydrogen are in good agreement, well within the 5% systematic error assigned to the beam polarization in the subsequent analysis of the oxygen data. The result for the ratio  $\mu \frac{G_E}{G_M}$  of hydrogen measured in this experiment at a  $Q^2$  of  $0.8 (\text{GeV}/c)^2$  is  $0.92 \pm 0.05$ , in agreement with previous results and with the values subsequently measured with higher precision [13].

Results for the transverse and longitudinal components of the polarization at the two central values of the missing momentum for the two bound states,  $p_{1/2}$  and  $p_{3/2}$ , and for the region of the unbound  $s_{1/2}$  state are shown in Fig. 1. The missing energy cuts on the latter were about 26 MeV wide. The polarizations are given in the scattering (lab) frame, defined by the incident and outgoing electron [12,15]. The errors shown are statistical. Systematic errors on the individual polarizations are about  $\pm 6\%$ , primarily due to the uncertainty in the polarization of the beam. Small acceptance averaging corrections are included [21], as are the effects of corrections to the

dipole approximation for spin transport of the proton in the hadron spectrometer [21,22]. Both corrections are generally less than about 2%. Radiative corrections, expected to be much smaller than the statistical errors here, have not been made [24].




FIG. 1. Measured values of the polarization transfer observables  $P'_l$  and  $P'_t$  for the  $^{16}\text{O}(\bar{e}, e'\bar{p})^{15}\text{N}$  reaction at  $Q^2 = 0.8 (\text{GeV}/c)^2$ . The theoretical curves represent plane-wave calculations (dotted) and distorted-wave calculations without spinor distortions (dashed) and with spinor distortions (dash-dot) by Kelly [18] and by Udias *et al.* [19] (solid).

The values of  $P'_l$  and  $P'_t$  for the free proton measured in this experiment (via the hydrogen in the waterfall target) are  $0.30 \pm 0.01$  and  $-0.20 \pm 0.01$ , with statistical errors. All the results for  $P'_l$  for  $^{16}\text{O}$  are within about one standard deviation of the free value, even those for the  $s_{1/2}$  region; their average value is 0.30. Several of the  $P'_t$  data points deviate somewhat from the free value; their average is -0.17. Such differences are not unexpected, even in the plane-wave impulse approximation (PWIA).

The curves in Fig. 1 represent theoretical calculations based upon one-body currents and free (MMD) [25] proton form factors. PWIA results, shown as dotted lines, are identical for the three states and, at  $p_m = 0$ , are equal to those for the free proton [15]. Final-state interactions included in DWIA calculations produce small state-dependent deviations from PWIA. The DWIA calculations by Kelly [15] are based upon a relativized Schrödinger equation and an effective momentum approximation (EMA) to the current operator. The dashed curves assume that lower and upper components of bound and ejectile spinors are related in the same way as for free protons [15]. The dash-dotted curves include

relativistic dynamics (spinor distortions) through the effect of Dirac scalar and vector potentials upon the effective current operator [18]. The solid curves show the results of calculations by Moya de Guerra and Udias [19] who solve the Dirac equation directly without using the EMA. All DWIA calculations shown used the same input as the calculations of unpolarized observables in Ref. [26]. These include the EDIAO optical model of Cooper *et al.* [27], NLSH bound-state wave functions [28], the Coulomb gauge, and the cc2 off-shell current operator. For modest  $p_m$ , the recoil polarization is relatively insensitive to variations of these choices.




FIG. 2. Measured values of the ratio of polarization transfer observables  $P'_t/P'_l$  for the  $^{16}\text{O}(\bar{e}, e'\bar{p})^{15}\text{N}$  reaction at  $Q^2 = 0.8 \text{ (GeV/c)}^2$ . The theoretical curves represent plane-wave calculations (dotted) and distorted-wave calculations without spinor distortions (dashed) and with spinor distortions (dash-dotted) [18]. The solid curves include the modifications of the nucleon predicted by Lu *et al.* [9] and spinor distortions.

All DWIA calculations are in reasonable agreement with the measured data. The two calculations which include relativistic dynamics are very similar over the relevant range of  $p_m$ . This is expected, since the results of the calculation of unpolarized observables in Ref. [26] suggested that Kelly's formulation is a good approximation to the more accurate approach of Udias *et al.* for  $p_m \lesssim 300 \text{ MeV/c}$ . The effects of relativity on the recoil polarization are small for this range of  $p_m$  and are dominated by distortion of the ejectile spinor.

Contributions from meson exchange (MEC) and isobar (IC) currents can also affect the recoil polarizations. Early calculations of the effects of MEC and IC were carried out by the Pavia group [14], and preliminary cal-

culations by M. Radici [16] of this group for the present kinematics have been made. Predictions of these effects using a different approach have been published recently by J. Ryckebusch *et al.* [17] for the present kinematics. The scale of these effects is typically comparable to the differences among the three DWIA curves shown. The DWIA with small corrections thus provides a firm baseline for considering changes in the form factor.

The ratios of the  $P'_t$  to  $P'_l$  data for the three states are plotted in Fig. 2. Three of the theoretical curves shown there correspond to those in Fig. 1, namely the PWIA (dots), and the DWIA without spinor distortion (dashes) and with spinor distortion (dash-dot) by Kelly [18]. The data are in good agreement with the three predictions, as expected from Fig. 1. The ratio of the experimental and theoretical values of  $P'_t/P'_l$  for the summed p state data is  $0.95 \pm 0.18$  using either DWIA calculation.

Deviations from unity significantly outside theoretical and experimental uncertainties would be evidence for changes in the nucleon form factor ratio in the nuclear medium. As noted in the introduction, the Adelaide group [9] obtained density-dependent form factors using a quark-meson coupling model and found changes in the form factor ratio for  $^{16}\text{O}$  of about 10% for  $Q^2 = 0.8 \text{ (GeV/c)}^2$ . The sensitivity of the  $(e, e'p)$  reaction to such changes has been estimated by Kelly using a local density approximation to the current operator. The fourth curve (solid) in Fig. 2 shows that the 10% changes in the form factors translate into changes of roughly 5% in the  $P'_t$  to  $P'_l$  ratio [18]. The reduced sensitivity of knockout at small  $p_m$  can be understood by comparing the averaging procedure used by Lu *et al.* with one more closely related to the matrix elements involved in the  $(e, e'p)$  reaction.

Lu *et al.* [9] estimated the effect of density dependence upon the electromagnetic form factors for a bound nucleon in orbital  $\phi_\alpha$  by using average form factors of the form

$$\bar{G}_\alpha(Q^2) \propto \int d^3r w_\alpha(r) G(Q^2, \rho_B(r)) \quad (1)$$

where  $\rho_B$  is the ground-state baryon density for the residual nucleus. Here proportionality denotes division by a similar integral omitting  $G$ . The static weighting factor  $w_\alpha(r) = |\phi_\alpha(r)|^2$  determines the effective density for different orbits. In the  $(e, e'p)$  reaction, however, Kelly [18] finds that the weighting factor is approximately

$$w_\alpha = \exp(i\mathbf{q} \cdot \mathbf{r}) \chi^{(-)}(\mathbf{p}', \mathbf{r})^* \phi_\alpha(\mathbf{r}) \quad (2)$$

where  $\chi$  is the distorted wave for ejectile momentum  $\mathbf{p}'$ ,  $\mathbf{q}$  is the momentum transfer, and  $\mathbf{p}_m = \mathbf{p}' - \mathbf{q}$ . In the interests of simplicity, recoil corrections and details of the current operator have been suppressed. In the plane-wave approximation, the weighting factor becomes

$$w_\alpha^{(\text{PWIA})} = \exp(-i\mathbf{p}_m \cdot \mathbf{r}) \phi_\alpha(\mathbf{r}) \quad (3)$$

and for  $p_m \rightarrow 0$  reduces to simply  $\phi_\alpha$ . In kinematic regimes explored thus far, this linear dependence upon  $\phi_\alpha$  reduces the effect of density dependence in the reaction, although the effect does increase with  $p_m$ . Furthermore, absorption and nonlocality corrections also reduce interior contributions to the average form factor.

Much more precise measurements of the  $P'_t/P'_l$  ratio are now possible. The polarized beam at JLab has shown a marked improvement in intensity, lifetime, and polarization since the commissioning experiment so that the statistical errors can be greatly reduced within reasonable running times, and systematic errors are already small. Conditions at the MAMI accelerator at Mainz are also appropriate for a high precision measurement at low  $Q^2$ , and an experiment was recently carried out there on  $^4\text{He}$  [29]. However, although recoil polarization provides a direct signal for medium modifications of nucleon form factors, the effect in  $(e,e'p)$  reactions is smaller than previously expected. A rigorous interpretation will require a unified relativistic treatment of the reaction and form factor models, including two-body currents.

The present experiment confirms the accuracy of the DWIA description of the reaction mechanism in this kinematical regime. The measured ratio of the transverse to longitudinal polarization transfers for the proton embedded in  $^{16}\text{O}$  at a  $Q^2$  of  $0.8 \text{ (GeV/c)}^2$  is in good agreement with calculations based on the free proton form factor with an experimental uncertainty of about 18%. The current generation of polarization transfer experiments should substantially improve this limit, but reliable identification of changes in the form factor in the medium remains an ambitious undertaking.

We are grateful to the entire technical staff at JLab for their dedicated and expert support of the commissioning experiment. We thank E. Moya De Guerra, M. Radici, J. Rycebusch and J. M. Udias for numerous discussions and detailed calculations. This work was supported in part by the National Science Foundation and the Department of Energy (U.S.), the National Institute for Nuclear Physics (Italy), the Atomic Energy Commission and the National Center for Scientific Research (France), and the Natural Sciences and Engineering Research Council (Canada).

- 
- [1] D. Eyl *et al.*, Z. Phys. **A352**, 211 (1995); B. D. Milbrath *et al.*, Phys. Rev. Lett. **80**, 452 (1998).
- [2] JLab Experiment E89-033, Spokespersons: C. C. Chang, C. Glashauser, S. Nanda, P. Rutt, unpublished.
- [3] I. Sick, Comm. Nucl. Part. Phys. **18**, 109 (1988).
- [4] R. Schiavilla, R. B. Wiringa, and J. Carlson, Phys. Rev. Lett. **70**, 3856 (1993).
- [5] J. Jourdan, Phys. Lett. **B353**, 189 (1995).
- [6] Z.-E. Meziani, private communication; Z.-E. Meziani and J. Morgenstern, submitted to Phys. Rev. Lett.
- [7] G. van der Steenhoven *et al.*, Phys. Rev. Lett. **57**, 182 (1986); G. van der Steenhoven *et al.*, Phys. Rev. Lett. **58**, 1727 (1987); D. Reffay-Pikeroen *et al.*, Phys. Rev. Lett. **60**, 776 (1988).
- [8] G. van der Steenhoven, Nucl. Phys. A527, 17c (1991), and references therein; L. Lapikas, Nucl. Phys. A553, 297c (1993); K.I. Blomqvist *et al.*, Z. Phys. **A351**, 353 (1995).
- [9] D. H. Lu *et al.*, Nucl. Phys. **A634**, 443 (1998); D. H. Lu *et al.*, Phys. Lett. **B417**, 217 (1998); D. H. Lu *et al.*, Phys. Rev. **C55**, 2628 (1998).
- [10] Ulf-G. Meissner, Phys. Rev. Lett. **62**, 1013 (1989); I. T. Cheon and M. T. Jeong, J. Phys. Soc. Jpn. **61**, 2726 (1992); M. R. Frank, B. K. Jennings and G. A. Miller, Phys. Rev. **C54**, 920 (1996).
- [11] A. I. Akhiezer and M. P. Rekalo, Sov. J. Part. Nucl. **3**, 277 (1974); R. Arnold, C. Carlson and F. Gross, Phys. Rev. **C23**, 363 (1981).
- [12] A. Raskin, T. W. Donnelly, Ann. Phys. **191**, 78 (1989).
- [13] JLab Experiment E93-027, C. F. Perdrisat, V. Punjabi and M. K. Jones, spokespersons; M. K. Jones *et al.*, Phys. Rev. Lett. **84**, 1398 (2000).
- [14] S. Boffi, C. Giusti, F.D. Pacati and M. Radici, Nucl. Phys. **A518**, 639 (1990).
- [15] J. J. Kelly, Phys. Rev. **C56**, 2672 (1997); Adv. in Nucl. Phys. **23**, 75 (1996).
- [16] M. Radici, private communication.
- [17] Jan Rycebusch, Dimitri Debruyne, Wim Van Nespren and Stijn Janssen, Phys. Rev. **C60**, 034604 (1999).
- [18] J. J. Kelly, Phys. Rev. **C60**, 044609 (1999).
- [19] E. Moya de Guerra and M. Udias, private communication; J. M. Udias, J. A. Caballero, E. Moya de Guerra, J. E. Amaro and T. W. Donnelly, Phys. Rev. Lett. **83**, 5451 (1999).
- [20] M. K. Jones *et al.*, in AIP Conference Proceedings 412, ed. T. W. Donnelly, 342 (1997).
- [21] K. Wijesooriya, Ph.D. thesis, College of William & Mary, (1999), unpublished.
- [22] S. Malov, Ph.D. thesis, Rutgers University (1999), unpublished.
- [23] M. McNaughton *et al.*, Nucl. Instr. Meth. **A241**, 435 (1985).
- [24] A. Afanasev, I. Akusevich and N. Merenkov, private communication.
- [25] P. Mergell, U.-G Meissner and D. Drechsel, Nucl. Phys. **A596**, 367 (1996).
- [26] J. Gao *et al.*, Phys. Rev. Lett. **84**, 3265, 2000.
- [27] E. D. Cooper *et al.*, Phys. Rev. **C47**, 297 (1993).
- [28] M. M. Sharma, M. A. Nagarajan and P. Ring, Phys. Lett. **B312**, 377 (1993).
- [29] MAMI Experiment A1/2-93 Addendum, J. Friedrich, R. Ransome, G. Rosner and H. Schmieden, 1998, unpublished.