



Presence of the solar de Vries cycle (205 years) during the last ice age

Gerhard Wagner, Jürg Beer, Jozef Masarik, Raimund Muscheler, Peter Kubik,
Werner Mende, Carlo Laj, Grant Raisbeck, Françoise Yiou

► To cite this version:

Gerhard Wagner, Jürg Beer, Jozef Masarik, Raimund Muscheler, Peter Kubik, et al.. Presence of the solar de Vries cycle (205 years) during the last ice age. *Geophysical Research Letters*, American Geophysical Union, 2001, 28 (2), pp.303-306. in2p3-00008529

HAL Id: in2p3-00008529

<http://hal.in2p3.fr/in2p3-00008529>

Submitted on 20 May 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Presence of the solar de Vries cycle (~205 years) during the last ice age

Gerhard Wagner, Jürg Beer, Jozef Masarik¹, Raimund Muscheler

Department of Surface Waters, EAWAG, Dübendorf, Switzerland

Peter W. Kubik

Paul Scherrer Institute, Zürich, Switzerland

Werner Mende

Freie Universität Berlin, Berlin, Germany

Carlo Laj

Laboratoire des Sciences du Climat et de l'Environnement, Gif-sur-Yvette, France

Grant M. Raisbeck and Francoise Yiou

Centre de Spectrométrie Nucléaire et de Spectrométrie de Masse, Orsay, France.

Abstract. Certain characteristic periodicities in the $\Delta^{14}\text{C}$ record from tree rings, such as the well-known 11-yr Schwabe cycle, are known to be of solar origin. The origin of longer-period cycles, such as the 205-yr de Vries cycle, in the $\Delta^{14}\text{C}$ record was less certain, and it was possible to attribute it either to solar or climatic variability. Here, we demonstrate that the de Vries cycle is present in ^{10}Be data from the GRIP ice core during the last ice age (25 to 50 kyr BP). Analysis of the amplitude of variation of this cycle shows it to be modulated by the geomagnetic field, indicating that the de Vries cycle is indeed of solar, rather than climatic, origin.

1. Introduction

The detection of solar cycles in ice cores using the cosmogenic radionuclides has important consequences for three distinct areas of research. Solar physicists are interested in the long-term behaviour of the length and the amplitude of the cycles in order to understand the underlying mechanisms that drive the solar dynamo [Beer *et al.*, 1998]. Climatologists on the other hand compare climate records with long records of solar variability in order to investigate what role solar variability plays in climate change [Beer *et al.*, 2000, Damon and Sonett, 1991]. Furthermore, climatologists are interested in the detection of solar cycles in different climate records because if the average cycle length is constant, they provide a natural clock that can be used to date different climate archives [Steig *et al.*, 1998].

Analysis of the cosmogenic radionuclides ^{10}Be and ^{36}Cl in ice cores permits the examination of solar variability through

the last 100 kyr or more, extending the $\Delta^{14}\text{C}$ record from tree rings [Stuiver *et al.*, 1998] by an order of magnitude. Several periodicities found in the $\Delta^{14}\text{C}$ data have been interpreted in terms of changing solar activity. These are the Gleissberg cycle (period of approximately 88 yr), the de Vries cycle (period of approximately 205 yr) and the Hallstatt cycle (period of approximately 2300 yr) [Stuiver and Braziunas, 1993, Damon and Sonett, 1991]. Preliminary results of spectral analysis of ^{10}Be profiles from the Dye 3 core (covering the last 550 yr) and the South Pole core (covering the last 1200 yr) suggest periodicities of about 90 and 200 yr [Beer *et al.*, 1994, Raisbeck *et al.*, 1990]. The well known 11-yr Schwabe cycle has been found in ^{10}Be records [Steig *et al.*, 1998, Beer *et al.*, 1990]. It is difficult to observe the Schwabe cycle in the $\Delta^{14}\text{C}$ data, because the amplitudes of short-term production changes are strongly attenuated due to the large sizes of the carbon reservoirs (atmosphere, biosphere and ocean). The solar origin of the Schwabe and the Gleissberg cycle signals in the proxy data is strongly supported by the historical sunspot record. The solar origin of the de Vries cycle is less well established due to lack of direct observational data. A solar relationship is suggested by the fact that maxima of the de Vries cycle in the $\Delta^{14}\text{C}$ data coincide with the Maunder (AD 1645-1715) and Spörer (AD 1420-1540) minima [Damon and Sonett, 1991, Eddy, 1976] of solar activity. The reconstruction of the long-term solar variability is not only fundamental to increasing our understanding of the solar dynamo, but also in determining the role of solar forcing in past and present climate changes [Beer *et al.*, 2000]. A possible solar forcing of climate change has been suggested by the coincidence of low $\Delta^{14}\text{C}$ values (Grand Maximum) with the "Medieval Warm Epoch" of the 11th through 13th centuries in western Europe. Further, the Maunder Minimum and the Spörer Minimum, which coincide with high $\Delta^{14}\text{C}$ values, fit with severe temperature dips during the Little Ice Age (14th to 19th centuries) [Eddy, 1976]. However new temperature reconstructions [Mann *et al.*, 1999] lead one to the question whether the "Medieval Warm Epoch" was a phenomenon that occurred all over the Northern

¹Now at Department of Nuclear Physics, Faculty of Mathematics and Physics, Comenius University, Mlynska dolina F/2, SK-842 14 Bratislava, Slovakia

Hemisphere. The de Vries cycle has been found e.g. in the tree ring widths [Sonett and Suess, 1984] of high altitude Bristercone pines dated from 3405 yr BC to 1885 yr AD and in the thermoluminescence record from a sea sediment covering the last 2500 yr [Cini Castagnoli et al., 1998]. These findings provide further evidence of a link between changing solar activity and climate on time scales of several thousands of years. Considering only the $\Delta^{14}\text{C}$ data it cannot be completely ruled out that the de Vries cycle is of climatic origin, in particular that it is due to changes of the carbon cycle. To separate climatic from production effects, it is useful to compare time series of ^{10}Be and $\Delta^{14}\text{C}$, because the relative production rate changes of these two isotopes due to solar and geomagnetic modulation are similar [Masarik and Beer, 1999, Lal and Peters, 1967], whereas their transport behaviour in the atmosphere is different [Bard et al., 1997, Beer et al., 1994]. ^{10}Be becomes fixed to aerosols and is washed out from the atmosphere by precipitation within about a year [Raisbeck et al., 1981]. In contrast, ^{14}C is oxidised to $^{14}\text{CO}_2$ and starts exchanging between atmosphere, biosphere and ocean [Siegenthaler et al., 1980]. Due to its relatively long atmospheric residence time it is well homogenised in the atmosphere. For a comparison of ^{10}Be and $\Delta^{14}\text{C}$, the high-resolution ^{10}Be data from the GRIP ice core provide an excellent data record for the last 100 kyr [Yiou et al., 1997a]. However, so far we have only analysed a continuous profile of ^{10}Be concentrations in the GRIP ice core for the last ice age, and not as yet for the last 10 kyr BP, which are covered by the $\Delta^{14}\text{C}$ data. Nevertheless it is possible to compare the ^{10}Be and $\Delta^{14}\text{C}$ records by considering their power spectra.

2. Data

The ^{10}Be concentration data from the GRIP ice core are based on ice samples of 55 cm length. In the time interval from 25 to 50 kyr BP, this corresponds to a mean time resolution of about 45 yr. Figure 1 shows in the upper panel the ^{10}Be concentration record as a function of time, using the depth age relation suggested by Johnsen et al. [1995]. The




Figure 1. ^{10}Be concentration and ^{10}Be flux as a function of age and depth for the time interval 25 to 50 kyr BP. The time scale and the used accumulation rate is based on the suggestions of Johnsen et al. [1995]. Note that the depth scale is not linear.




Figure 2. a) Lomb Scargle [Scargle, 1982; Haubold and Beer, 1992] power spectrum of the ^{10}Be flux in the time interval 25 to 50 kyr BP. b) Multitaper Method (MTM) harmonic analysis of the ^{10}Be flux. The parameter values are $N\Omega = 8$ (bandwidth), $K = 10$ (tapers). The left axis is the estimated MTM Amplitude (solid line) and the right axis is the statistical F-test (dotted line).

^{10}Be concentration in the GRIP ice core is the result of the interplay between production, transport and deposition. To compensate for changes in the ice accumulation rate the ^{10}Be concentrations were converted into fluxes (Figure 1) using the accumulation rate given by Johnsen et al. [1995].

3. Results and Discussion

The spectral analysis of the ^{10}Be flux record using the Lomb Scargle technique [Haubold and Beer, 1992, Scargle, 1982] is shown in Figure 2 a. The de Vries cycle is one of the most prominent periodicities in the ^{10}Be data, comparable to the power spectrum of the $\Delta^{14}\text{C}$ data [Stuiver and Braziunas, 1993]. After applying linear interpolation to produce equally spaced ^{10}Be flux data, other spectral analysis methods such as FFT (Fast Fourier Transformation), MEM (Maximum Entropy Method) and MTM (Multitaper Method) confirm the dominant period of 205 ± 5 yr obtained with the Lomb Scargle method. Figure 2 b shows the spectral analysis of the ^{10}Be flux using the MTM technique [Yiou et al., 1997b, Paillard et al., 1996]. A statistical F-test performed on the MTM amplitude shows that the de Vries cycle is significant on the 99 % level. To exclude the possibility that the spectral analysis of the ^{10}Be flux only shows the de Vries cycle due to a periodicity in the accumulation rate of approximately 205 yr (^{10}Be flux \equiv ^{10}Be concentration * snow accumulation rate), we repeated the spectral analysis for the ^{10}Be concentration data. The de Vries cycle is also present in the ^{10}Be concentration data, which strengthens our result. Nevertheless in this paper we refer to the ^{10}Be flux, because it is a better proxy of the ^{10}Be production rate.

Having identified the de Vries cycle in the ^{10}Be data, we now address the question of whether this cycle is of solar or

climatic origin. If it is of solar origin its amplitude should be modulated by the geomagnetic field. However amplitude modulation by the geomagnetic field is not expected if the de Vries cycle is of climatic origin. The dependence of the ^{10}Be production rate on the geomagnetic field and on the solar activity has been calculated by *Masarik and Beer* [1999]. Figure 3 shows the mean global production rate of ^{10}Be as a function of the geomagnetic field intensity and the solar activity. The solar activity is described by the solar modulation parameter Φ ($\Phi = 0$ MeV corresponds to a quiet and $\Phi \geq 1000$ MeV to a very active sun) [*Masarik and Beer*, 1999]. Figure 3 also shows that for a constant solar activity the production rate of ^{10}Be increases with decreasing geomagnetic field intensity. For example, a reduction of the geomagnetic field intensity from its present value to zero would increase the production rate of ^{10}Be by a factor of 2.7 for $\Phi = 0$ MeV and 1.8 for $\Phi = 1000$ MeV, respectively. Moreover Figure 3 shows that for a fixed variation of the solar modulation parameter a lower geomagnetic field intensity leads to a greater amplitude in the ^{10}Be production signal than a higher field intensity. For example a change of the solar modulation parameter from $\Phi = 1000$ MeV to $\Phi = 0$ MeV leads to a 3.8 times larger change in the production rate if the geomagnetic field is reduced from the present value to zero.

We now investigate how well these theoretical considerations agree with the experimental data. Assuming that the ^{10}Be flux reflects the mean global ^{10}Be production rate [*Wagner et al.*, 2000, *Steig et al.*, 1996, *Beer et al.*, 1990] and using published paleogeomagnetic field variations [*Laj et al.*, 2000; *Tric et al.*, 1992] the geomagnetic modulation effect on the de Vries cycle can be investigated. Support for this assumption is based on the fact that central Greenland receives a substantial part of its precipitation from lower latitudes [*Johnsen et al.*, 1989] and many chemical species found in the ice originate from eastern Asia [*Biscaye et al.*, 1997]. The main trends of the geomagnetic field intensity during the last ice age are fairly well known. In the lower panel of Figure 4 two estimations of the paleogeomagnetic field intensity relative to the level of today are shown, which are based on remanence measurements from six sediment




Figure 3. Dependence of mean global ^{10}Be production rate on the relative geomagnetic field intensity B [normalised to present intensity] and on the solar modulation parameter Φ [MeV].




Figure 4. Upper panel: Band pass filtered ^{10}Be flux [$(215 \text{ yr})^{-1} \leq \tau \leq (195 \text{ yr})^{-1}$] [*Paillard et al.*, 1996] over the time period of 25 to 50 kyr BP. Lower panel: paleomagnetic field intensities during the last ice age: The first (dotted line) is derived from remanence measurements of several Mediterranean sediment cores [*Tric et al.*, 1992]. The second (solid line) is based on remanence measurements of several North Atlantic sediment cores [*Laj et al.*, 2000]. The shaded area corresponds to the $\pm 2\sigma$ uncertainties. Both records are low pass filtered (cut off frequency = $1/3 \text{ kyr}^{-1}$) to obtain a comparable time resolution as the envelope of the band pass filtered ^{10}Be flux.

cores of the North Atlantic [*Laj et al.*, 2000] and from four sediment cores of the Mediterranean sea [*Tric et al.*, 1992]. The North Atlantic Paleointensity Stack since 75 ka (NAPIS-75) [*Laj et al.*, 2000], originally placed on the GISP2 time scale [*Groote et al.*, 1997] is transferred to the GRIP time scale [*Johnsen et al.*, 1995] by synchronising the $\delta^{18}\text{O}$ records. The paleogeomagnetic field intensity record based on the Mediterranean sediments is transferred to the GRIP time scale as described in *Wagner et al.* [2000]. The minimum of the geomagnetic field intensity at about 38.5 kyr BP corresponds to the so called Laschamp event. Figure 4 shows in the upper panel the band pass filtered ^{10}Be flux with a frequency range between $(215 \text{ yr})^{-1}$ and $(195 \text{ yr})^{-1}$. During the minimum of the magnetic field at about 38.5 kyr BP the amplitude of the de Vries cycle is about twice as high as during time periods, when the geomagnetic field intensity is approximately equal to today. The shared variance (r^2) between the envelope of the amplitude of the band pass filtered ^{10}Be signal and the geomagnetic field data is 30 % for the NAPIS-75 record and 33 % for the Mediterranean record, which is significant on the 99 % level. After low pass filtering the envelope of the amplitude with the same cut off frequency (3 kyr^{-1}) as the geomagnetic field data were low pass filtered, the shared variances increase to values of about 40 %. Taking into account that the amplitude of solar activity cycles changes with time (e.g. sun spot cycle) [*Beer et al.*, 2000, *Beer et al.*, 1998], a shared variance of 30–40 % can be considered as high.

This geomagnetic modulation of the amplitude of the de Vries cycle strongly points to a solar origin of the periodicity of approximately 205 yr in the ^{10}Be flux. In addition the presence of the de Vries cycle in the GRIP ice core confirms that the relative time scale between 25 and 50 kyr BP is precise.

Acknowledgments. We wish to thank C. Stengel, C. Obrist and S. Bollhalder and the crew of the ETH-Zürich AMS facility. We are indebted to J. Rippin and S. Baumgartner for fruitful discussions and to E. Steig for critical and helpful review. This work was supported by the Swiss National Science Foundation.

References

- Bard, E., G. M. Raisbeck, F. Yiou and J. Jouzel, Solar modulation of cosmogenic nuclide production over the last millennium: comparison between ^{14}C and ^{10}Be records, *Earth. Planet. Sci. Lett.* 150, 453-462, 1997.
- Beer, J., A. Blinov, G. Bonani, R.C. Finkel, H. J. Hofmann, B. Lehmann, H. Oeschger, A. Sigg, J. Schwander, T. Staffelbach, B. Stauffer, M. Suter and W. Wölfli, Use of ^{10}Be in polar ice to trace the 11-year cycle of solar activity, *Nature* 347, 164-166, 1990.
- Beer, J., C. F. Joos, C. Lukaszczuk, W. Mende, U. Siegenthaler, R. Stellmacher and M. Suter, ^{10}Be as an indicator of solar variability and climate, in: The solar engine and its influence on terrestrial atmosphere and climate, edited by E. Nesme-Ribes, pp. 221-233, Springer-Verlag, Berlin, 1994.
- Beer, J., S. M. Tobias and N. O. Weiss, An active Sun throughout the Maunder minimum, *Solar Physics* 181, 237-249, 1998.
- Beer, J., W. Mende, R. Stellmacher, The role of the sun in climate forcing, *Quaternary Science Reviews* 19, 403-415, 2000.
- Biscaye, P. E., F. E. Grousset, M. Revel, S. Van der Gaast, G. A. Zielinski, A. Vaars, and G. Kukla, Asian provenance of glacial dust (stage 2) in the Greenland Ice Sheet Project 2 Ice Core, Summit, Greenland, *J. Geophys. Res.* 102, NO C12, 26765-26781, 1997.
- Cini Castagnoli, G., G. Bonino, P. Della Monica, S. Procopio and C. Taricco, On the solar origin of the 200 yr Suess wiggles: evidence from thermoluminescence in sea sediments, *Il Nuovo Cimento* 21 C, 237-241, 1998.
- Damon, P. E. and C. P. Sonett, Solar and terrestrial components of the atmospheric C-14 variation spectrum, in: The sun in time, edited by C. P. Sonett, M. S. Giampapa and M. S. Matthews, pp. 360-388, The University of Arizona, Tucson, 1991.
- Eddy, J. A. The Maunder Minimum, *Science* 192, 1189-1201, 1976.
- Grootes, P. M., M. Stuiver, $^{18}\text{O}/^{16}\text{O}$ variability in Greenland snow and ice with 10^3 to 10^5 year time resolution, *J. Geophys. Res.* 102, NO C12, 26455-26470, 1997.
- Haubold, H. J. and J. Beer, Solar activity cycles revealed by time series analysis of argon-37, sunspot-number, and beryllium-10 records, Proc. IUGG, Vienna, Solar-Terrestrial Variability and Global Change, 11-34, 1992.
- Johnsen, S. J., W. Dansgaard and J. W. C. White, The origin of Arctic precipitation under present and glacial conditions, *Tellus* 41B, 452-468, 1989.
- Johnsen, S. J., D. Dahl-Jensen, W. Dansgaard and N. Gundestrup, Greenland palaeotemperatures derived from GRIP bore hole temperature and ice core isotope profiles, *Tellus* 47B, 624-629, 1995.
- Laj, C., C. Kissel, A. Mazaud, J. E. T. Channell and J. Beer, High resolution paleointensity stack from the North Atlantic for the 75-10 kyr interval (NAPIS-75) and the duration of the Laschamp event, *Phil. Trans. R. Soc. Lond. A* 358, 1009-1025, 2000.
- Lal, D and B. Peters, Cosmic Ray Produced Radioactivity on the Earth, in: Handbuch der Physik, S. Flügge, ed, Springer Verlag, Berlin, pp. 551-612, 1967.
- Mann, M. E., R. S. Bradley, and M. K. Hughes, Northern Hemisphere Temperatures During the Past Millennium: Inferences, Uncertainties, and Limitations, *Geophys. Res. Lett.* 26, 759-762, 1999.
- Masarik, J. and J. Beer, Simulation of particle fluxes and cosmogenic nuclide production in the Earth's atmosphere, *J. Geophys. Res.* 104, 12099 - 13012, 1999.
- Paillard, D., L. Labeyrie and P. Yiou, Macintosh program performs time-series analysis, *Eos Trans. AGU* 77, 379, 1996.
- Raisbeck, G. M., F. Yiou, M. Fruneau, J.M. Loiseaux, M. Liewuin and J.C. Ravel, Cosmogenic $^{10}\text{Be}/^{10}\text{Be}$ as a probe of atmospheric transport processes, *Geophys. Res. Lett.* 8, 1015-1018, 1981.
- Raisbeck, G. M., F. Yiou, J. Jouzel and J. R. Petit, ^{10}Be and $\delta^2\text{H}$ in polar ice cores as a probe of the solar variability's influence on climate, *Phil. Trans. R. Soc. Lond. A* 330, 65-72, 1990.
- Scargle, J. D., Studies in astronomical time series analysis. II. Statistical aspects of spectral analysis of unevenly spaced data, *Astrophys. J.* 263, 835-853, 1982.
- Siegenthaler, U., M. Heimann and H. Oeschger, ^{14}C Variations Caused by Changes in the Global Carbon Cycle, *Radiocarbon* 22, 177-191, 1980.
- Sonett, C. P. and H. E. Suess, Correlation of bristlecone pine ring widths with atmospheric ^{14}C variations: A climate-Sun relation, *Nature* 307, 141-143, 1984.
- Steig, E. J., P. J. Polissar, M. Stuiver, P. M. Grootes and R. C. Finkel, Large amplitude solar modulation cycles of ^{10}Be in Antarctica: implications for atmospheric mixing processes and interpretation of the ice-core record., *Geophys. Res. Lett.* 23, 523-526, 1996.
- Steig, E. J., D. L. Morse, E. D. Waddington and P. J. Polissar, Using the sunspot cycle to date ice cores, *Geophys. Res. Lett.* 25, 163-166, 1998.
- Stuiver, M. and T. F. Braziunas, Sun, Ocean, Climate and Atmospheric $^{14}\text{CO}_2$, an evaluation of causal and spectral relationships, *The Holocene* 3, 289-305, 1993.
- Stuiver, M., P. J. Reimer, E. Bard, J. W. Beck, G. S. Burr, K. A. Hughen, B. Kromer, G. McCormac, J. Van der Plicht and M. Spurk, INTCAL98 Radiocarbon age calibration, 24,000-0 cal BP, *Radiocarbon* 40, 1041-1083, 1998.
- Tric, E., J. P. Valet, P. Tucholka, M. Paterne, L. LaBeyrie, F. Guichard, L. Tauxe and M. Fontugne, Paleointensity of the geomagnetic field during the last 80,000 years, *J. Geophys. Res.* 97, 9337-9351, 1992.
- Wagner, G., J. Masarik, J. Beer, S. Baumgartner, D. Imboden, P. W. Kubik, H.-A. Synal and M. Suter, Reconstruction of the geomagnetic field between 20 and 60 ka from cosmogenic nuclides in the GRIP ice core, *Nuclear Instruments and Methods B172*, in press, 2000.
- Yiou, F., G. M. Raisbeck, S. Baumgartner, J. Beer, C. Hammer, S. Johnsen, J. Jouzel, P. W. Kubik, J. Lestringuez, M. Stiévenard, M. Suter and P. Yiou, Beryllium 10 in the Greenland Ice Core Project ice core at Summit, Greenland, *J. Geophys. Res.* 102, 26783-26794, 1997a. [Meanwhile in the considered depth interval of this paper about 160 more Be-10 samples (mostly processed with 0.45 micron filters) from the GRIP ice core were analysed in Zürich compared to the number of ^{10}Be samples presented in Yiou et al..]
- Yiou, P., K. Fuhrer, L. D. Meeker, J. Jouzel, S. Johnsen, and P. A. Mayewski, Paleoclimatic variability inferred from the spectral analysis of Greenland and Antarctic ice-core data., *J. Geophys. Res.* 102, NO C12, 26441-26454, 1997b.

G. Wagner, J. Beer, J. Masarik¹, R. Muscheler, Department of Surface Waters, EAWAG, Überlandstr.133, CH-8600 Dübendorf, Switzerland. (e-mail: gerhard.wagner@eawag.ch, beer@eawag.ch)

P. Kubik, Paul Scherrer Institute, c/o ETH-Hönggerberg, Building HPK-H33, CH-8093 Zürich, Switzerland.

W. Mende, Freie Universität Berlin, c/o Berlin Brandenburgische Akademie der Wissenschaften, Jägerstr. 22/23, D-100117 Berlin, Germany.

C. Laj, Laboratoire des Sciences du Climat et de l'Environnement, Unité Mixte CEA-CNRS, Avenue de la Terrasse, Batiment 12, 91198 Gif-sur-Yvette, France.

G. M. Raisbeck, F. Yiou, Centre de Spectrométrie Nucléaire et de Spectrométrie de Masse, IN2P3-CNRS, 91405 Orsay, France.

(Received April 4, 2000; revised August 2, 2000; accepted September 28, 2000)