

Higgs searches at L3

D. Teyssier

▶ To cite this version:

D. Teyssier. Higgs searches at L3. High Energy Physics Conference - HEP-MAD'01, Sep 2001, Antananariv, Madagascar. pp.67-72. in2p3-00011094

HAL Id: in2p3-00011094 https://hal.in2p3.fr/in2p3-00011094

Submitted on 15 Jan2002

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

HIGGS SEARCHES AT L3

DANIEL TEYSSIER

on behalf of the L3 Collaboration IPN Lyon, 4 rue Enrico Fermi, Campus de la Doua, 69622 Villeurbanne Cedex, FRANCE E-mail: teyssier@cern.ch

Standard and non-minimal Higgs searches were performed with data collected by the L3 detector in e^+e^- collisions at 189 – 202 GeV center-of-mass energies. The standard search methodology is described. An example of non-standard search is reviewed as well and results at the above mentionned center-of-mass energies are presented.

1 Standard Model Higgs searches.

1.1 Introduction.

The Standard Model (SM) provides fermions and gauge bosons masses through electroweak symmetry breaking, the so-called Higgs mecanism ¹. Introducing one complex scalar Higgs doublet, the three gauge bosons W^{\pm} and Z^0 get their longitudinal polarisation, and become massive ². It remains one degree of freedom, corresponding to a new fundamental scalar particle, the Higgs boson.

1.2 Production mecanism.

The main process to produce the Higgs boson at LEP2 is the Higgs-Strahlung. The corresponding Feynman diagram is presented below (Fig. 1). An off-shell Z boson is produced, and is decaying into an on-shell Z boson and a Higgs boson.



Figure 1. Higgs boson processes : left) Higgs-Strahlung, right) W or Z fusion.

proceedings: submitted to World Scientific on December 20, 2001

Especially in the missing and the leptonic channels, the Higgs boson can be produced by W or Z fusion (Fig. 1). But the cross section is small compared to Higgs-Strahlung process (Fig. 2).



Figure 2. Cross section for Higgs boson production.

1.3 Higgs boson decay.

The coupling between the Higgs boson and the charged fermions is proportionnal to their masses. In the range of mass of interest, the Higgs boson will decay preferentially into b quarks (Fig. 3). The Z boson branching ratios will determine the different topologies.

1.4 Four topologies.

The Z boson can decay into a pair of quarks (four-jet topology, 65% of the total number of events), or a pair of neutrinos (missins energy channel, 19% of events), or a pair of electrons or muons (leptonic channels, 6% of events). Finally, the Z or the Higgs boson can decay into a pair of taus, and the other particle into a pair of quarks (in this case, we get two jets and two mini-jets from the taus, 9% of events).

1.5 Analysis procedure.

First of all, loose cuts are used to reduce the most copious backgrounds, like the *two-photon* background $(e^+e^- \rightarrow e^+e^-q\bar{q})$. Afterwards, we apply tighter cuts for each specific topology. For instance, we demand two tagged jets in

proceedings: submitted to World Scientific on December 20, 2001

 $\mathbf{2}$



Figure 3. Higgs boson branching ratios.

all topologies as the Higgs boson will decay into $b\bar{b}$. So we are able to build a global variable, which can either be the output of a neural network or a likelihood function. Combining this information with the Higgs mass measurement, we obtain the final discriminant variable (Fig. 4), used to calculate the confidence level (CL) for each mass hypothesis.



Figure 4. $Hq\bar{q}$ search channel : left) Higgs mass distribution, right) likelihood output.

proceedings: submitted to World Scientific on December 20, 2001

1.6 Confidence levels.

We construct a statistical estimator (designed here by -2ln(Q)), depending on the Higgs mass hypothesis, so as to test the compatibility of the data with the *signal plus background* hypothesis, or the *background* only hypothesis. This estimator is taken to be :

$$-2ln(Q(m_H)) = 2s_{tot} - 2\sum N_i \cdot ln[1 + s_i(m_H)/b_i]$$
(1)

where m_H is the Higgs mass hypothesis, s_i and b_i the expected signal and background numbers of events respectively. N_i takes the values of d_i (data), b_i or $s_i + b_i$ following the hypothesis. By integrating the -2ln(Q) distributions, we can obtain :

• the $(1 - CL_b)$ (Fig. 5) representing the compatibility between the data and the SM background expectations.

• the CL_s (Fig. 5) which allows us to set a limit.



Figure 5. Confidence levels.

The observed lower limit on m_H is 112.0 GeV at the 95% CL, while the expected lower limit is 112.4 GeV³. For $m_H = 115$ GeV, the background probability $(1 - CL_b)$ is 32%. Preliminary combined searches from the four LEP experiments, Aleph, Delphi, Opal and L3 give an observed lower bound on its mass at a 95% CL⁴: $m_H > 114.1$ GeV. On the other hand, precision measurements of electroweak observables, which involve virtual Higgs boson exchanges, give an upper bound of ⁵: $m_H < 210$ GeV at a 95% CL.

proceedings: submitted to World Scientific on December 20, 2001

2 Beyond Standard Model Higgs searches.

2.1 Two Higgs doublet models of type II

In general two Higgs doublet models of type II ², five scalar Higgs bosons appear after symmetry breaking. Both searches for the production of the h and A Higgs bosons in the processes $e^+e^- \rightarrow hZ$ and $e^+e^- \rightarrow hA$ are used. We did not use any b-tagging in order to perform a flavour-independent search, since the coupling of the Higgs boson to the b quarks could be suppressed.

The dominant production mechanisms are the following :



Figure 6. hZ and hA processes

The cross sections can be parametrized as a function of the Standard Model (SM) cross sections as :

$$\sigma_{hZ} = \rho.\sigma_{hZ}^{SM} \tag{2}$$

$$\sigma_{hA} = \eta \cdot \lambda \cdot \sigma_{hZ}^{SM} \tag{3}$$

where σ_{hZ}^{SM} is the cross section for $e^+e^- \to hZ$ in the SM, ρ and η are the couplings, to the Z normalized to the SM ones

$$\rho = \frac{g_{hZZ}^2}{(g_{hZZ}^{SM})^2} \tag{4}$$

$$\eta = \frac{g_{hAZ}^2}{(g_{hZZ}^{SM})^2} \tag{5}$$

and λ is a phase space factor.

2.2 One example : the search for $h, A \rightarrow q\bar{q}\tau^+\tau^-$

We give the result for a search of the process $h, A \to q\bar{q}\tau^+\tau^-$, with an upper limit on the η parameter (Fig. 7). The analysis procedure is the same as that described in Ref. ⁶.

proceedings: submitted to World Scientific on December 20, 2001

 $\mathbf{5}$



Figure 7. Upper limit on the η parameter.

3 Conclusion.

There was an excellent LEP performance in 2000 : \sqrt{s} up to 209 GeV, and more than 200 pb^{-1} of integrated luminosity collected per experiment. The observed lower bound on SM Higgs boson found by L3 is 112.0 GeV. Other results concerning flavour-independent analyses and Minimal Supersymetric Standard Model searches can be found in : http://l3www.cern.ch/conferences/contributions.html

References

- F. Englert et R. Brout, *Phys.Rev.Lett.*13(1964)321, P.W. Higgs, *Phys.Rev.Lett.*12(1964)132, T. Kibble, *Phys.Rev.*155(1967)1554.
- J. Gunion, H. Haber, G. Kane et S. Dawson, *The Higgs Hunter's Guide*, Frontiers in Physics Series (vol. 80), éd. Addison-Wesley (1990).
- L3 Collaboration, P. Achard et al., Standard model Higgs boson with the L3 experiment at LEP, Phys. Lett. B 517(2001) 319-331.
- ALEPH, DELPHI, L3 et OPAL Collaborations, The LEP working group for Higgs boson searches, Search for the Standard Model Higgs boson at LEP, preliminary CERN-EP/2001.
- The LEP Collaborations ALEPH, DELPHI, L3 et OPAL, the LEP Electroweak Working Group and the SLD Heavy Flavour and Electroweak Groups, A Combination of Preliminary Electroweak Measurements and Constraints on the Standard Model, CERN-EP/2001-021.
- 6. L3 Collaboration, M. Acciarri et al., Phys. Lett. B 471 (1999) 321.

proceedings: submitted to World Scientific on December 20, 2001