

Search for excited leptons at LEP

P. Achard, O. Adriani, M. Aguilar-Benitez, J. Alcaraz, G. Alemanni, J. Allaby, A. Aloisio, M.G. Alviggi, H. Anderhub, V.P. Andreev, et al.

► **To cite this version:**

P. Achard, O. Adriani, M. Aguilar-Benitez, J. Alcaraz, G. Alemanni, et al.. Search for excited leptons at LEP. Physics Letters B, Elsevier, 2003, 568, pp.23-34. in2p3-00013742

HAL Id: in2p3-00013742

<http://hal.in2p3.fr/in2p3-00013742>

Submitted on 26 Aug 2003

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Search for Excited Leptons at LEP

The L3 Collaboration

Abstract

A search for charged and neutral excited leptons is performed in 217 pb^{-1} of data collected with the L3 detector at LEP at centre-of-mass energies up to 209 GeV. The pair- and single-production mechanisms are investigated and no signals are detected. Combining with L3 results from searches at lower centre-of-mass energies, gives improved limits on the masses and couplings of excited leptons.

Submitted to *Phys. Lett. B*

1 Introduction

Charged (e^* , μ^* , and τ^*) and neutral (ν_e^* , ν_μ^* and ν_τ^*) excited leptons are predicted by composite models where leptons and quarks have substructure [1–3]. These models address fundamental questions left open by the Standard Model [4], such as the number of families and the fermion mass values.

High energy electron-positron annihilations constitute an excellent environment for the search for excited leptons and several searches have been carried out at LEP [5–7]. Searches for excited electrons and neutrinos were also performed at the HERA [8] ep collider. This Letter describes the extension of previous L3 searches [5, 6] to the highest centre-of-mass energies attained by the LEP machine in its last year of operation, $\sqrt{s} = 202 - 209$ GeV.

A model [1] is assumed in which excited leptons are described as isospin doublets with left and right handed components:

$$L^* = \begin{pmatrix} \nu^* \\ \ell^* \end{pmatrix}_L + \begin{pmatrix} \nu^* \\ \ell^* \end{pmatrix}_R, \quad (1)$$

with $\ell = e, \mu, \tau$ and $\nu = \nu_e, \nu_\mu, \nu_\tau$. These excited leptons are accessible at LEP through pair production, $e^+e^- \rightarrow \ell^*\ell^*$, $\nu^*\nu^*$, or single production, $e^+e^- \rightarrow \ell\ell^*$, $\nu\nu^*$. Pair-production searches are sensitive to excited leptons of mass up to values close to the kinematic limit $\sqrt{s}/2$, while the single-production mechanism extends the search potential up to masses close to \sqrt{s} .

The pair production of excited leptons is described by the effective Lagrangian:

$$\mathcal{L}_{L^*L^*} = \bar{L}^*\gamma^\mu \left(g\frac{\vec{\tau}}{2}\vec{W}_\mu + g'YB_\mu \right) L^*. \quad (2)$$

while the single production of excited leptons and their decay into Standard Model leptons is modeled by:

$$\mathcal{L}_{L^*L} = \frac{1}{\Lambda}\bar{L}^*\sigma^{\mu\nu} \left(gf\frac{\vec{\tau}}{2}\partial_\mu\vec{W}_\nu + g'f'Y\partial_\mu B_\nu \right) \frac{1-\gamma^5}{2}L + (\text{h.c.}). \quad (3)$$

In these expressions, γ^μ are the Dirac matrices, $\sigma^{\mu\nu} = i[\gamma^\mu, \gamma^\nu]/2$, g and g' are the Standard Model SU(2) and U(1) coupling constants, $\vec{\tau}$ denotes the Pauli matrices, $Y = -1/2$ is the hypercharge and \vec{W} and B are the gauge fields associated with the SU(2) and U(1) groups respectively. L denotes the Standard Model leptons, Λ is the scale of the New Physics responsible for the existence of excited leptons and f and f' scale the SU(2) and U(1) couplings, respectively.

The cross section for pair production of excited leptons depends only on their mass and on \sqrt{s} . As an example, for masses of the excited leptons of 101 GeV, the cross section at $\sqrt{s} = 206$ GeV for $e^+e^- \rightarrow \mu^*\mu^*$ and $e^+e^- \rightarrow \nu_\mu^*\nu_\mu^*$ are 0.6 pb and 0.3 pb respectively. The single-production cross section for excited leptons depends also (equation 3) on f/Λ and f'/Λ .

Equation 3 also describes the decay of excited leptons into Standard Model leptons in association with a photon or a gauge boson. Three decays are possible: radiative decays, $\ell^* \rightarrow \ell\gamma$ and $\nu^* \rightarrow \nu\gamma$, charged-current decays, $\ell^* \rightarrow \nu W$ and $\nu^* \rightarrow \ell W$, and neutral-current decays $\ell^* \rightarrow \ell Z$ and $\nu^* \rightarrow \nu Z$. The branching fractions in these different modes depend on the relative values of f and f' . As an example, Table 1 lists these fractions for two mass values and the the two extreme cases $f = f'$ and $f = -f'$. Table 2 summarises all the final states for excited lepton production and decay that are considered in this Letter.

2 Data and Monte Carlo Samples

The data sample discussed in this letter comprises 216.9 pb^{-1} collected with the L3 detector [9] at $\sqrt{s} = 202 - 209 \text{ GeV}$ with an average centre-of-mass energy of 206 GeV .

Monte Carlo samples of pair-produced excited leptons are generated for a mass of 101 GeV , which is close to the expected kinematic limit. The single production of excited leptons is modeled for masses of $110, 160$ and 201 GeV . An interpolation, which includes samples at $101, 150$ and 195 GeV produced at $\sqrt{s} = 192 - 202 \text{ GeV}$ [6], allows the estimation of selection efficiencies in the mass range from 90 to 209 GeV . All possible final states listed in Table 2 are generated, including both hadronic and leptonic decays of the W and Z bosons. Differential cross sections are modeled according to Reference 1 and initial state radiation is taken into account in cross section calculations.

Standard Model background processes are simulated with several Monte Carlo generators. Radiative Bhabha events are generated using BHWIDE [10] and TEEGG [11]. KK2F [12] is used for the $e^+e^- \rightarrow \mu\mu(\gamma)$, $e^+e^- \rightarrow \tau\tau(\gamma)$, $e^+e^- \rightarrow \nu\nu(\gamma)$ and $e^+e^- \rightarrow q\bar{q}(\gamma)$ processes. The $e^+e^- \rightarrow W^+W^-$ process is modeled with KORALW [13], with the exception of the $q\bar{q}e\nu$ final state, described by EXCALIBUR [14] which is also used for the $e^+e^- \rightarrow llll$ and $e^+e^- \rightarrow ll\nu\nu$ processes. PYTHIA [15] is used for the final states coming from $e^+e^- \rightarrow ZZ$ not covered by EXCALIBUR. GGG [16] describes the $e^+e^- \rightarrow \gamma\gamma(\gamma)$ process. The production of hadrons and leptons in two-photon interactions is described by PHOJET [17] and DIAG36 [18], respectively.

The L3 detector response is simulated for all Monte Carlo samples using the GEANT program [19], which includes the effects of energy loss, multiple scattering and showering in the detector. Time dependent detector behaviour, as monitored during the data taking period, is also taken into account.

3 Selection Strategy

The search for charged and neutral excited leptons follows the one already performed at lower centre-of-mass energies [5,6]. Several selections are devised in order to cover all final states listed in Table 2. All those selections proceed from the identification of photons, leptons and jets and then assemble those constituents to single out the particular signature of each final state.

Photons and electrons are identified in the BGO electromagnetic calorimeter as clusters within a polar angle $|\cos\theta| < 0.95$ and with energy above 1 GeV . The shape of the shower in the crystals must be compatible with a photon or an electron. Electrons must be associated to tracks in the central tracker, while no track is allowed near photon candidates.

Muons are selected from tracks reconstructed in the central or forward-backward muon spectrometers. These tracks must point to the interaction vertex. Signals in the time-of-flight system are used to reject background from cosmic rays. In addition, muon candidates without a track in the spectrometer are also built starting from energy depositions in the calorimeters consistent with a minimum ionising particle matched to a track in the central tracker.

Tau leptons are reconstructed from low multiplicity narrow hadronic jets, or identified through their decay into electrons or muons with additional missing energy (\cancel{E}) and momentum.

Jets are reconstructed from charged tracks and energy clusters in the electromagnetic and hadronic calorimeters. The missing energy and missing momentum of the event, used to tag events with neutrinos in the final state, are calculated from all tracks, energy clusters and additional muons. The missing momentum is required to point away from the beam axis in

most of the selections, so as to minimise the background from two-photon interactions and fermion pair production in association with a high energy initial state photon radiated at low polar angle.

A short summary of the relevant features of the analyses for the pair- and single-production mechanisms is given in the following two sections. Details on the different selections are discussed elsewhere [6, 20].

4 Pair Production

The search for pair-produced charged and neutral excited leptons relies on many selections, aimed to maximise the sensitivity for excited lepton masses close to the kinematic limit, around $\sqrt{s}/2$. For these mass values, radiative and charged-current decays include over 80% of the decay modes, for any choice of the couplings. Neutral-current decays are hence not considered. Table 3 summarises the selections used for the analysis of each final state, as described below.

Radiative decays of charged excited leptons are searched for through exclusive final states with a pair of identified leptons and two identified photons. Final states with two photons and missing energy and momentum are used to tag radiative decays of excited neutrinos.

Four selections are devised to identify the pair of W bosons produced by the charged-current decays of both charged and neutral excited leptons. Three selections, collectively denoted as $q\ell\nu$, and different according to the lepton flavour, select final states with two hadronic jets and an isolated high energy lepton. The fourth selection, $qqqq$, identifies high multiplicity final states, with four hadronic jets. The $q\ell\nu$ and $qqqq$ selections are used for the charged-current decays of charged excited leptons and tau excited neutrinos. Figure 1a shows the hadronic mass for events selected by the $q\ell\nu$ selection. In Figure 1b is shown the sum of the invariant and recoil masses for two jets associated to a reconstructed W in the $qqqq$ selection. In the search for electron and muon excited neutrinos, an additional pair of electrons and muons is required in addition to the $q\ell\nu$ and $qqqq$ selections.

Events where one excited lepton decays radiatively and the other through the charged-current are investigated with a similar selection for both charged and neutral excited leptons. Events are selected that have an isolated lepton, a photon and missing energy. In addition, the decay products of a W boson are required as either two hadronic jets or an additional isolated lepton. Figures 1c and 1d show the masses of the excited electron and muon candidates, reconstructed as the photon-lepton invariant mass.

5 Single Production

The search for singly produced excited leptons complements the search through the pair-production mechanism and gives access to the mass range from $\sqrt{s}/2$ up to \sqrt{s} . To retain the highest efficiency, all decay modes are investigated: radiative, charged-current and neutral-current. Table 4 lists the association between all final states and the corresponding selections which are summarised below.

In the search for charged excited leptons decaying radiatively, final states with two leptons and a photon are selected. High sensitivity to the excited lepton mass is achieved via the invariant mass of a detected lepton and the photon, as presented in Figures 2a, 2b and 2c. Events with one photon and large missing energy and momentum are used to search for the

possible production of excited neutrinos. Figure 2d presents the normalised energy spectrum of the selected photons.

For the charged-current decay of charged and neutral excited leptons, the $qq\ell\nu$ selection described above is used. Figure 3 presents the distribution used for the reconstruction of the mass of the excited lepton candidates, namely the lepton recoil mass in the case of charged excited leptons and the invariant mass of the lepton and the detected jets in the case of excited neutrinos with a hadronic decay of the W boson. To retain the highest efficiency, two additional selections are devised. They identify events with hadronic activity and missing energy, $qq\cancel{E}$, as well as leptons and missing energy, $\ell\ell\cancel{E}$. These selections complement the $qq\ell\nu$ selection in the case in which the lepton is not detected or the W boson decays into leptons.

Neutral-current decays of charged excited leptons are searched for with the same selections used for the charged-current case. The neutral-current decay of excited neutrinos, produces a Z boson and missing energy. This final state is covered by the $qq\cancel{E}$ and $\ell\ell\cancel{E}$ selections.

6 Results

Tables 3 and 4 list the number of events observed in the data by each selection together with the Monte Carlo background expectations and the signal efficiencies. No evidence for the production of excited leptons is observed in any final state.

Systematic uncertainties affect the results in Tables 3 and 4. An uncertainty of 1.5 – 2.5%, depending on the selection, is associated to the background estimate. This includes the uncertainties on the cross sections of background processes, limited Monte Carlo statistics, detector simulation and the selection procedure. The limited Monte Carlo statistics and the detector simulation also affect the estimation of the signal efficiency. Depending on the selection, the systematic uncertainty from this source is around 2.5%, which also covers the accuracy of the efficiency interpolation for different excited lepton masses.

The absence of excited leptons in the data sample is expressed by means of upper limits on their masses and couplings. Absolute limits on the masses are derived from the pair-production process, whereas limits on the effective couplings as a function of the masses are derived from the single-production study. The limits discussed in the following also include the results from previous searches at $\sqrt{s} = 189$ GeV [5] and $\sqrt{s} = 192 - 202$ GeV [6]. All limits take systematic uncertainties into account and are reported for two extreme scenarios, $f = f'$ and $f = -f'$. For $f = f'$, the radiative decay is allowed for charged excited leptons whereas it is forbidden for excited neutrinos. The opposite holds for $f = -f'$. The final state topologies and hence the experimental sensitivity for the two scenarios are therefore very different. For the pair-production process, the numbers of observed and expected events, together with the signal efficiencies are translated into upper limits on the cross section for the production of excited leptons. A scan is performed for all possible values of the ratio f/f' , and the decay fractions of charged excited leptons and excited neutrinos are computed for the radiative, charged-current and mixed decay modes. Signal cross sections are calculated and the corresponding mass limits are derived, as presented in Table 5. More stringent limits are obtained for channels with low background. As an example, the limits on charged excited leptons in the $f = f'$ scenario and excited neutrinos in the $f = -f'$ scenario benefit from a large branching ratio in the clean radiative decay channel. In addition to those corresponding to the $f = f'$ and $f = -f'$ scenarios, limits are also given for all excited leptons which correspond to the lowest values obtained in the scan over f/f' , and hence valid for any choice of the couplings.

In the case of single-production searches, an upper limit on the cross section is obtained as

a function of the excited lepton mass. Different mass hypotheses are investigated by means of the distributions of the variables presented in Figures 2 and 3. These cross section limits are translated in the upper limits on the ratios $|f|/\Lambda$ and $|f'|/\Lambda$ shown in Figure 4. The edge of the curves at low mass indicates the lower mass limit derived from pair-production searches, whereas the rise at high mass reflects the decrease of the expected signal cross section and therefore of the experimental sensitivity. The limits corresponding to charged excited leptons in the $f = f'$ scenario and excited neutrinos in the $f = -f'$ scenario are derived mainly from the radiative decay searches, whose clean final states have a high signal sensitivity. These limits are more stringent than those obtained in the complementary scenarios in which the radiative decays are forbidden. The limits corresponding to excited leptons of the first generation are significantly tighter due to their higher cross section resulting from the t -channel contribution.

In conclusion, no evidence for charged and neutral excited leptons of any flavour is found in the LEP data, and lower mass limits as high as 101.5 GeV are derived for any value of the excited lepton couplings. Upper limits on $|f|/\Lambda$ and $|f'|/\Lambda$, ranging from 10^{-1} to 10^{-4} GeV $^{-1}$ according to the excited lepton flavour and mass, are set in the mass range from 100 to 200 GeV.

References

- [1] K. Hagiwara *et al.*, *Z. Phys.* **C 29** (1985) 115.
- [2] M.B. Voloshin *et al.*, *Sov. Phys. JETP* **64** (1986) 446;
M.B. Voloshin, *Phys. Lett.* **B 209** (1988) 360;
F. Boudjema *et al.*, *Phys. Lett.* **B 240** (1990) 485;
F. Boudjema *et al.*, *Z. Phys.* **C 57** (1993) 425.
- [3] F. Boudjema *et al.*, *Z Physics at LEP 1*, eds. J. Ellis *et al.*, Vol. 2, p. 188, Report CERN/89/08 (1989), and references therein.
- [4] S.L. Glashow, *Nucl. Phys.* **22** (1961) 579;
A. Salam, in *Elementary Particle Theory*, ed. N. Svartholm, (Almqvist and Wiksell, Stockholm, 1968);
S. Weinberg, *Phys. Rev. Lett.* **19** (1967) 1264.
- [5] L3 Coll., M. Acciarri *et al.*, *Phys. Lett.* **B 473** (2000) 177.
- [6] L3 Coll., M. Acciarri *et al.*, *Phys. Lett.* **B 502** (2001) 37.
- [7] ALEPH Coll., D. Buskulic *et al.*, *Phys. Lett.* **B 385** (1996) 445;
DELPHI Coll., P. Abreu *et al.*, *E. Phys. J.* **C 8** (1999) 41;
OPAL Coll., G. Abbiendi *et al.*, *Eur. Phys. J.* **C 14** (2000) 73, *Phys. Lett.* **B 544** (2002) 57.
- [8] H1 Coll., C. Adloff *et al.* *Phys. Lett.* **B 525** (2002) 9; *Phys. Lett.* **B 548** (2002) 35
ZEUS Coll., S. Chekanov *et al.*, *Phys. Lett.* **B 549** (2002) 32.
- [9] L3 Coll., B. Adeva *et al.*, *Nucl. Instr. Meth.* **A 289** (1990) 35;
L3 Coll., O. Adriani *et al.*, *Phys. Rep.* **236** (1993) 1;
M. Chemarin *et al.*, *Nucl. Instr. Meth.* **A 349** (1994) 345;
M. Acciarri *et al.*, *Nucl. Instr. Meth.* **A 351** (1994) 300;

- G. Basti *et al.*, Nucl. Instr. Meth. **A 374** (1996) 293;
A. Adam *et al.*, Nucl. Instr. Meth. **A 383** (1996) 342.
- [10] BHWIDE version 1.01 is used.
S. Jadach *et al.*, Phys. Rev. **D 40** (1989) 3582, Comp. Phys. Comm. **70** (1992) 305, Phys. Lett. **B 390** (1997) 298.
- [11] TEEGG Monte Carlo.
D. Karlen, Nucl. Phys. **B 289** (1987) 23.
- [12] KK2F version 4.12 is used.
S. Jadach *et al.*, Comp. Phys. Comm. **130** (2000) 260.
- [13] KORALW version 1.33 is used.
S. Jadach *et al.*, Comp. Phys. Comm. **94** (1996) 216;
M. Jadach *et al.*, Phys. Lett. **B 372** (1996) 289.
- [14] EXCALIBUR Monte Carlo.
F.A. Berends *et al.*, Comp. Phys. Comm. **85** (1995) 437.
- [15] PYTHIA versions 5.722 and 6.1 are used.
T. Sjöstrand, Preprint CERN-TH/7112/93 (1993), revised August 1995; Comp. Phys. Comm. **82** (1994) 74; Preprint hep-ph/0001032 (2000).
- [16] GGG Monte Carlo.
F.A. Berends *et al.*, Nucl. Phys. **B 186** (1981) 22;
CALKUL Coll., F.A. Berends *et al.*, Nucl. Phys. **B 239** (1984) 395.
- [17] PHOJET version 1.05 is used.
R. Engel, Z. Phys. **C 66** (1995) 203;
R. Engel *et al.*, Phys. Rev. **D 54** (1996) 4244.
- [18] DIAG36 Monte Carlo.
F.A. Berends *et al.*, Nucl. Phys. **B 253** (1985) 441.
- [19] GEANT version 3.15 is used.
R. Brun *et al.*, Preprint CERN-DD/EE/84-1 (1984), revised 1987.
The GHEISHA program (H. Fesefeldt, RWTH Aachen Report PITHA 85/02 (1985)) is used to simulate hadronic interactions.
- [20] M. Falagan, Ph.D thesis, Complutense Univeristy, Madrid (2000).

The L3 Collaboration:

P.Achard,²⁰ O.Adriani,¹⁷ M.Aguilar-Benitez,²⁴ J.Alcaraz,²⁴ G.Alemanni,²² J.Allaby,¹⁸ A.Aloisio,²⁸ M.G.Alvigi,²⁸ H.Anderhub,⁴⁶ V.P.Andreev,^{6,33} F.Anselmo,⁸ A.Arefiev,²⁷ T.Azmoon,³ T.Aziz,⁹ P.Bagnaia,³⁸ A.Bajo,²⁴ G.Baksay,²⁵ L.Baksay,²⁵ S.V.Baldew,² S.Banerjee,⁹ Sw.Banerjee,⁴ A.Barczyk,^{46,44} R.Barillere,¹⁸ P.Bartalini,²² M.Basile,⁸ N.Batalova,⁴³ R.Battiston,³² A.Bay,²² F.Becattini,¹⁷ U.Becker,¹³ F.Behner,⁴⁶ L.Bellucci,¹⁷ R.Berbeco,³ J.Berdugo,²⁴ P.Berges,¹³ B.Bertucci,³² B.L.Betev,⁴⁶ M.Biasini,³² M.Biglietti,²⁸ A.Biland,⁴⁶ J.J.Blaising,⁴ S.C.Blyth,³⁴ G.J.Bobbink,² A.Böhm,¹ L.Boldizar,¹² B.Borgia,³⁸ S.Bottai,¹⁷ D.Bourilkov,⁴⁶ M.Bourquin,²⁰ S.Braccini,²⁰ J.G.Branson,⁴⁰ F.Brochu,⁴ J.D.Burger,¹³ W.J.Burger,³² X.D.Cai,¹³ M.Capell,¹³ G.Cara Romeo,⁸ G.Carlinio,²⁸ A.Cartacci,¹⁷ J.Casaus,²⁴ F.Cavallari,³⁸ N.Cavallo,³⁵ C.Cecchi,³² M.Cerrada,²⁴ M.Chamizo,²⁰ Y.H.Chang,⁴⁸ M.Chemarin,²³ A.Chen,⁴⁸ G.Chen,⁷ G.M.Chen,⁷ H.F.Chen,²¹ H.S.Chen,⁷ G.Chiefari,²⁸ L.Cifarelli,³⁹ F.Cindolo,⁸ I.Clare,¹³ R.Clare,³⁷ G.Coignet,⁴ N.Colino,²⁴ S.Costantini,³⁸ B.de la Cruz,²⁴ S.Cucciarelli,³² J.A.van Dalen,³⁰ R.de Asmundis,²⁸ P.Déglon,²⁰ J.Debreczeni,¹² A.Degré,⁴ K.Dehmelt,²⁵ K.Deiters,⁴⁴ D.della Volpe,²⁸ E.Delmeire,²⁰ P.Denes,³⁶ F.DeNotaristefani,⁴ A.De Salvo,⁴⁶ M.Diemoz,³⁸ M.Dierckxsen,² C.Dionisi,³⁸ M.Dittmar,⁴⁶ A.Doria,²⁸ M.T.Dova,^{10,11} D.Duchesneau,⁴ M.Duda,¹ B.Echenard,²⁰ A.Eline,¹⁸ A.El Hage,¹ H.El Mamouni,²³ A.Engler,³⁴ F.J.Eppling,¹³ P.Extermann,²⁰ M.A.Falagan,²⁴ S.Falciano,³⁸ A.Favara,³¹ J.Fay,²³ O.Fedin,³³ M.Felcini,⁴⁶ T.Ferguson,³⁴ H.Fesefeldt,¹ E.Fiandrini,³² J.H.Field,²⁰ F.Filthaut,³⁰ P.H.Fisher,¹³ W.Fisher,³⁶ I.Fisk,⁴⁰ G.Forconi,¹³ K.Freudenreich,⁴⁶ C.Furetta,²⁶ Yu.Galaktionov,^{27,13} S.N.Ganguli,⁹ P.Garcia-Abia,²⁴ M.Gataullin,³¹ S.Gentile,³⁸ S.Giagu,³⁸ Z.F.Gong,²¹ G.Grenier,²³ O.Grimm,⁴⁶ M.W.Gruenewald,¹⁶ M.Guida,³⁹ R.van Gulik,² V.K.Gupta,³⁶ A.Gurtu,⁹ L.J.Gutay,⁴³ D.Haas,⁵ R.Sh.Hakobyan,³⁰ J.M.Hansen,¹⁸ D.Hatzifotiadiou,⁸ T.Hebbeker,¹ A.Hervé,¹⁸ J.Hirschfelder,³⁴ H.Hofer,⁴⁶ M.Hohlmann,²⁵ G.Holzner,⁴⁶ S.R.Hou,⁴⁸ Y.Hu,³⁰ B.N.Jin,⁷ L.W.Jones,³ P.de Jong,² I.Josa-Mutuberria,²⁴ D.Käfer,¹ M.Kaur,¹⁴ M.N.Kienzle-Focacci,²⁰ J.K.Kim,⁴² J.Kirkby,¹⁸ W.Kittel,³⁰ A.Klimentov,^{13,27} A.C.König,³⁰ M.Kopal,⁴³ V.Koutsenko,^{13,27} M.Kräber,⁴⁶ R.W.Kraemer,³⁴ A.Krüger,⁴⁵ A.Kunin,¹³ P.Ladron de Guevara,²⁴ I.Laktineh,²³ G.Landi,¹⁷ M.Lebeau,¹⁸ A.Lebedev,¹³ P.Lebrun,²³ P.Lecomte,⁴⁶ P.Lecoq,¹⁸ P.Le Coultre,⁴⁶ J.M.Le Goff,¹⁸ R.Leiste,⁴⁵ M.Levtchenko,²⁶ P.Levtchenko,³³ C.Li,²¹ S.Likhoded,⁴⁵ C.H.Lin,⁴⁸ W.T.Lin,⁴⁸ F.L.Linde,² L.Lista,²⁸ Z.A.Liu,⁷ W.Lohmann,⁴⁵ E.Longo,³⁸ Y.S.Lu,⁷ C.Luci,³⁸ L.Luminari,³⁸ W.Lustermann,⁴⁶ W.G.Ma,²¹ L.Malgeri,²⁰ A.Malinin,²⁷ C.Maña,²⁴ J.Mans,³⁶ J.P.Martin,²³ F.Marzano,³⁸ K.Mazumdar,⁹ R.R.McNeil,⁶ S.Mele,^{18,28} L.Merola,²⁸ M.Meschini,¹⁷ W.J.Metzger,³⁰ A.Mihul,¹¹ H.Milcent,¹⁸ G.Mirabelli,³⁸ J.Mnich,¹ G.B.Mohanty,⁹ G.S.Muanza,²³ A.J.M.Muijs,² B.Musicar,⁴⁰ M.Musy,³⁸ S.Nagy,¹⁵ S.Natale,²⁰ M.Napolitano,²⁸ F.Nessi-Tedaldi,⁴⁶ H.Newman,³¹ A.Nisati,³⁸ H.Nowak,⁴⁵ R.Ofierzynski,⁴⁶ G.Organtini,³⁸ I.Pal,⁴³ C.Palomares,²⁴ P.Paolucci,²⁸ R.Paramatti,³⁸ G.Passaleva,¹⁷ S.Patricelli,²⁸ T.Paul,¹⁰ M.Pauluzzi,³² C.Paus,¹³ F.Pauss,⁴⁶ M.Pedace,³⁸ S.Pensotti,²⁶ D.Perret-Gallix,⁴ B.Petersen,³⁰ D.Piccolo,²⁸ F.Pierella,⁸ M.Pioppi,³² P.A.Piroué,³⁶ E.Pistoiesi,²⁶ V.Plyaskin,²⁷ M.Pohl,²⁰ V.Pojidaev,¹⁷ J.Pothier,¹⁸ D.Prokofiev,³³ J.Quartieri,³⁹ G.Rahal-Callot,⁴⁶ M.A.Rahaman,⁹ P.Raics,¹⁵ N.Raja,⁹ R.Ramelli,⁴⁶ P.G.Rancoita,²⁶ R.Ranieri,¹⁷ A.Raspereza,⁴⁵ P.Razis,²⁹ D.Ren,⁴⁶ M.Rescigno,³⁸ S.Reucroft,¹⁰ S.Riemann,⁴⁵ K.Riles,³ B.P.Roe,³ L.Romero,²⁴ A.Rosca,⁴⁵ S.Rosier-Lees,⁴ S.Roth,¹ C.Rosenbleck,¹ J.A.Rubio,¹⁸ G.Ruggiero,¹⁷ H.Rykaczewski,⁴⁶ A.Sakharov,⁴⁶ S.Saremi,⁶ S.Sarkar,³⁸ J.Salicio,¹⁸ E.Sanchez,²⁴ C.Schäfer,¹⁸ V.Schegelsky,³³ H.Schopper,⁴⁷ D.J.Schotanus,³⁰ C.Sciacca,²⁸ L.Servoli,³² S.Shevchenko,³¹ N.Shivarov,⁴¹ V.Shoutko,¹³ E.Shumilov,²⁷ A.Shvorob,³¹ D.Son,⁴² C.Souga,²³ P.Spillantini,¹⁷ M.Steuer,¹³ D.P.Stickland,³⁶ B.Stoyanov,⁴¹ A.Straessner,¹⁸ K.Sudhakar,⁹ G.Sultanov,⁴¹ L.Z.Sun,²¹ S.Sushkov,¹ H.Suter,⁴⁶ J.D.Swain,¹⁰ Z.Szillasi,^{25,1} X.W.Tang,⁷ P.Tarjan,¹⁵ L.Tauscher,⁵ L.Taylor,¹⁰ B.Tellili,²³ D.Teyssier,²³ C.Timmermans,³⁰ Samuel C.C.Ting,¹³ S.M.Ting,¹³ S.C.Tonwar,⁹ J.Tóth,¹² C.Tully,³⁶ K.L.Tung,⁷ J.Ulbricht,⁴⁶ E.Valente,³⁸ R.T.Van de Walle,³⁰ R.Vasquez,⁴³ V.Veszpremi,²⁵ G.Vesztergombi,¹² I.Vetlitsky,²⁷ D.Vicinanza,³⁹ G.Viertel,⁴⁶ S.Villa,³⁷ M.Vivargent,⁴ S.Vlachos,⁵ I.Vodopianov,²⁵ H.Vogel,³⁴ H.Vogt,⁴⁵ I.Vorobiev,^{34,27} A.A.Vorobyov,³³ M.Wadhwa,⁵ Q.Wang,³⁰ X.L.Wang,²¹ Z.M.Wang,²¹ M.Weber,¹ P.Wienemann,¹ H.Wilkins,³⁰ S.Wynhoff,³⁶ L.Xia,³¹ Z.Z.Xu,²¹ J.Yamamoto,³ B.Z.Yang,²¹ C.G.Yang,⁷ H.J.Yang,³ M.Yang,⁷ S.C.Yeh,⁴⁹ An.Zalite,³³ Yu.Zalite,³³ Z.P.Zhang,²¹ J.Zhao,²¹ G.Y.Zhu,⁷ R.Y.Zhu,³¹ H.L.Zhuang,⁷ A.Zichichi,^{8,18,19} B.Zimmermann,⁴⁶ M.Zöller.¹

- 1 III. Physikalisches Institut, RWTH, D-52056 Aachen, Germany[§]
 - 2 National Institute for High Energy Physics, NIKHEF, and University of Amsterdam, NL-1009 DB Amsterdam, The Netherlands
 - 3 University of Michigan, Ann Arbor, MI 48109, USA
 - 4 Laboratoire d'Annecy-le-Vieux de Physique des Particules, LAPP,IN2P3-CNRS, BP 110, F-74941 Annecy-le-Vieux CEDEX, France
 - 5 Institute of Physics, University of Basel, CH-4056 Basel, Switzerland
 - 6 Louisiana State University, Baton Rouge, LA 70803, USA
 - 7 Institute of High Energy Physics, IHEP, 100039 Beijing, China[△]
 - 8 University of Bologna and INFN-Sezione di Bologna, I-40126 Bologna, Italy
 - 9 Tata Institute of Fundamental Research, Mumbai (Bombay) 400 005, India
 - 10 Northeastern University, Boston, MA 02115, USA
 - 11 Institute of Atomic Physics and University of Bucharest, R-76900 Bucharest, Romania
 - 12 Central Research Institute for Physics of the Hungarian Academy of Sciences, H-1525 Budapest 114, Hungary[‡]
 - 13 Massachusetts Institute of Technology, Cambridge, MA 02139, USA
 - 14 Panjab University, Chandigarh 160 014, India.
 - 15 KLTE-ATOMKI, H-4010 Debrecen, Hungary[¶]
 - 16 Department of Experimental Physics, University College Dublin, Belfield, Dublin 4, Ireland
 - 17 INFN Sezione di Firenze and University of Florence, I-50125 Florence, Italy
 - 18 European Laboratory for Particle Physics, CERN, CH-1211 Geneva 23, Switzerland
 - 19 World Laboratory, FBLJA Project, CH-1211 Geneva 23, Switzerland
 - 20 University of Geneva, CH-1211 Geneva 4, Switzerland
 - 21 Chinese University of Science and Technology, USTC, Hefei, Anhui 230 029, China[△]
 - 22 University of Lausanne, CH-1015 Lausanne, Switzerland
 - 23 Institut de Physique Nucléaire de Lyon, IN2P3-CNRS, Université Claude Bernard, F-69622 Villeurbanne, France
 - 24 Centro de Investigaciones Energéticas, Medioambientales y Tecnológicas, CIEMAT, E-28040 Madrid, Spain[‡]
 - 25 Florida Institute of Technology, Melbourne, FL 32901, USA
 - 26 INFN-Sezione di Milano, I-20133 Milan, Italy
 - 27 Institute of Theoretical and Experimental Physics, ITEP, Moscow, Russia
 - 28 INFN-Sezione di Napoli and University of Naples, I-80125 Naples, Italy
 - 29 Department of Physics, University of Cyprus, Nicosia, Cyprus
 - 30 University of Nijmegen and NIKHEF, NL-6525 ED Nijmegen, The Netherlands
 - 31 California Institute of Technology, Pasadena, CA 91125, USA
 - 32 INFN-Sezione di Perugia and Università Degli Studi di Perugia, I-06100 Perugia, Italy
 - 33 Nuclear Physics Institute, St. Petersburg, Russia
 - 34 Carnegie Mellon University, Pittsburgh, PA 15213, USA
 - 35 INFN-Sezione di Napoli and University of Potenza, I-85100 Potenza, Italy
 - 36 Princeton University, Princeton, NJ 08544, USA
 - 37 University of California, Riverside, CA 92521, USA
 - 38 INFN-Sezione di Roma and University of Rome, “La Sapienza”, I-00185 Rome, Italy
 - 39 University and INFN, Salerno, I-84100 Salerno, Italy
 - 40 University of California, San Diego, CA 92093, USA
 - 41 Bulgarian Academy of Sciences, Central Lab. of Mechatronics and Instrumentation, BU-1113 Sofia, Bulgaria
 - 42 The Center for High Energy Physics, Kyungpook National University, 702-701 Taegu, Republic of Korea
 - 43 Purdue University, West Lafayette, IN 47907, USA
 - 44 Paul Scherrer Institut, PSI, CH-5232 Villigen, Switzerland
 - 45 DESY, D-15738 Zeuthen, Germany
 - 46 Eidgenössische Technische Hochschule, ETH Zürich, CH-8093 Zürich, Switzerland
 - 47 University of Hamburg, D-22761 Hamburg, Germany
 - 48 National Central University, Chung-Li, Taiwan, China
 - 49 Department of Physics, National Tsing Hua University, Taiwan, China
- [§] Supported by the German Bundesministerium für Bildung, Wissenschaft, Forschung und Technologie
[‡] Supported by the Hungarian OTKA fund under contract numbers T019181, F023259 and T037350.
[¶] Also supported by the Hungarian OTKA fund under contract number T026178.
[‡] Supported also by the Comisión Interministerial de Ciencia y Tecnología.
[‡] Also supported by CONICET and Universidad Nacional de La Plata, CC 67, 1900 La Plata, Argentina.
[△] Supported by the National Natural Science Foundation of China.

Decay Channel	Branching Ratios			
	$M = 102 \text{ GeV}$		$M = 200 \text{ GeV}$	
	$f = f'$	$f = -f'$	$f = f'$	$f = -f'$
$\ell^* \rightarrow \ell\gamma$	70%	–	35%	–
$\ell^* \rightarrow \nu W$	28%	83%	55%	63%
$\ell^* \rightarrow \ell Z$	2%	17%	10%	37%
$\nu^* \rightarrow \nu\gamma$	–	70%	–	35%
$\nu^* \rightarrow \ell W$	83%	28%	63%	55%
$\nu^* \rightarrow \nu Z$	17%	2%	37%	10%

Table 1: Predicted branching ratios for charged and neutral excited lepton decays, for different choices of masses (M) and couplings.

Decay mode	Pair production		Single production	
	$e^+e^- \rightarrow \ell^*\ell^*$	$e^+e^- \rightarrow \nu^*\nu^*$	$e^+e^- \rightarrow \ell\ell^*$	$e^+e^- \rightarrow \nu\nu^*$
Radiative	$\ell^*\ell^* \rightarrow \ell\ell\gamma\gamma$	$\nu^*\nu^* \rightarrow \nu\nu\gamma\gamma$	$\ell\ell^* \rightarrow \ell\ell\gamma$	$\nu\nu^* \rightarrow \nu\nu\gamma$
Charged-current	$\ell^*\ell^* \rightarrow \nu\nu WW$	$\nu_\ell^*\nu_\ell^* \rightarrow \ell\ell WW$	$\nu_\ell\nu_\ell^* \rightarrow \nu_\ell\ell W$	$\ell\ell^* \rightarrow \ell\nu_\ell W$
Neutral-current	–	–	$\ell\ell^* \rightarrow \ell\ell Z$	$\nu\nu^* \rightarrow \nu\nu Z$
Mixed	$\ell^*\ell^* \rightarrow \ell\gamma\nu W$	$\nu_\ell^*\nu_\ell^* \rightarrow \nu\gamma\ell W$	–	–

Table 2: The final states for excited lepton production considered in this Letter, where ℓ runs on the lepton flavour: e , μ and τ .

Pair production				
Signal	Selection	N_D	N_B	ε (%)
$e^*e^* \rightarrow ee\gamma\gamma$	Leptons	0	0.1	45
$\mu^*\mu^* \rightarrow \mu\mu\gamma\gamma$	and	0	0.3	44
$\tau^*\tau^* \rightarrow \tau\tau\gamma\gamma$	photons	0	0.4	40
$\ell^*\ell^* \rightarrow \nu\nu WW$	qqqq or qq $\ell\nu$	3012	2974	66
$e^*e^* \rightarrow e\gamma\nu W$	$e + \gamma + \cancel{E} + [\text{qq or } \ell]$	9	10	56
$\mu^*\mu^* \rightarrow \mu\gamma\nu W$	$\mu + \gamma + \cancel{E} + [\text{qq or } \ell]$	6	5	49
$\tau^*\tau^* \rightarrow \tau\gamma\nu W$	jet + $\gamma + \cancel{E} + [\text{qq or } \ell]$	45	43	35
$\nu_e^*\nu_e^* \rightarrow eeWW$	$ee + [\text{qqqq or qq}\ell\nu]$	1	0.2	15
$\nu_\mu^*\nu_\mu^* \rightarrow \mu\mu WW$	$\mu\mu + [\text{qqqq or qq}\ell\nu]$	2	0.6	19
$\nu_\tau^*\nu_\tau^* \rightarrow \tau\tau WW$	qqqq or qq $\ell\nu$	3012	2974	70
$\nu^*\nu^* \rightarrow \nu\nu\gamma\gamma$	$\gamma\gamma + \cancel{E}$	2	1.9	45
$\nu_e^*\nu_e^* \rightarrow \nu\gamma eW$	$e + \gamma + \cancel{E} + [\text{qq or } \ell]$			35
$\nu_\mu^*\nu_\mu^* \rightarrow \nu\gamma\mu W$	$\mu + \gamma + \cancel{E} + [\text{qq or } \ell]$	10	11	29
$\nu_\tau^*\nu_\tau^* \rightarrow \nu\gamma\tau W$	jet + $\gamma + \cancel{E} + [\text{qq or } \ell]$			23

Table 3: Selections used in the search for pair-produced excited leptons and the corresponding numbers of observed events, N_D , expected background, N_B , and selection efficiency, ε , for an excited lepton mass of 101 GeV.

Single Production				
Signal	Selection	N_D	N_B	ε (%)
$ee^* \rightarrow ee\gamma$	Leptons and a photon	672	737	54, 65, 65
$\mu\mu^* \rightarrow \mu\mu\gamma$		55	66	62, 63, 60
$\tau\tau^* \rightarrow \tau\tau\gamma$		48	60	42, 45, 43
$ee^* \rightarrow e\nu_e W$	qq $l\nu$ or	749	773	65, 64, 61
$\mu\mu^* \rightarrow \mu\nu_\mu W$	qq \cancel{E} or	652	675	60, 64, 62
$\tau\tau^* \rightarrow \tau\nu_\tau W$	$ll\cancel{E}$	1287	1307	59, 48, 50
$ee^* \rightarrow eeZ$	qq $l\nu$ or	673	698	65, 37, 63
$\mu\mu^* \rightarrow \mu\mu Z$	qq \cancel{E} or	568	596	58, 41, 59
$\tau\tau^* \rightarrow \tau\tau Z$	$ll\cancel{E}$	1217	1236	32, 22, 37
$\nu\nu^* \rightarrow \nu\nu\gamma$	$\gamma + \cancel{E}$	176	194	60, 67, 70
$\nu_e\nu_e^* \rightarrow \nu_e e W$	qq $l\nu$ or	749	773	61, 68, 67
$\nu_\mu\nu_\mu^* \rightarrow \nu_\mu\mu W$	qq \cancel{E} or	652	675	63, 63, 60
$\nu_\tau\nu_\tau^* \rightarrow \nu_\tau\tau W$	$ll\cancel{E}$	1287	1307	46, 57, 60
$\nu\nu^* \rightarrow \nu\nu Z$	qq \cancel{E} or $ll\cancel{E}$	343	350	18, 15, 40

Table 4: Selections used in the search for singly-produced excited leptons and corresponding numbers of observed events, N_D , expected background, N_B , and selection efficiency, ε , for excited lepton masses of 110, 160 and 201 GeV.

Excited Lepton	95% CL Mass Limit (GeV)		
	$f = f'$	$f = -f'$	Any Coupling
e^*	102.8	96.6	96.5
μ^*	102.8	96.6	96.6
τ^*	102.8	96.6	95.6
ν_e^*	101.7	102.6	101.5
ν_μ^*	101.8	102.6	101.4
ν_τ^*	92.9	102.6	91.3

Table 5: Lower mass limits at 95% confidence level for charged and neutral excited lepton as obtained from pair-production searches.




Figure 1: Distributions of a) the hadronic invariant mass, M_{jj} , in the $qq\ell\nu$ selection; b) the sum of M_{jj} and the recoil mass, M_{rec} , for two jets in the $qqqq$ selection; c) the electron photon invariant mass in the $e\gamma\nu W$ selection and d) the muon photon invariant mass in the $\mu\gamma\nu W$ selection. The expected signal for excited leptons produced in pairs with a mass of 101 GeV is shown together with data and Standard Model background.




Figure 2: The invariant mass distributions for the a) $e-\gamma$, b) $\mu-\gamma$, and c) $\tau-\gamma$ systems. d) The normalised energy distribution of single photon events. The expected signal for a singly-produced excited lepton with a mass of 160 GeV is shown together with data and the Standard Model background expectation. The signals are plotted for the arbitrary choice of couplings displayed in the Figures.




Figure 3: Lepton recoil mass distributions for the a) $qqe\nu$, b) $qq\mu\nu$ and c) $qq\tau\nu$ selections. Invariant mass distributions for the d) $qqe\nu$, e) $qq\mu\nu$ and f) $qq\tau\nu$ selections. The expected signal for a singly-produced excited lepton with a mass of 160 GeV is shown together with data and Standard Model background expectations. The signals are plotted for the arbitrary choice of couplings displayed in the Figures.




Figure 4: 95% confidence level upper limits on $|f|/\Lambda$, as a function of the excited lepton mass with $f = f'$ for a) e^* , μ^* and τ^* , b) ν_e^* , ν_μ^* and ν_τ^* , and with $f = -f'$ for c) e^* , μ^* and τ^* , d) ν_e^* , ν_μ^* and ν_τ^* .