

Status of the NA57 experiment at CERN PS

F. Antinori, A. Badala, R. Barbera, I.J. Bloodworth, G.E. Bruno, R. Caliandro, M. Campbell, E. Cantatore, W. Carena, N. Carrer, et al.

► **To cite this version:**

F. Antinori, A. Badala, R. Barbera, I.J. Bloodworth, G.E. Bruno, et al.. Status of the NA57 experiment at CERN PS. *Journal of Physics G: Nuclear and Particle Physics*, IOP Publishing, 2001, 27, pp.383-390. in2p3-00019670

HAL Id: in2p3-00019670

<http://hal.in2p3.fr/in2p3-00019670>

Submitted on 5 Sep 2001

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Status of the NA57 experiment at CERN SPS

Presented by V Manzari for the NA57 Collaboration:

F Antinori¹, A Badalà², R Barbera², I J Bloodworth³, G E Bruno⁴,
R Caliendo⁴, M Campbell⁵, E Cantatore⁵, W Carena⁵, N Carrer¹,
A P de Haas⁶, D Di Bari⁴, S Di Liberto⁷, R Divia⁵, D Elia⁴, D Evans³,
K Fanebust⁸, F Fayazzadeh⁹, J Fedorisin¹⁰, G A Feofilov¹¹, R A Fini⁴,
J Ftáčnik¹², B Ghidini⁴, G Grella¹³, M Gulino², H Helstrup¹⁴,
M Henriquez⁹, A K Holme⁹, A Jacholkowski⁴, G T Jones³, P Jovanovic¹²,
A Jusko¹⁰, R Kamermans⁶, J B Kinson³, W Klempt⁵, K Knudson⁵,
A A Kolojvari¹¹, V Kondratiev¹¹, I Králik¹⁰, P Kuijer⁶, V Lenti⁴,
R Lietava¹², G Løvholden⁹, M Lupták¹⁰, V Manzari⁴, G Martinska¹⁰,
M A Mazzoni⁷, F Meddi⁷, A Michalon¹⁵, M E Michalon-Mentzer¹⁵,
M Morando¹, D Muigg⁶, E Nappi⁴, F Navach⁴, P I Norman³, A Palmeri²,
G S Pappalardo², B Pastirčák¹⁰, J Pisut¹², N Pisutova¹², F Posa⁴,
E Quercigh⁵, F Riggi², D Röhrich⁸, G Romano¹³, K Šafařík⁵, L Šándor¹⁰,
E Schillings⁶, G Segato¹, M Senè¹⁶, R Senè¹⁶, W Snoeys⁵, P Staroba¹⁷,
M Thompson³, T A Tulina¹¹, R Turrisi¹, T S Tveter⁹, J Urbán¹⁰,
F F Valiev¹¹, A van den Brink⁶, P van de Ven⁶, P Vande Vyvre⁵,
N van Eijndhoven⁶, L Vannucci¹⁸, A Vascotto⁵, T Vik⁹,
O Villalobos Baillie³, I Vinogradov¹¹, T Virgili¹³, M F Votruba³,
J Vrlakova¹⁰ and P Závada¹⁷

¹ University of Padua and INFN, Padua, Italy

² University of Catania and INFN, Catania, Italy

³ University of Birmingham, Birmingham, UK

⁴ Dipartimento IA di Fisica dell'Università e del Politecnico di Bari and INFN, Bari, Italy

⁵ CERN, European Laboratory for Particle Physics, Geneva, Switzerland

⁶ Utrecht University and NIKHEF, Utrecht, The Netherlands

⁷ University 'La Sapienza' and INFN, Rome, Italy

⁸ Fysisk Institutt, Universitetet i Bergen, Bergen, Norway

⁹ University of Oslo, Oslo, Norway

¹⁰ Institute of Experimental Physics SAS and Šafařík University, Kosice, Slovakia

¹¹ State University of St Petersburg, St Petersburg, Russia

¹² Comenius University, Bratislava, Slovakia

¹³ Dipartimento di Scienze Fisiche 'E R Caianiello' dell'Università and INFN, Salerno, Italy

¹⁴ Høgskolen i Bergen, Bergen, Norway

¹⁵ IReS/ULP, Strasbourg, France

¹⁶ Collège de France and IN2P3, Paris, France

¹⁷ Institute of Physics, Prague, Czech Republic

¹⁸ INFN, Laboratori Nazionali di Legnaro, Legnaro, Italy

Received 2 November 2000

Abstract

The NA57 experiment aims to investigate the production of strange and multi-strange particles in nucleus–nucleus collisions at CERN SPS. NA57 has been specifically designed to extend the study of Λ , Ξ and Ω as well as the K_s^0 (and

their antiparticles) production pattern in ultra-relativistic nuclear collisions, as a function of the centre-of-mass energy and in a wider centrality range than its predecessor WA97. In this paper we recall the main features of the NA57 set-up and we report on the status of the processing and analysis of data samples collected in 1998 with a lead beam at $158 A \text{ GeV } c^{-1}$ and in 1999 with lead and proton beams at $40 A \text{ GeV } c^{-1}$.

(Some figures in this article are in colour only in the electronic version; see www.iop.org)

1. Introduction

The WA97 experiment [1] has shown that the multi-strange baryon and antibaryon yields per wounded nucleon at central rapidity are enhanced when going from p–Be to Pb–Pb collisions. The enhancement increases with the strange content of the produced particles. Such an effect has been predicted [2] as a characteristic signature of the phase transition to a quark–gluon plasma, and has so far not been reproduced by microscopic hadronic collision models. The NA57 experiment has been designed to study the strange and multi-strange particle production in nucleus–nucleus collisions as a function of the centre-of-mass energy and collision centrality [3]. Its main physics goal is to search for the onset of the observed strangeness enhancement by:

- extending the centrality range of WA97 for Pb–Pb collisions at $158 A \text{ GeV } c^{-1}$
- measuring the pattern of multi-strange hyperon production in Pb–Pb collisions at lower values of the energy density.

2. The NA57 set-up

The NA57 set-up is designed to measure the decays of strange baryons and antibaryons in the high-multiplicity environment of the Pb–Pb collisions at the SPS. The full apparatus is shown schematically in figure 1: a detailed description can be found elsewhere [4].

We recall here that the heart of the apparatus is a telescope tracker made of silicon pixel detectors, placed inside a 1.4 T magnetic field. Strange and multi-strange particles are identified by reconstructing their weak decays into final states containing only charged particles. The telescope is inclined with respect to the beam line and points back to the target: the inclination angle can be changed to select different laboratory rapidity settings. The full acceptance coverage corresponds to about one unit of rapidity at medium transverse momentum and it is the same for particles and antiparticles.

The centrality trigger for Pb–Pb collisions, based on a scintillator petal system placed 10 cm downstream of the target, selects the most central 60% of the total inelastic cross section (to be compared with 40% in WA97). The centrality of the collision is measured via the charged particle multiplicity, sampled at central rapidity by two multiplicity strip detector (MSD) stations, as in WA97.

3. Data samples

After a commissioning proton run in 1997, the following samples of data for physics analysis have been collected by NA57 in 1998 and 1999:




Figure 2. Wounded nucleon distributions for the five NA57 multiplicity classes.




Figure 3. Scatter plot of ΛK versus $\Lambda\pi$ invariant mass.




Figure 4. $\Lambda\pi^-$ and $\bar{\Lambda}\pi^+$ invariant mass distributions.

Consequently, the detector alignment procedure and the track finding and event reconstruction software packages had to be significantly changed to match the NA57 specific new features.




Figure 5. ΔK^- and $\bar{\Lambda} K^+$ invariant mass distributions.

5. Signals from the 1998 data sample

5.1. Centrality range

The study of the collision centrality has been improved with respect to the WA97 data [6]. In the NA57 experiment a precise measurement of the trigger cross section allowed us to estimate




Figure 6. Acceptance windows for Ξ and Ω .

the number of wounded nucleons without assumptions on its scaling with the measured charged particle multiplicity.

The NA57 sample of events has been divided into five multiplicity classes: the corresponding distributions of the number of wounded nucleons are shown in figure 2.

The most peripheral events in the class 0 correspond to the new region to be investigated with respect to the WA97 studies.

5.2. Mass spectra

Figure 3 shows the correlation of the $\Lambda\pi$ and ΛK invariant masses, after preliminary analysis cuts have been applied: clear signals from $\Xi \rightarrow \Lambda\pi$ and $\Omega \rightarrow \Lambda K$ particle decays are visible.

Figures 4 and 5 show the mass spectra in 2 MeV bins for $\Lambda\pi^-$ and ΛK^- and charge-conjugate channels separately. For the Ω^- and $\bar{\Omega}^+$ signals an additional cut is applied to reject the background due to Ξ decays kinematically ambiguous with Ω decays.

The distributions peak at the nominal Ξ and Ω masses, with FWHM of about 10 and 12 MeV, respectively.

The $y_{Lab}-p_T$ distributions for the selected Ξ s and Ω s are shown in figure 6. A weighting procedure, based on a Monte Carlo simulation of the full apparatus, will then be applied to each selected particle to account for losses due to geometrical acceptance, detection and reconstruction efficiencies.

6. Summary and future plans

The status of the NA57 analysis has been reported. First results are shown for the centrality analysis, together with the first Ξ and Ω signals from the 1998 Pb–Pb run at $158 A \text{ GeV } c^{-1}$. Low-energy p–Be and Pb–Pb data from 1999 runs at $40 A \text{ GeV } c^{-1}$ beam momentum are being processed.

The year 2000 run should provide a substantial increase of the statistics for the Pb–Pb data at $158 \text{ GeV } c^{-1}$. In the year 2001 we plan a $40 A \text{ GeV } c^{-1}$ proton run in order to increase the present p–Be statistics.

References

- [1] Fini R A *et al* (WA97 Collaboration) 2001 *J. Phys. G: Nucl. Part. Phys.* **27** 375
- [2] Rafelski J and Müller B 1982 *Phys. Rev. Lett.* **48** 1066
- [3] Caliendo R *et al* (NA57 Collaboration) 1996 *Tech. Proposal CERN/SPSLC/96-40*, SPSLC/P300
- [4] Manzari V *et al* (NA57 Collaboration) 1999 *J. Phys. G: Nucl. Part. Phys.* **25** 473
- [5] Vande Vyvre P *et al* (CERN ALICE DAQ group) 1999 *ALICE Internal Note/DAQ 99/46*
- [6] Carrer N *et al* (WA97 and NA57 Collaborations) 2001 *J. Phys. G: Nucl. Part. Phys.* **27** 391