



Test of CP Invariance in $Z \rightarrow \mu^+ \mu^- \gamma$ decay

M. Acciarri, O. Adriani, M. Aguilar-Benitez, S. Ahlen, J. Alcaraz, G. Alemanni, A. Aloisio, M G. Alviggi, G. Ambrosi, H. Anderhub, et al.

► To cite this version:

M. Acciarri, O. Adriani, M. Aguilar-Benitez, S. Ahlen, J. Alcaraz, et al.. Test of CP Invariance in $Z \rightarrow \mu^+ \mu^- \gamma$ decay. Physics Letters B, 1998, 436, pp.428-436. 10.1016/S0370-2693(98)00965-4 . in2p3-00023167

HAL Id: in2p3-00023167

<https://hal.in2p3.fr/in2p3-00023167>

Submitted on 6 Nov 1998

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Test of CP Invariance in $Z \rightarrow \mu^+ \mu^- \gamma$ decay

The L3 Collaboration

Abstract

We report on the first test of CP invariance in Z decays with hard photon radiation. The data recorded with the L3 detector at centre-of-mass energies near the Z resonance are used to search for CP violation in the reaction $e^+e^- \rightarrow \mu^+\mu^-\gamma$. No evidence for CP violation is found and limits on the CP-violating form factors f_A^μ and f_V^μ are derived.

Submitted to *Phys. Lett. B*

Introduction

Violation of CP invariance in Z decays would point to new physics beyond the Standard Model [1]. In the Standard Model the CP-violating contributions to the Z decay amplitude are of the order of 10^{-7} at most [2]. They are therefore not measureable at current experiments. At LEP experimental tests of the CP symmetry have been performed analysing the decays $Z \rightarrow \tau^+\tau^-$ [3] and $Z \rightarrow b\bar{b}g$ [4]. Recently the authors of Reference [5] proposed to test the CP invariance of the radiative decays $Z \rightarrow \mu^+\mu^-\gamma$ and $Z \rightarrow \tau^+\tau^-\gamma$. The angular correlations of the three particle final state allow a measurement of chirality-conserving 4-particle couplings, which could be induced by CP-violating interactions beyond the Standard Model [6].

CP violation in the decays $Z \rightarrow \mu^+\mu^-\gamma$ and $Z \rightarrow \tau^+\tau^-\gamma$ can be described using the following effective Lagrangian

$$\mathcal{L}(x) = \sum_{\ell} \left\{ \begin{aligned} & [f_V^{\ell} \bar{\ell}(x) \gamma^{\nu} \ell(x) + f_A^{\ell} \bar{\ell}(x) \gamma^{\nu} \gamma_5 \ell(x)] Z^{\mu}(x) [\partial_{\mu} A_{\nu}(x) - \partial_{\nu} A_{\mu}(x)] \\ & - \frac{i}{2} d_{\gamma}^{\ell} \bar{\ell}(x) \sigma^{\mu\nu} \gamma_5 \ell(x) [\partial_{\mu} A_{\nu}(x) - \partial_{\nu} A_{\mu}(x)] \\ & - \frac{i}{2} d_Z^{\ell} \bar{\ell}(x) \sigma^{\mu\nu} \gamma_5 \ell(x) [\partial_{\mu} Z_{\nu}(x) - \partial_{\nu} Z_{\mu}(x)] \end{aligned} \right\}, \quad (1)$$

where d_{γ}^{ℓ} and d_Z^{ℓ} are the electric and weak dipole moments of the lepton ($\ell = \mu, \tau$), respectively. The real coupling constants f_V^{ℓ} and f_A^{ℓ} correspond to a CP-violating, but chirality-conserving $Z\ell\ell\gamma$ vertex.

The electric dipole moment of the muon is known to be smaller than 10^{-18} ecm [7], a limit which cannot be improved by using LEP data. For the weak dipole moment only indirect limits exist [5], which use the partial width of $Z \rightarrow \mu^+\mu^-(\gamma)$. It is found to be smaller than 10^{-17} ecm. For the tau lepton a better limit [8] has been obtained from LEP data by analysing the topology of tau decays. This kind of analysis cannot be performed using muons, as the spin-analysing muon decay occurs outside the detector.

Therefore, we concentrate on the measurement of the real coupling constants f_V^{μ} and f_A^{μ} , which are of mass dimension -2 and parametrise an anomalous, CP-violating, but chirality-conserving $Z\mu\mu\gamma$ vertex (see Figure 1). Following Reference [2] we introduce the dimensionless parameters $\hat{f}_V^{\mu}$ and $\hat{f}_A^{\mu}$ defined by

$$f_{V/A}^{\mu} = \frac{e^2}{\sin \theta_w \cos \theta_w m_Z^2} \hat{f}_{V/A}^{\mu} = 2.62 \cdot 10^{-5} \text{ GeV}^{-2} \hat{f}_{V/A}^{\mu}. \quad (2)$$

We measure these parameters from the rate of $\mu^+\mu^-\gamma$ final states and by investigating CP-odd observables in our $e^+e^- \rightarrow \mu^+\mu^-\gamma$ sample obtained at centre-of-mass energies around the Z resonance.

Event Selection and Data Sample

The L3 detector is described elsewhere [9]. The selection of muon pair events requires either two reconstructed tracks in the muon spectrometer or one muon track and the signature of a minimum ionising particle in the inner sub-detectors. Details of the muon identification are presented in Reference [10].

The two muons should be in the polar angle region $|\cos\theta| < 0.8$. The sum of the energies of the two muons and all photons in the event must exceed 80% of the centre-of-mass energy. In order to reject cosmic ray muons at least one scintillator hit is required with a time measurement that coincides within $\pm 3\text{ns}$ with the beam crossing. Additionally at least one track in the vertex chamber is required with a distance of closest approach to the beam axis of less than 5 mm. We apply no acollinearity or acoplanarity cut.

The data were taken in the years from 1991 to 1995, when LEP was operated at centre-of-mass energies in the vicinity of the Z resonance. From these data, corresponding to a total integrated luminosity of 133 pb^{-1} , we select 92072 muon-pair events.

Analysis of the Event Topologies

Energy and momentum conservation implies that the kinematics of the $\mu^+\mu^-\gamma$ final state is fully determined by measuring the directions of the three particles. In the following we denote the direction of the muons by the unit vectors $\hat{\mathbf{k}}_{\mu^+}$ and $\hat{\mathbf{k}}_{\mu^-}$. For the search for CP violation the following CP-odd observables were found to be useful [5]

$$\begin{aligned} T &= (\hat{\mathbf{k}}_{\mu^+} - \hat{\mathbf{k}}_{\mu^-}) \cdot \hat{\mathbf{p}}_{e^+} (\hat{\mathbf{k}}_{\mu^+} \times \hat{\mathbf{k}}_{\mu^-}) \cdot \hat{\mathbf{p}}_{e^+} , \\ V &= (\hat{\mathbf{k}}_{\mu^+} \times \hat{\mathbf{k}}_{\mu^-}) \cdot \hat{\mathbf{p}}_{e^+} , \end{aligned} \quad (3)$$

where $\hat{\mathbf{p}}_{e^+}$ is the direction of the incoming positron. In this context T is the component of a tensor and V of a vector observable.

Figure 2 shows the distributions of the observables T and V for the data and events generated with the KORALZ [11] Monte-Carlo program and including full detector simulation using the GEANT [13] detector simulation program. KORALZ simulates the radiation of up to two hard photons in the initial or final state and includes photon exponentiation according to the Yennie-Frautschi-Suura ansatz [12]. Good agreement is observed over the full ranges of the two variables. Above all no asymmetry between negative and positive values, *i.e.* no sign of CP violation, is observed.

To compare the data with the prediction from CP-violating $Z\mu\mu\gamma$ couplings we use a Monte-Carlo program [14]. For the calculation of the observables T and V , it is convenient to replace the parameters $\hat{f}_V^\mu$ and $\hat{f}_A^\mu$ by the linear combinations

$$\begin{aligned} \hat{f}_1^\mu &= g_V^\mu \hat{f}_A^\mu - g_A^\mu \hat{f}_V^\mu , \\ \hat{f}_2^\mu &= g_V^\mu \hat{f}_V^\mu - g_A^\mu \hat{f}_A^\mu , \end{aligned} \quad (4)$$

where $g_V^\mu = 2\sin^2\theta_w - \frac{1}{2}$ and $g_A^\mu = -\frac{1}{2}$ are the neutral current coupling constants of the muon. As $\hat{f}_1^\mu$ influences only tensor observables and $\hat{f}_2^\mu$ is related to vector observables, these two parameters can be measured independently by using either the observable T or V . The sensitivity of these observables on the electric and weak dipole moment of the muon is two orders of magnitude smaller [5] and therefore neglected.

In Figure 3 the relative asymmetries between negative and positive values of the observables T and V are shown for intervals in $|T|$ and $|V|$, respectively. Within their statistical errors all measurements are consistent with zero. Also included are the theoretical expectations for $\hat{f}_1^\mu = \pm 1$ and $\hat{f}_2^\mu = \pm 1$, which corresponds to a CP-violating coupling on the order of the Fermi constant. The expected effect of the anomalous CP-violating couplings is larger for the events with large values of $|T|$ or $|V|$. The events at $|T| = |V| = 0$ are those without photon radiation,

$Z \rightarrow \mu^+ \mu^-$. In this two-particle final state it is impossible to measure CP violation without information on the muon spin. But these events can be used to test the CP invariance of the detector and of the acceptance cuts. From Figure 3 we conclude that such effects that could mimic CP violation are not present in the data.

To extract limits on the parameters $\hat{f}_1^\mu$ and $\hat{f}_2^\mu$, two sub-samples with $|T| > 0.025$ and $|V| > 0.1$ are considered, containing in total 3680 and 1586 events, respectively. To predict the number of events in each $|T|$ or $|V|$ bin according to the Standard Model, N_i^{SM} , we use the events that were generated with the KORALZ program and include the full simulation of the L3 detector. The Monte-Carlo program [14] is used to calculate the expected change of the number of events in the different bins, $N_i(\hat{f}^\mu)/N_i(0)$, and to predict the relative asymmetry, $A_i(\hat{f}^\mu)$, which is shown in Figure 3. With this procedure the detector effects and radiative corrections calculated with the KORALZ Monte Carlo are taken into account.

The difference between the number of events with positive T or V and negative T or V in bin number i is predicted as

$$N_i^{\text{SM}} \cdot \frac{N_i(\hat{f}^\mu)}{N_i(0)} \cdot A_i(\hat{f}^\mu) .$$

The calculated asymmetry is shown in Figure 4 for data and theory. It is proportional to the interference term between the Standard Model amplitude and the anomalous CP-odd amplitude. Varying the anomalous couplings not only changes the asymmetry, but also the total number of radiative events in the different bins, which is proportional to the absolute square of the CP-odd amplitude and therefore CP even. The number of events in bin number i is calculated as

$$N_i^{\text{SM}} \cdot \frac{N_i(\hat{f}^\mu)}{N_i(0)} .$$

We compare our data and the predictions for different values of $\hat{f}_1^\mu$ and $\hat{f}_2^\mu$, using a χ^2 fit for different ranges of T and V , as indicated by the arrow in Figure 4. Using the asymmetries the parameters are measured to be

$$\hat{f}_1^\mu = 0.08 \pm 0.64 \quad \text{and} \quad \hat{f}_2^\mu = -1.61 \pm 1.34 .$$

Exploiting in addition the spectra of radiative events improves the measurement. We then get

$$\hat{f}_1^\mu = 0.03 \pm 0.19 \quad \text{and} \quad \hat{f}_2^\mu = -0.12_{-0.21}^{+0.34} ,$$

both consistent with zero.

In our event sample the average invariant mass of the muon photon pair, $m_{\mu\gamma}$, and therefore the mass of the radiating virtual muon, is about 15 GeV. The ratio between the CP-violating Z decay amplitude and the Standard Model amplitude coming from bremsstrahlung amounts to $\hat{f}^\mu (m_{\mu\gamma}/m_Z)^2$. Our measurement constrains this ratio to values below 10^{-2} at the 95% confidence level.

For the CP-violating coupling constants f_V^μ and f_A^μ we get limits of

$$\begin{aligned} -1.7 \cdot 10^{-5} \text{ GeV}^{-2} &< f_V^\mu < 1.9 \cdot 10^{-5} \text{ GeV}^{-2} & \text{and} \\ -2.4 \cdot 10^{-5} \text{ GeV}^{-2} &< f_A^\mu < 2.2 \cdot 10^{-5} \text{ GeV}^{-2} \end{aligned}$$

at the 95% confidence level.

Following Reference [2] the measurements of f_V^μ and f_A^μ are used to set a lower limit on the scale parameter for CP-violating interactions, Λ_{CP} , of 229 GeV and 204 GeV, respectively.

Acknowledgements

We wish to express our gratitude to the CERN accelerator divisions for the excellent performance of the LEP machine. We acknowledge the contributions of all the engineers and technicians who have participated in the construction and maintenance of this experiment. We would like to thank W. Bernreuther, O. Nachtmann and P. Overmann for fruitful discussions.

References

- [1] S.L. Glashow, Nucl. Phys. **22** (1961) 579;
S. Weinberg, Phys. Rev. Lett. **19** (1967) 1264;
A. Salam, “Elementary Particle Theory”, Ed. N. Svartholm, “Almqvist and Wiksell” (1968), 367.
- [2] W. Bernreuther, U. Löw, J.P. Ma, O. Nachtmann, Z. Phys. **C 43** (1989) 117.
- [3] ALEPH Collab., D. Buskulic *et al.*, Phys. Lett. **B 346** (1995) 371;
OPAL Collab., K. Ackerstaff *et al.*, Z. Phys. **C 74** (1997) 403.
- [4] ALEPH Collab., D. Buskulic *et al.*, Phys. Lett. **B 384** (1996) 365.
- [5] D. Bruß, O. Nachtmann, P. Overmann, Eur. Phys. J. **C 1** (1998) 191.
- [6] W. Bernreuther, A. Brandenburg, P. Haberl, O. Nachtmann, Phys. Lett. **B 387** (1996) 155.
- [7] Particle Data Group, Phys. Rev. **D 54** (1996) 21.
- [8] OPAL Collab., K. Ackerstaff *et al.*, Z. Phys. **C 74** (1997) 403;
L3 Collab., M. Acciarri *et al.*, Phys. Lett. **B 426** (1998) 207.
- [9] L3 Collab., B. Adeva *et al.*, Nucl. Inst. Meth. **A 289** (1990) 35;
M. Chemarin *et al.*, Nucl. Inst. Meth. **A 349** (1994) 345;
M. Acciarri *et al.*, Nucl. Inst. Meth. **A 351** (1994) 300;
G. Basti *et al.*, Nucl. Inst. Meth. **A 374** (1996) 293;
I. Brock *et al.*, Nucl. Inst. Meth. **A 381** (1996) 236;
A. Adam *et al.*, Nucl. Inst. Meth. **A 383** (1996) 342.
- [10] L3 Collab., M. Acciarri *et al.*, Z. Phys. **C 62** (1994) 551.
- [11] S. Jadach, B.F.L. Ward, Z. Wąs, Comp. Phys. Comm. **79** (1994) 503.
- [12] B.F.L. Ward *et al.*, Acta Phys. Polon. **B 25** (1994) 245.
- [13] R. Brun *et al.*, “GEANT 3”, CERN DD/EE/84-1 Revised (1987).
- [14] The Monte-Carlo program is provided by the authors of Reference [5];
P. Overmann, private communication.

The L3 Collaboration:

M. Acciarri,²⁷ O. Adriani,¹⁶ M. Aguilar-Benitez,²⁶ S. Ahlen,¹¹ J. Alcaraz,²⁶ G. Alemani,²² J. Allaby,¹⁷ A. Aloisio,²⁹ M.G. Alvigi,²⁹ G. Ambrosi,¹⁹ H. Anderhub,⁴⁸ V.P. Andreev,³⁷ T. Angelescu,¹³ F. Anselmo,⁹ A. Arefiev,²⁸ T. Azemoon,³ T. Aziz,¹⁰ P. Bagnaia,³⁶ L. Baksay,⁴³ R.C. Ball,³ S. Banerjee,¹⁰ Sw. Banerjee,¹⁰ K. Banicz,⁴⁵ A. Barczyk,^{48,46} R. Barillere,¹⁷ L. Barone,³⁶ P. Bartalini,²² A. Baschirotto,²⁷ M. Basile,⁹ R. Battiston,³³ A. Bay,²² F. Becattini,¹⁶ U. Becker,¹⁵ F. Behner,⁴⁸ J. Berdugo,²⁶ P. Berges,¹⁵ B. Bertucci,³³ B.L. Betev,⁴⁸ S. Bhattacharya,¹⁰ M. Biasini,³³ A. Biland,⁴⁸ G.M. Bilei,³³ J.J. Blaising,⁴ S.C. Blyth,³⁴ G.J. Bobbink,² R. Bock,¹ A. Böhm,¹ L. Boldizar,¹⁴ B. Borgia,^{17,36} D. Bourilkov,⁴⁸ M. Bourquin,¹⁹ D. Boutigny,⁴ S. Braccini,¹⁹ J.G. Branson,³⁹ V. Brigljevic,⁴⁸ I.C. Brock,³⁴ A. Buffini,¹⁶ A. Buijs,⁴⁴ J.D. Burger,¹⁵ W.J. Burger,³³ J. Busenitz,⁴³ X.D. Cai,¹⁵ M. Campanelli,⁴⁸ M. Capell,¹⁵ G. Cara Romeo,⁹ G. Carlino,²⁹ A.M. Cartacci,¹⁶ J. Casaus,²⁶ G. Castellini,¹⁶ F. Cavallari,³⁶ N. Cavallo,²⁹ C. Cecchi,¹⁹ M. Cerrada,²⁶ F. Cesaroni,²³ M. Chamizo,²⁶ Y.H. Chang,⁵⁰ U.K. Chaturvedi,¹⁸ M. Chemarin,²⁵ A. Chen,⁵⁰ G. Chen,⁷ G.M. Chen,⁷ H.F. Chen,²⁰ H.S. Chen,⁷ M. Chen,¹⁵ G. Chiefari,²⁹ C.Y. Chien,⁵ L. Cifarelli,³⁸ F. Cindolo,⁹ C. Civinini,¹⁶ I. Clare,¹⁵ R. Clare,¹⁵ G. Coignet,⁴ A.P. Colijn,² N. Colino,²⁶ S. Costantini,⁸ F. Cotorobai,¹³ B. de la Cruz,²⁶ A. Csilling,¹⁴ T.S. Dai,¹⁵ R.D. Alessandro,¹⁶ R. de Asmundis,²⁹ A. Degre,⁴ K. Deiters,⁴⁶ P. Denes,³⁵ F. DeNotaristefani,³⁶ M. Diemoz,³⁶ D. van Dierendonck,² F. Di Lodovico,⁴⁸ C. Dionisi,^{17,36} M. Dittmar,⁴⁸ A. Dominguez,³⁹ A. Doria,²⁹ M.T. Dova,^{18,‡} E. Drago,²⁹ D. Duchesneau,⁴ P. Duinker,² I. Duran,⁴⁰ S. Easo,³³ H. El Mamouni,²⁵ A. Engler,³⁴ F.J. Eppling,¹⁵ F.C. Erne,² J.P. Ernenwein,²⁵ P. Extermann,¹⁹ M. Fabre,⁴⁶ R. Faccini,³⁶ M.A. Falagan,²⁶ S. Falciano,³⁶ A. Favara,¹⁶ J. Fay,²⁵ O. Fedin,³⁷ M. Felcini,⁴⁸ T. Ferguson,³⁴ F. Ferroni,³⁶ H. Fesefeldt,¹ E. Fiandrini,³³ J.H. Field,¹⁹ F. Filthaut,¹⁷ P.H. Fisher,¹⁵ I. Fisk,³⁹ G. Forconi,¹⁵ L. Fredj,¹⁹ K. Freudenreich,⁴⁸ C. Furetta,²⁷ Yu. Galaktionov,^{28,15} S.N. Ganguli,¹⁰ P. Garcia-Abia,⁶ M. Gataullin,³² S.S. Gau,¹² S. Gentile,³⁶ J. Gerald,⁵ N. Gheordanescu,¹³ S. Giagu,³⁶ S. Goldfarb,²² J. Goldstein,¹¹ Z.F. Gong,²⁰ A. Gougas,⁵ G. Gratta,³² M.W. Gruenewald,⁸ R. van Gulik,² V.K. Gupta,³⁵ A. Gurtu,¹⁰ L.J. Gutay,⁴⁵ D. Haas,⁶ B. Hartmann,¹ A. Hasan,³⁰ D. Hatzifotiadiou,⁹ T. Hebbeker,⁸ A. Herve,¹⁷ P. Hidas,¹⁴ J. Hirschfelder,³⁴ W.C. van Hoek,³¹ H. Hofer,⁴⁸ H. Hoorani,³⁴ S.R. Hou,⁵⁰ G. Hu,⁵ I. Iashvili,⁴⁷ B.N. Jin,⁷ L.W. Jones,³ P. de Jong,¹⁷ I. Josa-Mutuberria,²⁶ A. Kasser,²² R.A. Khan,¹⁸ D. Kamrad,⁴⁷ J.S. Kapustinsky,²⁴ Y. Karyotakis,⁴ M. Kaur,^{18,◇} M.N. Kienzle-Focacci,¹⁹ D. Kim,³⁶ D.H. Kim,⁴² J.K. Kim,⁴² S.C. Kim,⁴² W.W. Kinnison,²⁴ A. Kirkby,³² D. Kirkby,³² J. Kirkby,¹⁷ D. Kiss,¹⁴ W. Kittel,³¹ A. Klimentov,^{15,28} A.C. König,³¹ A. Kopp,⁴⁷ I. Korolko,²⁸ V. Koutsenko,^{15,28} R.W. Kraemer,³⁴ W. Krenz,¹ A. Kunin,^{15,28} P. Lacentre,^{47,‡,‡} P. Ladron de Guevara,²⁶ G. Landi,¹⁶ C. Lapoint,¹⁵ K. Lassila-Perini,⁴⁸ P. Laurikainen,²¹ A. Lavorato,³⁸ M. Lebeau,¹⁷ A. Lebedev,¹⁵ P. Lebrun,²⁵ P. Lecomte,⁴⁸ P. Lecocq,¹⁷ P. Le Coultre,⁴⁸ H.J. Lee,⁸ C. Leggett,³ J.M. Le Goff,¹⁷ R. Leiste,⁴⁷ E. Leonardi,³⁶ P. Levchenko,³⁷ C. Li,²⁰ C.H. Lin,⁵⁰ W.T. Lin,⁵⁰ F.L. Linde,^{2,17} L. Lista,²⁹ Z.A. Liu,⁷ W. Lohmann,⁴⁷ E. Longo,³⁶ W. Lu,³² Y.S. Lu,⁷ K. Lübelmeyer,¹ C. Luci,^{17,36} D. Luckey,¹⁵ L. Luminari,³⁶ W. Lustermaier,⁴⁸ W.G. Ma,²⁰ M. Maity,¹⁰ G. Majumder,¹⁰ L. Malgeri,¹⁷ A. Malinin,²⁸ C. Mañá,²⁶ D. Mangeol,³¹ P. Marchesini,⁴⁸ G. Marian,^{43,¶} A. Marin,¹¹ J.P. Martin,²⁵ F. Marzano,³⁶ G.G.G. Massaro,² K. Mazumdar,¹⁰ S. Mele,¹⁷ L. Merola,²⁹ M. Meschini,¹⁶ W.J. Metzger,³¹ M. von der Mey,¹ Y. Mi,²² D. Miganj,⁹ A. Mihul,³ A.J.W. van Mil,³¹ H. Milcent,¹⁷ G. Mirabelli,³⁶ J. Mnich,¹⁷ P. Molnar,⁸ B. Monteleoni,¹⁶ R. Moore,³ T. Moulík,¹⁰ R. Mount,³² F. Muheim,¹⁹ A.J.M. Muijs,² S. Nahn,¹⁵ M. Napolitano,²⁹ F. Nessi-Tedaldi,⁴⁸ H. Newman,³² T. Niessen,¹ A. Nippe,²² A. Nisati,³⁶ H. Nowak,⁴⁷ Y.D. Oh,⁴² G. Organtini,³⁶ R. Ostonen,²¹ S. Palit,¹² C. Palomares,²⁶ D. Pandoulas,¹ S. Paoletti,^{36,17} P. Paolucci,²⁹ H.K. Park,³⁴ I.H. Park,⁴² G. Pascale,³⁶ G. Passaleva,¹⁷ S. Patricelli,²⁹ T. Paul,¹² M. Pauluzzi,³³ C. Paus,¹⁷ F. Pauss,⁴⁸ D. Peach,¹⁷ Y.J. Pei,¹ S. Pensotti,²⁷ D. Perret-Gallix,⁴ B. Petersen,³¹ S. Petrák,⁸ A. Pevsner,⁵ D. Piccolo,²⁹ M. Pieri,¹⁶ P.A. Piroué,³⁵ E. Pistolesi,²⁷ V. Plyaskin,²⁸ M. Pohl,⁴⁸ V. Pojidaev,^{28,16} H. Postema,¹⁵ J. Pothier,¹⁷ N. Produit,¹⁹ D. Prokofiev,³⁷ J. Quartieri,³⁸ G. Rahal-Callot,⁴⁸ N. Raja,¹⁰ P.G. Rancoita,²⁷ M. Rattaggi,²⁷ G. Raven,³⁹ P. Raziš,³⁰ D. Ren,⁴⁸ M. Rescigno,³⁶ S. Reucroft,¹² T. van Rhee,⁴⁴ S. Riemann,⁴⁷ K. Riles,³ O. Rind,³ A. Robohm,⁴³ J. Rodin,³ B.P. Roe,³ L. Romero,²⁶ S. Rosier-Lees,⁴ Ph. Rosselet,²² S. Roth,¹ J.A. Rubio,¹⁷ D. Ruschmeier,⁸ H. Rykaczewski,⁴⁸ S. Sakar,³⁶ J. Salicio,¹⁷ E. Sanchez,²⁶ M.P. Sanders,³¹ M.E. Sarakinos,²¹ G. Sauvage,⁴ C. Schäfer,¹ V. Schegelsky,³⁷ S. Schmidt-Kaerst,¹ D. Schmitz,¹ M. Schneegans,⁴ N. Scholz,⁴⁸ H. Schopper,⁴⁹ D.J. Schotanus,³¹ J. Schwenke,¹ G. Schwering,¹ C. Sciacca,²⁹ D. Sciarrino,¹⁹ L. Servoli,³³ S. Shevchenko,³² N. Shivarov,⁴¹ V. Shoutko,²⁸ J. Shukla,²⁴ E. Shumilov,²⁸ A. Shvorob,³² T. Siedenbueg,¹ D. Son,⁴² V. Soulimov,²⁹ B. Smith,¹⁵ P. Spillantini,¹⁶ M. Steuer,¹⁵ D.P. Stickland,³⁵ H. Stone,³⁵ B. Stoyanov,⁴¹ A. Straessner,¹ K. Sudhakar,¹⁰ G. Sultanov,¹⁸ L.Z. Sun,²⁰ G.F. Susinno,¹⁹ H. Suter,⁴⁸ J.D. Swain,¹⁸ X.W. Tang,⁷ L. Tauscher,⁶ L. Taylor,¹² C. Timmermans,³¹ Samuel C.C. Ting,¹⁵ S.M. Ting,¹⁵ S.C. Tonwar,¹⁰ J. Tóth,¹⁴ C. Tully,³⁵ K.L. Tung,⁷ Y. Uchida,¹⁵ J. Ulbricht,⁴⁸ E. Valente,³⁶ G. Vesztegombi,¹⁴ I. Vetlitsky,²⁸ G. Viertel,⁴⁸ M. Vivargent,⁴ S. Vlachos,⁶ H. Vogel,³⁴ H. Vogt,⁴⁷ I. Vorobiev,^{17,28} A.A. Vorobyov,³⁷ A. Vorvolakos,³⁰ M. Wadhwa,⁶ W. Wallraff,¹ J.C. Wang,¹⁵ X.L. Wang,²⁰ Z.M. Wang,²⁰ A. Weber,¹ S.X. Wu,¹⁵ S. Wynhoff,⁷ J. Xu,¹¹ Z.Z. Xu,²⁰ B.Z. Yang,²⁰ C.G. Yang,⁷ H.J. Yang,⁷ M. Yang,⁷ J.B. Ye,²⁰ S.C. Yeh,⁵¹ J.M. You,³⁴ An. Zalite,³⁷ Yu. Zalite,³⁷ P. Zemp,⁴⁸ Y. Zeng,¹ Z.P. Zhang,²⁰ B. Zhou,¹¹ Y. Zhou,³ G.Y. Zhu,⁷ R.Y. Zhu,³² A. Zichichi,^{9,17,18} F. Ziegler,⁴⁷ G. Zilizi,^{43,¶}

- 1 I. Physikalisches Institut, RWTH, D-52056 Aachen, FRG[§]
 - III. Physikalisches Institut, RWTH, D-52056 Aachen, FRG[§]
 - 2 National Institute for High Energy Physics, NIKHEF, and University of Amsterdam, NL-1009 DB Amsterdam, The Netherlands
 - 3 University of Michigan, Ann Arbor, MI 48109, USA
 - 4 Laboratoire d'Annecy-le-Vieux de Physique des Particules, LAPP, IN2P3-CNRS, BP 110, F-74941 Annecy-le-Vieux CEDEX, France
 - 5 Johns Hopkins University, Baltimore, MD 21218, USA
 - 6 Institute of Physics, University of Basel, CH-4056 Basel, Switzerland
 - 7 Institute of High Energy Physics, IHEP, 100039 Beijing, China[△]
 - 8 Humboldt University, D-10099 Berlin, FRG[§]
 - 9 University of Bologna and INFN-Sezione di Bologna, I-40126 Bologna, Italy
 - 10 Tata Institute of Fundamental Research, Bombay 400 005, India
 - 11 Boston University, Boston, MA 02215, USA
 - 12 Northeastern University, Boston, MA 02115, USA
 - 13 Institute of Atomic Physics and University of Bucharest, R-76900 Bucharest, Romania
 - 14 Central Research Institute for Physics of the Hungarian Academy of Sciences, H-1525 Budapest 114, Hungary[‡]
 - 15 Massachusetts Institute of Technology, Cambridge, MA 02139, USA
 - 16 INFN Sezione di Firenze and University of Florence, I-50125 Florence, Italy
 - 17 European Laboratory for Particle Physics, CERN, CH-1211 Geneva 23, Switzerland
 - 18 World Laboratory, FBLJA Project, CH-1211 Geneva 23, Switzerland
 - 19 University of Geneva, CH-1211 Geneva 4, Switzerland
 - 20 Chinese University of Science and Technology, USTC, Hefei, Anhui 230 029, China[△]
 - 21 SEFT, Research Institute for High Energy Physics, P.O. Box 9, SF-00014 Helsinki, Finland
 - 22 University of Lausanne, CH-1015 Lausanne, Switzerland
 - 23 INFN-Sezione di Lecce and Università Degli Studi di Lecce, I-73100 Lecce, Italy
 - 24 Los Alamos National Laboratory, Los Alamos, NM 87544, USA
 - 25 Institut de Physique Nucléaire de Lyon, IN2P3-CNRS, Université Claude Bernard, F-69622 Villeurbanne, France
 - 26 Centro de Investigaciones Energeticas, Medioambientales y Tecnológicas, CIEMAT, E-28040 Madrid, Spain[‡]
 - 27 INFN-Sezione di Milano, I-20133 Milan, Italy
 - 28 Institute of Theoretical and Experimental Physics, ITEP, Moscow, Russia
 - 29 INFN-Sezione di Napoli and University of Naples, I-80125 Naples, Italy
 - 30 Department of Natural Sciences, University of Cyprus, Nicosia, Cyprus
 - 31 University of Nijmegen and NIKHEF, NL-6525 ED Nijmegen, The Netherlands
 - 32 California Institute of Technology, Pasadena, CA 91125, USA
 - 33 INFN-Sezione di Perugia and Università Degli Studi di Perugia, I-06100 Perugia, Italy
 - 34 Carnegie Mellon University, Pittsburgh, PA 15213, USA
 - 35 Princeton University, Princeton, NJ 08544, USA
 - 36 INFN-Sezione di Roma and University of Rome, "La Sapienza", I-00185 Rome, Italy
 - 37 Nuclear Physics Institute, St. Petersburg, Russia
 - 38 University and INFN, Salerno, I-84100 Salerno, Italy
 - 39 University of California, San Diego, CA 92093, USA
 - 40 Dept. de Física de Partículas Elementales, Univ. de Santiago, E-15706 Santiago de Compostela, Spain
 - 41 Bulgarian Academy of Sciences, Central Lab. of Mechatronics and Instrumentation, BU-1113 Sofia, Bulgaria
 - 42 Center for High Energy Physics, Adv. Inst. of Sciences and Technology, 305-701 Taejeon, Republic of Korea
 - 43 University of Alabama, Tuscaloosa, AL 35486, USA
 - 44 Utrecht University and NIKHEF, NL-3584 CB Utrecht, The Netherlands
 - 45 Purdue University, West Lafayette, IN 47907, USA
 - 46 Paul Scherrer Institut, PSI, CH-5232 Villigen, Switzerland
 - 47 DESY-Institut für Hochenergiephysik, D-15738 Zeuthen, FRG
 - 48 Eidgenössische Technische Hochschule, ETH Zürich, CH-8093 Zürich, Switzerland
 - 49 University of Hamburg, D-22761 Hamburg, FRG
 - 50 National Central University, Chung-Li, Taiwan, China
 - 51 Department of Physics, National Tsing Hua University, Taiwan, China
- § Supported by the German Bundesministerium für Bildung, Wissenschaft, Forschung und Technologie
- ‡ Supported by the Hungarian OTKA fund under contract numbers T019181, F023259 and T024011.
- ¶ Also supported by the Hungarian OTKA fund under contract numbers T22238 and T026178.
- ‡ Supported also by the Comisión Interministerial de Ciencia y Tecnología.
- ‡ Also supported by CONICET and Universidad Nacional de La Plata, CC 67, 1900 La Plata, Argentina.
- ‡ Supported by Deutscher Akademischer Austauschdienst.
- △ Also supported by Panjab University, Chandigarh-160014, India.
- △ Supported by the National Natural Science Foundation of China.




Figure 1: CP-violating and chirality-conserving $Z\mu\mu\gamma$ vertex.





Figure 2: Distributions of the CP-odd observables T and V in data (points) and Monte Carlo (histogram).





Figure 3: Measurement of the relative asymmetry for T and V . The dashed line shows the theory prediction for a variation of ± 1 of the CP-violating parameters $\hat{f}_1^\mu$ and $\hat{f}_2^\mu$.





Figure 4: Measurement of the absolute asymmetry for T and V . The dashed line shows the theory prediction for a variation of the CP-violating parameters as in Figure 3. The fit range for the determination of the CP-violating parameters $\hat{f}_1^\mu$ and $\hat{f}_2^\mu$ is indicated by the arrow.