

# Historical and new spectral indicators from the Nearby Supernova Factory

N. Chotard, E. Gangler, G. Smadja

► **To cite this version:**

N. Chotard, E. Gangler, G. Smadja. Historical and new spectral indicators from the Nearby Supernova Factory. Annual meeting of the French Society of Astronomy and Astrophysics, SF2A-2009, Jun 2009, Besançon, France. pp.203-205. in2p3-00590627

**HAL Id: in2p3-00590627**

**<http://hal.in2p3.fr/in2p3-00590627>**

Submitted on 5 May 2011

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## HISTORICAL AND NEW SPECTRAL INDICATORS FROM THE NEARBY SUPERNOVA FACTORY

Chotard, N.<sup>1</sup>, Gangler, E.<sup>1</sup>, Smadja, G.<sup>1</sup> and the SNFactory collaboration

**Abstract.** From a sample of 58 SNe Ia spectra obtained by the Nearby Supernova Factory with the SuperNovae Integral Field Spectrograph (SNIFS), we present measurements of spectral indicators proposed in the literature. In addition, we probe the direct correlation of spectral indicators to the Hubble diagram residuals and the relevance of such indicators as substitutes to stretch and colour.

### 1 The SNe Ia sample

The data presented here has been obtained by the Nearby SNF collaboration with SNIFS on the UH 2.2 *m* (Aldering et al. 2002). The light curves in synthetic BVR filters are reconstructed from the absolute spectra of each SN. The present study is performed on a sample of 58 SNe Ia selected for the quality of their SALT2 (Guy et al. 2007) light-curve fit, the phase coverage, and for which a spectrum in a window of  $\pm 2.5$  days around the peak luminosity was obtained. The parameters of the fit (stretch ( $x1$ ), colour ( $c$ )) are also used in this analysis.

### 2 Classical spectral indicators

Two regions of interests between 3500 and 4200 Å and between 5500 and 6500 Å have been shown to exhibit calcium and silicon features correlated to the supernova intrinsic parameters (Nugent et al. 1995; Bongard et al. 2005). They defined the following classical spectral indicators :

$$R_{Ca} = \frac{F_{max}^{3950}}{F_{max}^{3650}} \quad R_{Si} = \frac{\text{depth}_{5800}}{\text{depth}_{6200}} \quad R_{SiSS} = \frac{\int_{5500}^{5700} F(\lambda) d\lambda}{\int_{6200}^{6450} F(\lambda) d\lambda}$$

where  $F(\lambda)$  is the flux ( $[erg/cm^2/s/\text{Å}]$ ),  $F_{max}^\lambda$  is the flux at the peak position found around the given wavelength, and  $\text{depth}_\lambda$  is the difference between the continuum flux and the real minimum flux of the absorption feature. Computing these indicators in an objective way requires an automatic smoothing procedure which preserves spectral extrema, as shown in Fig. 1.

Other spectral indicators such as equivalent widths ( $EW$ ) (Hachinger et al. 2006; Bronder et al. 2007) were computed in relevant regions of the SN Ia spectrum. In our sample,  $EW_{SiII}(4000)$  (shown in Fig. 1 on the top right of the control plot) was found to correlate strongly with the SALT2  $x1$  parameter . The general definition of  $EW$  is:

$$EW = \sum_{i=1}^N \left( 1 - \frac{f_\lambda(\lambda_i)}{f_c(\lambda_i)} \right) \Delta\lambda_i$$

---

<sup>1</sup> Institut de Physique Nucl aire de Lyon, 4 rue Enrico Fermi, domaine scientifique de la DOUA 69622 Villeurbanne CEDEX, France




Fig. 1. Automatic control plot for a spectrum of the sample.

### 3 The correlation of spectral indicators with SNe Ia parameters

The relative standardization power of these indicators were studied with their correlations with  $x1$ , colour and a proxy to their absolute magnitude,  $\Delta M_B$ : the latter was derived from the residual of a fit to the Hubble diagram for a flat  $\Lambda$ CDM.

The Table 1 shows the correlation coefficients between parameters and spectral indicators. A SALT2 colour cut is applied on the sample ( $c < 0.1$ ), to study the intrinsic part of the SN Ia variability. 44 over our 58 SNe pass the cut. Fig. 2 shows the correlation between  $EW_{SiII}(4000)^{cut}$  and  $\Delta M_B$  after the colour cut.


	$\Delta M_B$	$x1$	colour	$\Delta M_B^{corr}$	phase
$RCa$	0.55	-0.45	0.35	0.17	0.05
$RSi$	0.26	-0.60	0.04	0.00	0.06
$RSiSS$	-0.77	-0.57	-0.54	-0.16	0.27
$EW_{SiII}(4000)$	0.33	-0.76	-0.01	0.17	0.23
$EW_{SiII}(4000)^{cut}$	<b>0.80</b>	<b>-0.84</b>	0.01	0.28	0.21

Table 1. Correlation coefficients between spectral indicators and other SN Ia parameters.

The  $EW_{SiII}(4000)$ 's correlations with Hubble residuals and  $x1$  **increase after the colour cut** (see Table 1 and Fig. 2).  $EW_{SiII}(4000)$  is independant of colour (the correlation coefficient is 0.01) and can be a good proxy for  $x1$ , it thus proves to be particularly useful for standardizing SNe Ia.

### 4 Corrected Hubble residuals

In the SALT2 approach, the luminosity is corrected for stretch and colour with  $\alpha$  and  $\beta$  tuned to minimize the residuals to the cosmological fit to the data:  $\Delta M_B^{corr} = \Delta M_B + \alpha \times x1 - \beta \times c$ . The spectral correction can be



**Fig. 2.** Correlation between  $\Delta M_B$  and  $EW_{SiII}(4000)^{cut}$ , after the colour cut ( $c < 0.1$ )

used in the similar way. Table 2 gives the RMS of the Hubble diagram residuals uncorrected and corrected with spectral indicators. The Hubble residual with colour and  $EW_{SiII}(4000)$  corrections are as effective as colour and  $x1$  before the colour cut and perform better after.  $EW_{SiII}(4000)$  is then an excellent candidate for the estimate the intrinsic part of the SN Ia variability and replaces or strengthens the  $x1$  parameter.

Correction	None	c & x1	c & $EW_{SiII}(4000)$	None	c & x1	c & $EW_{SiII}(4000)$
RMS	0.406	0.161	0.164	0.217	0.153	0.123
nMAD	0.264	0.159	0.177	0.243	0.139	0.148

**Table 2.** Standard deviation and normalized median absolute deviation of hubble diagram residual without any cut (left values) and with a colour cut (right values).

## 5 Conclusion

Traditional spectral indicators are correlated to the SNe Ia absolute luminosity, and some of them such as  $EW_{SiII}(4000)$  are strongly correlated to the intrinsic variability of SNe Ia. Another spectral indicator presented in (Bailey et al. 2009) using the same sample of SNe Ia from the SNFactory collaboration also shows a very strong correlation (0.95) between the spectral ratio  $R(642/443)$  and  $\Delta M_B$  even without any colour cut. It proves to be a strong competitor of  $x1$  and colour for the luminosity standardization and reaches a final RMS on the hubble diagram of  $0.128 \pm 0.012$ .

## References

- Aldering, G. 2002, Proceedings of the SPIE., vol. 4836 of Presented at SPIE Conference, 61-72  
 Bongard, S. 2005, arXiv:astro-ph/0512229v1  
 Bronder, T.J. 2007, arXiv:astro-ph/0709.0859v1  
 Guy, J., 2007 arXiv:astro-ph/0701828v1  
 Hachinger, S. 2006, arXiv:astro-ph/0604472v2  
 Nugent, P. 1995, ApJ, 455, L147