

Partial dynamical symmetry in Bose-Fermi systems

P. van Isacker, J. Jolie, T. Thomas, A. Leviatan

► **To cite this version:**

P. van Isacker, J. Jolie, T. Thomas, A. Leviatan. Partial dynamical symmetry in Bose-Fermi systems. Physical Review C, American Physical Society, 2015, 92, pp.011301. 10.1103/PhysRevC.92.011301. in2p3-01166162

HAL Id: in2p3-01166162

<http://hal.in2p3.fr/in2p3-01166162>

Submitted on 22 Jun 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Partial dynamical symmetry in Bose-Fermi systems

P. Van Isacker

Grand Accélérateur National d'Ions Lourds, CEA/DSM-CNRS/IN2P3, B.P. 55027, F-14076 Caen Cedex 5, France

J. Jolie and T. Thomas

Institute of Nuclear Physics, University of Cologne, Zùlpicherstrasse 77, 50937 Cologne, Germany

A. Leviatan

Racah Institute of Physics, The Hebrew University, Jerusalem 91904, Israel

(Dated: June 22, 2015)

We generalize the notion of partial dynamical symmetry (PDS) to a system of interacting bosons and fermions. In a PDS, selected states of the Hamiltonian are solvable and preserve the symmetry exactly, while other states are mixed. As a first example of such novel symmetry construction, spectral features of the odd-mass nucleus ^{195}Pt are analyzed.

PACS numbers: 21.60.Fw, 21.10.Re, 21.60.Ev, 27.80+w

During the last several decades, the concept of dynamical symmetry (DS) has become the cornerstone of algebraic modeling of dynamical systems. It has been applied in many branches of physics, such as hadronic [1], nuclear [2, 3], atomic [4] and molecular physics [5, 6]. Its basic paradigm is to write the Hamiltonian of the system in terms of Casimir operators of a chain of nested algebras, $G_{\text{dyn}} \supset G \supset \cdots \supset G_{\text{sym}}$, where G_{dyn} is the dynamical algebra, in terms of which any model operator of a physical observable can be expressed, and G_{sym} is the symmetry algebra. A given DS defines a class of many-body Hamiltonians that admit an analytic solution for all states, with closed expressions for the energy eigenvalues, quantum numbers for classification and definite selection rules for transition processes.

An exact DS provides considerable insights into complex dynamics and its merits are self evident. However, in most applications to realistic systems, its predictions are rarely fulfilled and one is compelled to break it. The DS spectrum imposes constraints on the pattern of level-splitting which many times is at variance with the empirical data. More often one finds that the assumed symmetry is not obeyed uniformly, *i.e.*, is fulfilled by some of the states but not by others. The required symmetry breaking is achieved by including in the Hamiltonian terms associated with different subalgebra chains of G_{dyn} , resulting in a loss of solvability and pronounced mixing. The need to address such situations, but still preserve important symmetry remnants, has led to the introduction of partial dynamical symmetry (PDS) [7, 8]. The essential idea is to relax the stringent conditions of *complete* solvability so that only part of the eigenspectrum retains analyticity and/or good quantum numbers, in the spirit of quasi-solvable models [9]. Various types of PDSs were proposed [8, 10–13] and algorithms for constructing Hamiltonians with such property have been developed [7, 14]. Bosonic Hamiltonians with PDS have been applied to nuclear spectroscopy [8, 10–17], where extensive tests provide empirical evidence for their relevance to a broad range of nuclei. Similar PDS Hamiltonians

have been used in molecular spectroscopy [18] and in the study of quantum phase transitions [19, 20] and of mixed regular and chaotic dynamics [20, 21]. Fermionic Hamiltonians with PDS have been identified within the nuclear shell model and applied to light nuclei [22] and seniority isomers [23, 24]. The growing number of empirical manifestations suggests a more pervasive role of PDSs in dynamical systems than heretofore realized.

All examples of PDS considered so far, were confined to systems of a given statistics (bosons or fermions). In this Rapid Communication, we extend the PDS concept to mixed systems of bosons and fermions, and present an empirical example of this novel construction. Systems with such composition of constituents are of broad interest and arise, for example, in the study of rotation-vibration-electronic spectra in molecules, collective states in odd-mass nuclei, electron-phonon phenomena in crystals, and spin-boson models in quantum optics.

If the separate numbers of bosons N and fermions M are conserved, the dynamical algebra of a Bose-Fermi system is of product form

$$\begin{array}{ccc} \text{U}^{\text{B}}(\Omega_{\text{B}}) & \otimes & \text{U}^{\text{F}}(\Omega_{\text{F}}) \\ \downarrow & & \downarrow \\ [N] & & [1^M] \end{array}, \quad (1)$$

where Ω_{B} (Ω_{F}) is the number of states available to a single boson (fermion). The statistics among the particles is imposed by an appropriate choice of irreducible representation (irrep), symmetric and anti-symmetric, for the bosons and fermions, respectively, as indicated in Eq. (1). There exist several strategies to define DSs with $\text{U}^{\text{B}}(\Omega_{\text{B}}) \otimes \text{U}^{\text{F}}(\Omega_{\text{F}})$ as a starting point [3]. They all define a chain of nested subalgebras, relying on the existence of isomorphisms between boson and fermion algebras and ending in the symmetry algebra.

Let us for the sake of concreteness consider a particular example while emphasizing that results of this Rapid Communication are of a generic nature that apply to any quantum-mechanical problem of interacting bosons and

fermions, as long as it can be formulated in an algebraic language. We consider N bosons with angular momentum $\ell = 0$ (s) or $\ell = 2$ (d) coupled to a single ($M = 1$)

fermion with angular momentum $j = 1/2, 3/2$, or $5/2$. This corresponds to the choice $\Omega_B = 6$ and $\Omega_F = 12$, and a possible classification is as follows:

$$\begin{array}{ccccccc} \text{U}^{\text{B}}(6) \otimes \text{U}^{\text{F}}(12) \supset & \left(\text{U}^{\text{BF}}(6) \supset \text{SO}^{\text{BF}}(6) \supset \text{SO}^{\text{BF}}(5) \supset \text{SO}^{\text{BF}}(3) \right) & \otimes & \text{SU}^{\text{F}}(2) \supset \text{Spin}^{\text{BF}}(3) \\ \downarrow & \downarrow & & \downarrow & \downarrow & \downarrow & \downarrow \\ [N] & [1^M] & [N_1, N_2] & \langle \sigma_1, \sigma_2 \rangle & (\tau_1, \tau_2) & L & \tilde{s} & J \end{array}, \quad (2)$$

where underneath each algebra the associated irrep labels are indicated, and G^{BF} is the direct sum of G^{B} and G^{F} . For $M = 0$, the classification (2) reduces to the $\text{SO}(6)$ limit of the interacting boson model [25] which is of relevance for the even-even platinum isotopes [26]. For $M = 1$, the classification (2) is proposed in the context of the interacting boson-fermion model (IBFM) [3] to describe odd-mass isotopes of platinum with the odd neutron in the orbits $3p_{1/2}$, $3p_{3/2}$, and $2f_{5/2}$, which are dominant for these isotopes [27, 28]. Since we are interested here in Bose-Fermi systems, we apply the classification (2) for $M = 1$, which implies $\tilde{s} = 1/2$, and refer to it as the $\text{SO}^{\text{BF}}(6)$ limit.

The eigenstates (2) are obtained with a Hamiltonian that is a combination of Casimir operators $\hat{C}_n[G]$ of order n of an algebra G appearing in the chain. Up to a constant energy, this Hamiltonian is of the form

$$\hat{H}_{\text{DS}} = a \hat{C}_2[\text{U}^{\text{BF}}(6)] + b \hat{C}_2[\text{SO}^{\text{BF}}(6)] + c \hat{C}_2[\text{SO}^{\text{BF}}(5)] + d \hat{C}_2[\text{SO}^{\text{BF}}(3)] + d' \hat{C}_2[\text{Spin}^{\text{BF}}(3)]. \quad (3)$$

The associated eigenvalue problem is analytically solvable, leading to the energy expression

$$E_{\text{DS}} = a f_5(N_1, N_2) + b f_4(\sigma_1, \sigma_2) + c f_3(\tau_1, \tau_2) + d L(L+1) + d' J(J+1), \quad (4)$$

with $f_i(s_1, s_2) \equiv s_1(s_1 + i) + s_2(s_2 + i - 2)$. The energy spectrum of the Hamiltonian (3) is then determined once the allowed values of $[N_1, N_2]$, $\langle \sigma_1, \sigma_2 \rangle$, (τ_1, τ_2) , L , and J for a given N and $M = 1$ are found. Such branching rules can be obtained with standard group-theoretical techniques [29]. While \hat{H}_{DS} (3) is *completely* solvable, the question arises whether terms can be added that preserve solvability for *part* of its spectrum. This can be achieved by the construction of a PDS.

The algorithm to construct a PDS [14] starts from the character under the classification (2) of the boson and fermion creation operators $b_{\ell m_\ell}^\dagger$ and $a_{j m_j}^\dagger$ [3]. Annihilation operators $b_{\ell m_\ell}$ and $a_{j m_j}$ transform in the same manner under orthogonal algebras if they are modified according to $\tilde{b}_{\ell m_\ell} \equiv (-)^{\ell+m_\ell} b_{\ell, -m_\ell}$ and $\tilde{a}_{j m_j} \equiv (-)^{j+m_j} a_{j, -m_j}$. The single-fermion angular momentum ($j = 1/2, 3/2, 5/2$) can be divided into a pseudo-orbital angular momentum ($\tilde{\ell} = 0, 2$) coupled to a pseudo-spin

($\tilde{s} = 1/2$). The resulting $\tilde{\ell}$ - \tilde{s} basis is given by $c_{\tilde{\ell} \tilde{m}_\ell; \tilde{s} \tilde{m}_s}^\dagger = \sum_{j, m} (\tilde{\ell}, \tilde{m}_\ell; \tilde{s}, \tilde{m}_s | j, m) a_{j m}^\dagger$.

Composite operators with definite tensor character under the classification (2) can be constructed by use of generalized coupling coefficients which can be written as a product of $\text{U}(6) \supset \text{SO}(6)$, $\text{SO}(6) \supset \text{SO}(5)$, and $\text{SO}(5) \supset \text{SO}(3)$ isoscalar factors [29]. For the two-particle operators (needed for the construction of a two-body interaction) the tensor character is uniquely specified by the $\text{SO}(6)$ and $\text{SO}(5)$ labels $\langle \sigma_1, \sigma_2 \rangle$ and (τ_1, τ_2) , together with the $\text{SO}^{\text{BF}}(3)$ and $\text{Spin}^{\text{BF}}(3)$ labels \mathcal{L} and \mathcal{J} . For example, operators that create a boson and a fermion with tensor character $[N_1, N_2] \langle \sigma_1, \sigma_2 \rangle (\tau_1, \tau_2) \mathcal{L} \mathcal{J}$ are

$$\sum_{\tau \tilde{\tau}} \left\langle \begin{array}{c} \langle 1 \rangle \\ (\tau) \end{array} \begin{array}{c} \langle 1 \rangle \\ (\tilde{\tau}) \end{array} \middle| \begin{array}{c} \langle \sigma_1, \sigma_2 \rangle \\ (\tau_1, \tau_2) \end{array} \right\rangle \sum_j \bar{\mathcal{L}} \bar{\mathcal{J}} \left\{ \begin{array}{c} \ell \\ \tilde{s} \end{array} \begin{array}{c} \tilde{\ell} \\ \mathcal{J} \end{array} \begin{array}{c} \mathcal{L} \\ j \end{array} \right\} (b_{\ell}^\dagger a_{j}^\dagger)_{M_{\mathcal{J}}}^{(\mathcal{J})}$$

with $(\ell, \tilde{\ell}) = 0, 2$, for $(\tau, \tilde{\tau}) = 0, 1$, respectively, $j = \tilde{\ell} \pm 1/2$, $\bar{x} \equiv \sqrt{2x+1}$ and the symbol $\langle : : | : : \rangle$ is an $\text{SO}(6) \supset \text{SO}(5)$ isoscalar factor, tabulated in Ref. [3]. Expressions for the two-particle creation operators of interest here are given in Table I. The corresponding annihilation operators with the correct tensor properties follow from $\tilde{\mathcal{T}}_{-, M_{\mathcal{J}}}^{\mathcal{L}(\mathcal{J})} = (-)^{\mathcal{J}+M_{\mathcal{J}}} \left(\mathcal{T}_{+, -M_{\mathcal{J}}}^{\mathcal{L}(\mathcal{J})} \right)^\dagger$, where $\mathcal{T} = \mathcal{U}$ or \mathcal{V} .

The lowest-lying states in the spectrum of an odd-mass nucleus, described in terms of N bosons and one fermion, can be written as $|[N+1] \langle N+1 \rangle (\tau) L J M_J \rangle$; the next class of state belongs to $|[N, 1] \langle N, 1 \rangle (\tau_1, \tau_2) L J M_J \rangle$ while there is also some evidence from one-neutron transfer for $|[N, 1] \langle N-1 \rangle (\tau) L J M_J \rangle$ states [30]. All two-particle operators listed in Table I annihilate particular states, hence lead to a PDS of some kind. For example, the operators with $\text{U}^{\text{BF}}(6)$ labels $[N_1, N_2] = [1, 1]$ satisfy

$$\tilde{\mathcal{U}}_{-, M_{\mathcal{J}}}^{\mathcal{L}(\mathcal{J})} |[N+1] \langle \sigma \rangle (\tau) L J M_J \rangle = 0, \quad (5)$$

for all permissible $(\sigma \tau L J M_J)$. This is so because a state with $N-1$ bosons and no fermion has the $\text{U}^{\text{BF}}(6)$ label $[N-1]$. Given the multiplication $[N-1] \otimes [1, 1] = [N, 1] \oplus [N-1, 1, 1]$, the action of a $\mathcal{U}_{+, -M_{\mathcal{J}}}^{\mathcal{L}(\mathcal{J})}$ operator on an $(N-1)$ -boson state can never yield a boson-fermion state with the $\text{U}^{\text{BF}}(6)$ labels $[N+1]$. Similar arguments involving $\text{SO}(6)$ multiplication lead to the following properties for

TABLE I: Two-particle tensor operators in the $\text{SO}^{\text{BF}}(6)$ limit. The superscript $\mathcal{L}(\mathcal{J})$ stands for the coupling $\mathcal{J} = \mathcal{L} \pm 1/2$.

N	M	$[N_1, N_2]$	$\langle \sigma_1, \sigma_2 \rangle$	(τ_1, τ_2)	\mathcal{L}	\mathcal{J}	Tensor operator $\mathcal{T}_{+,M\mathcal{J}}^{\mathcal{L}(\mathcal{J})}$
2	0	[2, 0]	$\langle 0, 0 \rangle$	(0, 0)	0	0	$\mathcal{V}_{+,0}^{0(0)} \equiv \sqrt{\frac{5}{12}}(d^\dagger d^\dagger)_0^{(0)} - \sqrt{\frac{1}{12}}(s^\dagger s^\dagger)_0^{(0)}$
1	1	[2, 0]	$\langle 0, 0 \rangle$	(0, 0)	0	1/2	$\mathcal{V}_{+,\mu}^{0(1/2)} \equiv -\sqrt{\frac{1}{6}}(s^\dagger a_{1/2}^\dagger)_\mu^{(1/2)} - \sqrt{\frac{1}{3}}(d^\dagger a_{3/2}^\dagger)_\mu^{(1/2)} + \sqrt{\frac{1}{2}}(d^\dagger a_{5/2}^\dagger)_\mu^{(1/2)}$
1	1	[1, 1]	$\langle 1, 1 \rangle$	(1, 1)	1	1/2	$\mathcal{U}_{+,\mu}^{1(1/2)} \equiv \sqrt{\frac{3}{5}}(d^\dagger a_{3/2}^\dagger)_\mu^{(1/2)} + \sqrt{\frac{2}{5}}(d^\dagger a_{5/2}^\dagger)_\mu^{(1/2)}$
1	1	[1, 1]	$\langle 1, 1 \rangle$	(1, 1)	1	3/2	$\mathcal{U}_{+,\mu}^{1(3/2)} \equiv -\sqrt{\frac{3}{10}}(d^\dagger a_{3/2}^\dagger)_\mu^{(3/2)} + \sqrt{\frac{7}{10}}(d^\dagger a_{5/2}^\dagger)_\mu^{(3/2)}$
1	1	[1, 1]	$\langle 1, 1 \rangle$	(1, 0)	2	3/2	$\mathcal{U}_{+,\mu}^{2(3/2)} \equiv \sqrt{\frac{1}{2}}(s^\dagger a_{3/2}^\dagger)_\mu^{(3/2)} - \sqrt{\frac{1}{2}}(d^\dagger a_{1/2}^\dagger)_\mu^{(3/2)}$
1	1	[1, 1]	$\langle 1, 1 \rangle$	(1, 0)	2	5/2	$\mathcal{U}_{+,\mu}^{2(5/2)} \equiv \sqrt{\frac{1}{2}}(s^\dagger a_{5/2}^\dagger)_\mu^{(5/2)} - \sqrt{\frac{1}{2}}(d^\dagger a_{1/2}^\dagger)_\mu^{(5/2)}$
1	1	[1, 1]	$\langle 1, 1 \rangle$	(1, 1)	3	5/2	$\mathcal{U}_{+,\mu}^{3(5/2)} \equiv \sqrt{\frac{4}{5}}(d^\dagger a_{3/2}^\dagger)_\mu^{(5/2)} + \sqrt{\frac{1}{5}}(d^\dagger a_{5/2}^\dagger)_\mu^{(5/2)}$
1	1	[1, 1]	$\langle 1, 1 \rangle$	(1, 1)	3	7/2	$\mathcal{U}_{+,\mu}^{3(7/2)} \equiv -\sqrt{\frac{1}{10}}(d^\dagger a_{3/2}^\dagger)_\mu^{(7/2)} + \sqrt{\frac{9}{10}}(d^\dagger a_{5/2}^\dagger)_\mu^{(7/2)}$

the \mathcal{V} operators which have $\text{SO}(6)$ tensor character $\langle 0, 0 \rangle$:

$$\tilde{\mathcal{V}}_{-,M\mathcal{J}}^{0(\mathcal{J})} |[N+1]\langle N+1 \rangle (\tau) LJM_J\rangle = 0, \quad (6a)$$

$$\tilde{\mathcal{V}}_{-,M\mathcal{J}}^{0(\mathcal{J})} |[N, 1]\langle N, 1 \rangle (\tau_1, \tau_2) LJM_J\rangle = 0. \quad (6b)$$

An alternative way of constructing Hamiltonians with PDS for an algebra G , is to identify n -particle operators which annihilate a lowest-weight state of a prescribed G -irrep [7]. In the IBFM, such a state, which transforms as $[N+1]$ and $\tilde{s} = 1/2$ under $\text{U}^{\text{BF}}(6) \otimes \text{SU}^{\text{F}}(2)$, is given by

$$|\Psi_g\rangle \propto [b_c^\dagger(\beta)]^N f_{\tilde{m}_s}^\dagger(\beta) |0\rangle, \quad (7)$$

where $b_c^\dagger(\beta) \propto (\beta d_0^\dagger + s^\dagger)$ and $f_{\tilde{m}_s}^\dagger(\beta) \propto (\beta c_{2,0;1/2,\tilde{m}_s}^\dagger + c_{0,0;1/2,\tilde{m}_s}^\dagger)$ in the $\tilde{\ell}\tilde{s}$ basis defined above. $|\Psi_g\rangle$ is a condensate of N bosons and a single fermion, and represents an intrinsic state for the ground band with deformation β . The Hermitian conjugate of the following two-particle operators

$$\mathcal{V}_+^{0(0)} = \sqrt{\frac{5}{12}}(d^\dagger d^\dagger)_0^{(0)} - \frac{\beta^2}{\sqrt{12}}(s^\dagger s^\dagger)_0^{(0)}, \quad (8a)$$

$$\mathcal{V}_{+,\mu}^{0(1/2)} = \sqrt{\frac{5}{6}}(d^\dagger c_{2;1/2}^\dagger)_\mu^{0(1/2)} - \frac{\beta^2}{\sqrt{6}}(s^\dagger c_{0;1/2}^\dagger)_\mu^{0(1/2)}, \quad (8b)$$

$$\mathcal{U}_{+,\mu}^{\mathcal{L}(\mathcal{J})} = (d^\dagger c_{2;1/2}^\dagger)_\mu^{\mathcal{L}(\mathcal{J})}, \quad \mathcal{L} = 1, 3, \quad (8c)$$

$$\mathcal{U}_{+,\mu}^{2(\mathcal{J})} = (s^\dagger c_{2;1/2}^\dagger)_\mu^{2(\mathcal{J})} - (d^\dagger c_{0;1/2}^\dagger)_\mu^{2(\mathcal{J})}, \quad (8d)$$

satisfy $\tilde{\mathcal{T}}_{-,\mu}^{\mathcal{L}(\mathcal{J})} |\Psi_g\rangle = 0$. The \mathcal{V} operators of Eqs. (8a)-(8b) satisfy also $\tilde{\mathcal{V}}_{-,\mu}^{0(\mathcal{J})} |\Psi_e\rangle = 0$, where

$$|\Psi_e\rangle \propto [b_c^\dagger(\beta) c_{2,1;1/2,\tilde{m}_s}^\dagger - d_1^\dagger f_{\tilde{m}_s}^\dagger(\beta)] [b_c^\dagger(\beta)]^{N-1} |0\rangle \quad (9)$$

is an intrinsic state, with $\text{U}^{\text{BF}}(6)$ label $[N, 1]$, representing an excited band in the odd-mass nucleus. For $\beta = 1$, $|\Psi_g\rangle$ and $|\Psi_e\rangle$ become the lowest-weight states in the $\text{SO}^{\text{BF}}(6)$ irreps $\langle N+1 \rangle$ and $\langle N, 1 \rangle$, respectively, from which the $|(\tau_1, \tau_2) LJM_J\rangle$ states of Eq. (6) can be obtained by $\text{SO}^{\text{BF}}(5)$ projection, and the operators (8) coincide with those listed in Table I.

The combined effect of normal-ordered interactions constructed out of the \mathcal{T} operators in Table I added to the DS Hamiltonian (3), gives rise to a rich variety of possible PDSs. In the current application to ^{195}Pt we take a restricted Hamiltonian of the form

$$\hat{H}_{\text{PDS}} = \hat{H}_{\text{DS}} + a_0 \hat{V}_0^{1/2} + a'_1 (2\hat{U}_1^{1/2} - \hat{U}_1^{3/2}) + a_2 (\hat{U}_2^{3/2} + \hat{U}_2^{5/2}) + a_3 (\hat{U}_3^{5/2} + \hat{U}_3^{7/2}), \quad (10)$$

in terms of the interactions $\hat{T}_{\mathcal{L}}^{\mathcal{J}} \equiv \tilde{\mathcal{J}}(\mathcal{T}_+^{\mathcal{L}(\mathcal{J})} \tilde{\mathcal{T}}_-^{\mathcal{L}(\mathcal{J})})_0^{(0)}$, where $\mathcal{T} = \mathcal{U}$ or \mathcal{V} and $T = U$ or V . These interactions can be transcribed as tensors with total pseudo-orbital \tilde{L} and pseudo-spin \tilde{S} coupled to zero total angular momentum. In particular, the a'_1 term in Eq. (10) has $\tilde{L} = \tilde{S} = 1$, while the a_0, a_2 and a_3 terms have $\tilde{L} = \tilde{S} = 0$.

The experimental spectrum of ^{195}Pt is shown in Fig. 1, compared with the DS and PDS calculations. The coefficients c, d , and d' in \hat{H}_{DS} (3) are adjusted to the excitation energies of the $[7, 0]\langle 7, 0 \rangle$ levels which are reproduced with a root-mean-square (rms) deviation of 12 keV. The remaining two coefficients a and b are obtained from an overall fit. The resulting (DS) values are (in keV): $a = 45.3, b = -41.5, c = 49.1, d = 1.7$, and $d' = 5.6$. The fit for the PDS calculation proceeds in stages. First, the parameters c, d , and d' in Eq. (3) are taken at their DS values. This ensures the same spectrum for the $[7, 0]\langle 7, 0 \rangle$ levels (drawn in black in Fig. 1) which remain eigenstates of \hat{H}_{PDS} (10). Next, one considers the $[6, 1]\langle 6, 1 \rangle$ levels and improves their description by adding the three PDS U interactions. The resulting coefficients are (in keV): $a'_1 = 10, a_2 = -97$, and $a_3 = 112$. Eq. (5) ensures that the energies of the $[7, 0]\langle 7, 0 \rangle$ levels do not change while the agreement for the $[6, 1]\langle 6, 1 \rangle$ levels is improved (blue levels in Fig. 1). The rms deviation for both classes of levels is 20 keV. In particular, unlike in the DS calculation, it is possible to reproduce the observed inversion of the $1/2^- - 3/2^-$ doublets without changing the order of other doublets. The additional PDS terms necessitate a readjustment of the a coefficient in Eq. (3), for which the final (PDS) value is $a = 37.7$ keV, while the coefficient b is kept unchanged. Finally, the position of the $[6, 1]\langle 5, 0 \rangle$




FIG. 1: (Color online). Left panel: Observed and calculated energy spectrum of ^{195}Pt . The levels in black are the solvable $[7,0]\langle 7,0\rangle$ eigenstates of \hat{H}_{DS} (3), whose structure and energy remain unaffected by the added PDS interactions in Eq. (10). The levels in blue (red) are the $[6,1]\langle 6,1\rangle$ ($[6,1]\langle 5,0\rangle$) eigenstates of \hat{H}_{DS} (3) and are subsequently perturbed by the PDS interactions in Eq. (10). Right panel: $\text{SO}^{\text{BF}}(6)$ decomposition of the eigenstates of \hat{H}_{PDS} (10), shown in red on the left panel.

TABLE II: Observed $B(E2; J_i \rightarrow J_f)$ values between negative-parity states in ^{195}Pt compared with the DS and PDS predictions of the $\text{SO}^{\text{BF}}(6)$ limit. The solvable (mixed) states are members of the ground (excited) bands shown in Fig. 1. The E2 operator employed is defined in the text.

E_i (keV)	J_i	E_f (keV)	J_f	$B(E2; J_i \rightarrow J_f)$ ($10^{-3} e^2 b^2$)		
				Exp	DS	PDS
Solvable \rightarrow solvable						
212	3/2	0	1/2	190(10)	179	179
239	5/2	0	1/2	170(10)	179	179
525	3/2	0	1/2	17(1)	0	0
525	3/2	239	5/2	≤ 19	72	72
544	5/2	0	1/2	8(4)	0	0
612	7/2	212	3/2	170(70)	215	215
667	9/2	239	5/2	200(40)	239	239
Solvable \rightarrow mixed						
239	5/2	99	3/2	60(20)	0	0
525	3/2	99	3/2	≤ 33	7	3
525	3/2	130	5/2	9(5)	3	2
612	7/2	99	3/2	5(3)	9	11
667	9/2	130	5/2	12(3)	10	12
Mixed \rightarrow solvable						
99	3/2	0	1/2	38(6)	35	34
130	5/2	0	1/2	66(4)	35	33
420	3/2	0	1/2	15(1)	0	0
456	5/2	0	1/2	≤ 0.04	0	0
508	7/2	212	3/2	55(17)	20	18
563	9/2	239	5/2	91(22)	22	22
199	3/2	0	1/2	25(2)	0	0
390	5/2	0	1/2	7(1)	0	0
Mixed \rightarrow mixed						
420	3/2	99	3/2	5(4)	177	165
508	7/2	99	3/2	240(50)	228	263
563	9/2	130	5/2	240(40)	253	284
390	5/2	99	3/2	200(70)	219	179
390	5/2	130	5/2	≤ 14	55	35

levels (red levels in Fig. 1) is corrected by considering the PDS V interaction with $a_0 = 306$ keV which, due to Eq. (6), has a marginal effect on lower bands. The rms deviation for all levels shown in Fig. 1 is 23 keV. While the states $[7,0]\langle 7,0\rangle$ of the ground band are pure, other eigenstates of \hat{H}_{PDS} in excited bands can have substantial $\text{SO}^{\text{BF}}(6)$ mixing (see right panel of Fig. 1).

A large amount of information also exists on electromagnetic transition rates and spectroscopic strengths. In Table II, 25 measured $B(E2)$ values in ^{195}Pt are compared with the DS and PDS predictions. The same E2 operator is used as in previous studies [31, 32] of the $\text{SO}^{\text{BF}}(6)$ limit, $\hat{T}_\mu(E2) = e_b \hat{Q}_\mu^{\text{B}} + e_f \hat{Q}_\mu^{\text{F}}$, where $\hat{Q}_\mu^{\text{B}} = s^\dagger \tilde{d}_\mu + d_\mu^\dagger s$ is the boson quadrupole operator, \hat{Q}_μ^{F} is its fermion analogue [3], and e_b and e_f are effective boson and fermion charges, with the values $e_b = -e_f = 0.151$ eb. Table II is subdivided in four parts according to whether the initial and/or final state in the transition has a DS structure (as in Refs. [31, 32]) or whether it is mixed by the PDS interaction. It is seen that when both have a DS structure the $B(E2)$ value does not change, only slight differences occur when either the initial or the final state is mixed, and the biggest changes arise when both are mixed.

In summary, we have proposed a novel extension of the PDS notion to Bose-Fermi systems and exemplified it in ^{195}Pt . The analysis highlights the ability of a PDS to select and add to the Hamiltonian, in a controlled fashion, required symmetry-breaking terms, yet retain a good DS for a segment of the spectrum. These virtues greatly enhance the scope of applications of algebraic modeling of complex systems. The operators (8) with $\beta \neq 1$, can be used to explore additional PDSs in odd-mass nuclei. Partial supersymmetry, of relevance to nuclei [33], can be studied by embedding the algebras of Eq. (1) in a graded Lie algebra. Work in these directions is in progress.

We thank Stefan Heinze for his help with the numerical calculations. J.J. acknowledges financial support by GANIL and by the Interuniversity Attraction Poles programme initiated by the Belgian Science Policy Officer

under grant BrixNetwork P7/12 during a sabbatical stay at GANIL. A.L. acknowledges the hospitality of the Theoretical Division at LANL during a sabbatical stay, and the support by the Israel Science Foundation.

-
- [1] A. Bohm, Y. Néeman, and A. O. Barut Eds. *Dynamical Groups and Spectrum Generating Algebras* (World Scientific, Singapore, 1988).
- [2] F. Iachello and A. Arima, *The Interacting Boson Model* (Cambridge University Press, Cambridge, 1987).
- [3] F. Iachello and P. Van Isacker, *The Interacting Boson-Fermion Model* (Cambridge University Press, Cambridge, 1991).
- [4] K. Kikoin, M. Kiselev, and Y. Avishai, *Dynamical Symmetries for Nanostructures* (Springer-Verlag, Berlin, 2012).
- [5] F. Iachello and R. D. Levine, *Algebraic Theory of Molecules* (Oxford University Press, Oxford, 1994).
- [6] A. Frank and P. Van Isacker, *Algebraic Methods in Molecular and Nuclear Physics*, (Wiley, New York, 1994).
- [7] Y. Alhassid and A. Leviatan, *J. Phys. A* **25**, L1265 (1992).
- [8] A. Leviatan, *Phys. Rev. Lett.* **77**, 818 (1996).
- [9] A.G. Ushveridze, *Quasi-Exactly Solvable Models in Quantum Mechanics* (Taylor and Francis Group, New York 1994).
- [10] A. Leviatan, A. Novoselsky, and I. Talmi, *Phys. Lett. B* **172**, 144 (1986).
- [11] P. Van Isacker, *Phys. Rev. Lett.* **83**, 4269 (1999).
- [12] A. Leviatan and P. Van Isacker, *Phys. Rev. Lett.* **89**, 222501 (2002).
- [13] A. Leviatan, *Prog. Part. Nucl. Phys.* **66**, 93 (2011).
- [14] J. E. García-Ramos, A. Leviatan, and P. Van Isacker, *Phys. Rev. Lett.* **102**, 112502 (2009).
- [15] A. Leviatan, J. E. García-Ramos, and P. Van Isacker, *Phys. Rev. C* **87**, 021302(R) (2013).
- [16] R. F. Casten, R. B. Cakirli, K. Blaum, and A. Couture, *Phys. Rev. Lett.* **113**, 112501 (2014); A. Couture, R. F. Casten, and R. B. Cakirli, *Phys. Rev. C* **91**, 014312 (2015).
- [17] C. Kremer, J. Beller, A. Leviatan, N. Pietralla, G. Rainovski, R. Trippel, and P. Van Isacker, *Phys. Rev. C* **89**, 041302(R) (2014).
- [18] J. L. Ping and J. Q. Chen, *Ann. Phys. (N.Y.)* **255**, 75 (1997).
- [19] A. Leviatan, *Phys. Rev. Lett.* **98**, 242502 (2007).
- [20] M. Macek and A. Leviatan, *Ann. Phys. (N.Y.)* **351**, 302 (2014).
- [21] N. Whelan, Y. Alhassid, and A. Leviatan, *Phys. Rev. Lett.* **71**, 2208 (1993); A. Leviatan and N. D. Whelan, *Phys. Rev. Lett.* **77**, 5202 (1996).
- [22] J. Escher and A. Leviatan, *Phys. Rev. Lett.* **84**, 1866 (2000); *Phys. Rev. C* **65**, 054309 (2002).
- [23] D. J. Rowe and G. Rosensteel, *Phys. Rev. Lett.* **87**, 172501 (2001); G. Rosensteel and D. J. Rowe, *Phys. Rev. C* **67**, 014303 (2003).
- [24] P. Van Isacker and S. Heinze, *Phys. Rev. Lett.* **100**, 052501 (2008); *Ann. Phys. (N.Y.)* **349**, 73 (2014).
- [25] A. Arima and F. Iachello, *Ann. Phys. (N.Y.)* **123**, 468 (1979).
- [26] J. A. Cizewski, R. F. Casten, G. J. Smith, M. L. Stelts, W. R. Kane, H. G. Börner, and W. F. Davidson, *Phys. Rev. Lett.* **40**, 167 (1978).
- [27] P. Van Isacker, A. Frank, and H.-Z. Sun, *Ann. Phys. (N.Y.)* **157**, 183 (1984).
- [28] R. Bijker and F. Iachello, *Ann. Phys. (N.Y.)* **161**, 360 (1985).
- [29] B. G. Wybourne, *Classical Groups for Physicists* (Wiley, New York, 1974).
- [30] A. Metz, Y. Eisermann, A. Gollwitzer, R. Hertenberger, B. D. Valnion, G. Graw, and J. Jolie, *Phys. Rev. C* **61**, 064313 (2000); *Phys. Rev. C* **67**, 049901(E) (2003).
- [31] A. M. Bruce, W. Gelletly, J. Lukasiak, W. R. Phillips, and D. D. Warner, *Phys. Lett. B* **165**, 43 (1985).
- [32] A. Mauthofer, K. Stelzer, J. Gerl, Th. W. Elze, Th. Happ, G. Eckert, T. Faestermann, A. Frank, and P. Van Isacker, *Phys. Rev. C* **34**, 1958 (1986).
- [33] A. Metz, J. Jolie, G. Graw, R. Hertenberger, J. Gröger, C. Günther, N. Warr, and Y. Eisermann, *Phys. Rev. Lett.* **83**, 1542 (1999).