

## Long term variability of the blazar PKS 2155-304

J. Chevalier, M.A. Kastendieck, F. Rieger, G. Maurin, J.-P. Lenain, G.  
Lamanna

► **To cite this version:**

J. Chevalier, M.A. Kastendieck, F. Rieger, G. Maurin, J.-P. Lenain, et al.. Long term variability of the blazar PKS 2155-304. 34th International Cosmic Ray Conference (ICRC2015), Jul 2015, The Hague, Netherlands. PoS Proceedings of Science. <in2p3-01186534>

**HAL Id: in2p3-01186534**

**<http://hal.in2p3.fr/in2p3-01186534>**

Submitted on 25 Aug 2015

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## 1 Long term variability of the blazar PKS 2155-304

---

### Jill Chevalier\*

Laboratoire d'Annecy-le-Vieux de Physique des Particules, Université Savoie Mont-Blanc,  
CNRS/IN2P3, F-74941 Annecy-le-Vieux, France

E-mail: [jill.chevalier@lapp.in2p3.fr](mailto:jill.chevalier@lapp.in2p3.fr)

### Max Anton Kastendieck

Universität Hamburg, Institut für Experimentalphysik, Luruper Chaussee 149, D 22761  
Hamburg, Germany

### Frank Rieger

Max-Planck-Institut für Kernphysik, P.O. Box 103980, D 69029 Heidelberg, Germany

### Gilles Maurin

Laboratoire d'Annecy-le-Vieux de Physique des Particules, Université Savoie Mont-Blanc,  
CNRS/IN2P3, F-74941 Annecy-le-Vieux, France

### Jean-Philippe Lenain

LPNHE, Université Pierre et Marie Curie Paris 6, Université Denis Diderot Paris 7,  
CNRS/IN2P3, 4 Place Jussieu, F-75252, Paris Cedex 5, France

### Giovanni Lamanna

Laboratoire d'Annecy-le-Vieux de Physique des Particules, Université Savoie Mont-Blanc,  
CNRS/IN2P3, F-74941 Annecy-le-Vieux, France

Time variability of the photon flux is a known feature of active galactic nuclei (AGN) and in particular of blazars. The high frequency peaked BL Lac (HBL) object PKS 2155-304 is one of the brightest sources in the TeV band and has been monitored regularly with different instruments and in particular with the H.E.S.S. experiment above 200 GeV for more than 11 years. These data together with the observations of other instruments and monitoring programs like SMARTS (optical), *Swift*-XRT/*RXTE*/*XMM-Newton* (X-ray) and *Fermi*-LAT (100 MeV < E < 300 GeV) are used to characterize the variability of this object in the quiescent state over a wide energy range. Variability studies are made by looking at the lognormality of the light curves and at the fractional root mean square (rms) variability  $F_{\text{var}}$  in several energy bands. Lognormality is found in every energy range and the evolution of  $F_{\text{var}}$  with the energy shows a similar increase both in X-rays and in TeV bands.

*The 34th International Cosmic Ray Conference,  
30 July- 6 August, 2015  
The Hague, The Netherlands*

---

\*Speaker.

## 1. Introduction

As many blazars, PKS 2155-304 ( $z = 0.117$ ) features a questioning and interesting behavior: its variability which can be seen over a great energy range, from radio to TeV. Characterizing the variability of blazars is important to study the underlying acceleration processes which are, at the moment, poorly known. Multiwavelength (MWL) data, from experiments and instruments as H.E.S.S., *Fermi*-LAT, *RXTE*, *Swift*-XRT, XMM-Newton and SMARTS are used to study the variability of this object.

Lognormality and intrinsic variability of the MWL light curves are investigated. Lognormal flux variability was first found in galactic accreting sources like X-ray binaries, linking lognormal processes to accretion disks [1]. In a lognormal process, the fluctuations of the flux are on average proportional, or at least correlated, to the flux itself, ruling out additive processes in favor of multiplicative ones. For blazars, a lognormal flux behaviour could as well be indicative of the accretion disk variability's imprint onto the jet [2]. BL Lacertae was the first blazar to show a lognormal behavior in the X-ray [3]. For PKS 2155-304, the first detection of a lognormal behavior was during the flaring state of 2006 in H.E.S.S. data [4].

Studying the evolution of the variability with the energy is one element to further distinguish between different acceleration models.

## 2. Data sets


### 2.1 Very High Energy (VHE): H.E.S.S.

The H.E.S.S. (High Energy Stereoscopic System) experiment is an array of 5 ground-based Cherenkov Telescopes which detect the Cherenkov light emitted by the shower created by VHE  $\gamma$ -rays with energies from  $\sim 100$  GeV to  $> 10$  TeV. The four first telescopes are operated since 2003 (HESS-I phase) and a fifth telescope had been added to the array in 2012 (HESS-II phase).

The present analysis is based on the data taken between MJD 53200 and 56246 using only the HESS-I phase of the array (without the fifth telescope) collected with 3 or 4 telescopes. The flaring state of July/August 2006 from MJD 53943 to 53949 has been removed, in order to have a light curve of the quiescent state of PKS 2155-304 [4]. In total, about 260h of data passed standard quality cuts as defined in [5], with a mean zenith angle of  $21^\circ$  and an energy threshold of 180GeV. The data are analyzed with the Model analysis chain (Model++) [6] between 0.2 and 125 TeV, yielding to a total detection significance  $> 300\sigma$ .

The light curve of nightly fluxes used here is calculated assuming a power-law spectrum  $dN/dE \propto E^{-\Gamma}$ ,  $\Gamma$  being the power law index. However, it has been reported in [4] that PKS 2155-304 shows some spectral variability. Therefore the fluxes are derived with a different fitted spectral index each year to take into account this variability. This leads to 9 different values for  $\Gamma$  (one per year) with a mean of  $3.45 \pm 0.05$  and a standard deviation of 3%.

The resulting quiescent nightly light curve, presented in the first panel of figure 1, has a mean flux of  $6.9 \times 10^{-11} \text{ph.cm}^{-2}.\text{s}^{-1}$ . This analysis procedure has been applied to the same data set using the following energy cuts: 0.2-0.63 and 0.63-125 TeV, the cut at 0.63 TeV being chosen to have the same statistics in each energy band.



**Figure 1:** Light curves from top to bottom : H.E.S.S. in  $\text{cm}^{-2}.\text{s}^{-1}$  from 0.2 to 125 TeV, *Fermi*-LAT in  $\text{cm}^{-2}.\text{s}^{-1}$  from 0.1 to 300 GeV, X-ray in  $\text{erg}.\text{cm}^{-2}.\text{s}^{-1}$  from 2 to 10 keV (*Swift*-XRT in black, *RXTE* in blue and *XMM*-Newton in green) and SMARTS (R band) in Jy.

## 41 2.2 High Energy (HE): *Fermi*-LAT

42 The *Fermi* Large Area Telescope (LAT) [7] on board the *Fermi* satellite is a pair-conversion  
43 telescope designed to detect HE  $\gamma$ -rays in the energy range from  $\sim 20$  MeV to more than 500 GeV.

44 The data for PKS 2155-304 used here goes from MJD 54687 to 56687 and are taken be-  
45 tween 100 MeV and 300 GeV. The detector is described by the P7REP\_SOURCE\_V15 instrument  
46 response functions<sup>1</sup> and the analysis has been performed by the tool *enrico* [8] using *Fermi* Sci-  
47 ence Tools v9r32p5. The prefactor of the diffuse galactic background and the normalisation of the  
48 isotropic diffuse emission are left free to vary in the likelihood fit. All sources from the 3FGL  
49 catalog [9] within  $15^\circ$  around PKS 2155-304 are included in the model. The spectra of PKS 2155-  
50 304 and the nearby FSRQ (Flat Spectrum Radio Quasar) PKS 2149-306 are modeled by a simple  
51 power-law, and their spectral indices and prefactors are left free. All other components are fixed to  
52 the values in the 3FGL catalog.

53 PKS 2155-304 is detected with a significance of  $> 150\sigma$  with an overall spectral index of

<sup>1</sup>[http://fermi.gsfc.nasa.gov/ssc/data/analysis/documentation/Cicerone/Cicerone\\_Data\\_Exploration/](http://fermi.gsfc.nasa.gov/ssc/data/analysis/documentation/Cicerone/Cicerone_Data_Exploration/)

54  $1.83 \pm 0.01$ . The photon counts are integrated into 201 bins of 10 days, resulting in a light curve,  
 55 presented in the second panel of figure 1, with a mean flux of  $1.3 \times 10^{-7} \text{ph.cm}^{-2}.\text{s}^{-1}$ . The same  
 56 analysis procedure has been applied to the same data set using the following energy cuts : 0.1-3,  
 57 3-30 and 30-300 GeV.

### 58 **2.3 X-ray: *RXTE*, *Swift*-XRT and XMM-Newton**

59 *RXTE* (*Rossi X-ray Timing Explorer*) already analyzed light curves are publicly available<sup>2</sup>  
 60 between 2 and 10 keV, and in three other energy bands: 2-4, 4-7 and 7-10 keV.

61 *Swift* X-ray Telescope (XRT) data were analyzed using the package HEASOFT 6.16 between  
 62 2 and 10 keV, and also in four other energy bands: 0.3-2, 2-4, 4-7 and 7-10 keV. The data were recal-  
 63 ibrated using the last update of CALDB and reduced using the standard procedures `xrtpipeline`  
 64 and `xrtproducts`. Caution has been taken to properly account for pile-up effects for corre-  
 65 sponding affected exposures, and spectral fits were performed using `Xspec` 12.8.2 assuming a  
 66 power-law spectrum.

67 XMM-Newton (X-ray Multi Mirror Mission) public EPIC (European Photon Imaging Cam-  
 68 era) data were reduced using the SAS software package, version 14.0, and analyzed following [10]  
 69 between 2 and 10 keV, and also in four other energy bands: 0.3-2, 2-4, 4-7 and 7-10 keV, to match  
 70 *RXTE* and *Swift*-XRT data.

71 *Swift*-XRT and XMM-Newton data have been corrected from galactic absorption with  $n_{\text{H}} =$   
 72  $1.48 \times 10^{20} \text{cm}^{-2}$  [11].

73 In all common energy bins, data of the three instruments are combined to construct an all-over  
 74 X-ray light curve. In order to have an X-ray light curve of the quiescent state of PKS 2155-304,  
 75 all ToO (Target of Opportunities) data were removed. The final light curve, presented in the third  
 76 panel of figure 1, goes from MJD 53501 to 55474 with a mean flux of  $2.7 \times 10^{-11} \text{erg.cm}^{-2}.\text{s}^{-1}$ .

### 77 **2.4 Optical wavelengths : SMARTS**

78 SMARTS (Small and Moderate Aperture Research Telescope System) already analyzed data  
 79 are publicly available<sup>3</sup>. Magnitudes are corrected for the absorption of the galactic foreground  
 80 using [12] before being converted to a spectral flux density using zero magnitude flux values of  
 81 [13]. The light curves are taken in the J, R, V and B bands from MJD 54603 to 56622. For the sake  
 82 of simplicity, only the light curve for the R band is presented in the last panel of figure 1.

## 83 **3. Results**

### 84 **3.1 Lognormality**

85 The lognormal behavior of PKS 2155-304 is investigated by looking at the distribution of  
 86 fluxes and the correlation between the excess rms and the average flux.

87 The flux and log flux distributions of each light curve are fitted by a Gaussian using a  $\chi^2$  fit.  
 88 The results of the fits are summarized in table 1. In all cases, a Gaussian fits the log flux distribution

<sup>2</sup><http://cass.ucsd.edu/rxteagn/>

<sup>3</sup><http://www.astro.yale.edu/smarts/glast/home.php#>

	$\Phi$		$\log \Phi$	
	$\chi^2/\text{d.o.f.}$	Prob	$\chi^2/\text{d.o.f.}$	Prob
H.E.S.S. (> 200GeV)	2.71	$10^{-4}$	0.716	0.75
<i>Fermi</i> -LAT (0.1-300GeV)	2.51	$10^{-3}$	1.36	0.18
X-ray (2-10keV)	6.67	$10^{-10}$	1.36	0.20
SMARTS (B band)	4.35	$10^{-3}$	0.422	0.96
SMARTS (V band)	2.30	$10^{-9}$	0.790	0.65
SMARTS (R band)	5.42	$10^{-9}$	0.823	0.62
SMARTS (J band)	2.33	$10^{-3}$	1.28	0.22

**Table 1:** Values of the  $\chi^2$  and associated probability (probability that the model is good) for the Gaussian fit of the flux and log flux distributions for each light curve.

89 better than the flux distribution, indicating that MLW fluxes of PKS 2155-304 follow a lognormal  
90 distribution.


91 The scatter plots of the excess rms  $\sigma_{\text{XS}} = \sqrt{S^2 - \overline{\sigma_{\text{err}}^2}}$ ,  $S^2$  being the variance and  $\overline{\sigma_{\text{err}}^2}$  the mean  
92 square error of the light curve [15], versus the mean flux  $\overline{\Phi}$  are shown for each light curve in figure  
93 2. Each point is calculated with at least 20 light curve points. To test the rms-flux relation, scatter  
94 plots are fitted by a constant and a linear adjustments. The fit results are summarized in table 2. The  
95 results show more than a  $5\sigma$  preference for the linear fit except for the *Fermi*-LAT data. However  
96 the correlation between  $\sigma_{\text{XS}}$  and  $\overline{\Phi}$  is not expected to be linear, especially when the points are  
97 spread in the plot, so the correlation factor  $\rho$  and the Kendall rank  $\tau$  (which measure the ordering  
98 of the points [14]) are calculated. In all cases  $\rho > 0.80$  and  $\tau > 0.65$  meaning that  $\sigma_{\text{XS}}$  and  $\overline{\Phi}$  show  
99 a strong correlation, leading to the conclusion that the fluctuations of the flux are correlated and/or  
100 proportional to the flux.

	cst	cst+a × x		$\rho$	$\tau$
	$\chi^2/\text{d.o.f.}$	$\chi^2/\text{d.o.f.}$	$\sigma$		
H.E.S.S. (> 200GeV)	4.15	0.773	5.25	$0.84 \pm 0.15$	$0.72 \pm 0.27$
<i>Fermi</i> -LAT (0.1-300GeV)	0.937	0.107	2.75	$0.93 \pm 0.19$	$0.69 \pm 0.25$
X-ray (2-10keV)	78.2	28.9	23	$0.85 \pm 0.02$	$0.67 \pm 0.24$
SMARTS (B band)	4222	892	225	$0.92 \pm 0.03$	$0.68 \pm 0.18$
SMARTS (V band)	4472	810	236	$0.92 \pm 0.02$	$0.75 \pm 0.19$
SMARTS (R band)	7542	946	326	$0.93 \pm 0.02$	$0.74 \pm 0.18$
SMARTS (J band)	3339	398	197	$0.94 \pm 0.03$	$0.80 \pm 0.21$

**Table 2:** Values of the reduced  $\chi^2$  of the constant and linear fits of the scatter plots shown in figure 2 for each light curve, with values for the significance  $\sigma$ , the correlation factor  $\rho$  and the Kendall rank  $\tau$ .

### 101 3.2 Variability energy distribution (VED)

The fractional variability  $F_{\text{var}}$  is used to quantify the variability of the flux taking into account



**Figure 2:** Scatter plot of the excess rms and the average flux for the MWL data. A linear fit is shown in red.

the errors [15]:

$$F_{\text{var}} = \sqrt{\frac{S^2 - \sigma_{\text{err}}^2}{\bar{\Phi}^2}} \pm \sqrt{\left(\sqrt{\frac{1}{2N}} \frac{\sigma_{\text{err}}^2}{\bar{\Phi}^2 F_{\text{var}}}\right)^2 + \left(\sqrt{\frac{\sigma_{\text{err}}^2}{N}} \frac{1}{\bar{\Phi}}\right)^2}$$

102 It is computed for each data set at the different energies presented in section 2. The VED plot  
 103 is shown on figure 3. This figures shows that the SMARTS variability level is somewhat constant  
 104 while a clear increase is visible in the X-ray band. Then the variability is smaller in the *Fermi*-LAT  
 105 range and increases again in the H.E.S.S. range.

#### 106 4. Conclusion

107 The distribution of the flux of the quiescent state of PKS 2155-304 follows a lognormal dis-  
 108 tribution and the variations of the flux are correlated with the flux level itself in all energy band  
 109 probed, indicating that lognormality is an intrinsic characteristic of PKS 2155-304. This results  
 110 have already been shown for the flaring state of 2006 in the TeV band [4]. Others BL Lac type  
 111 sources have present the same lognormal behavior, like BL Lacertae in the X-ray range [3] and  
 112 Markarian 501 at TeV energies [16]. Interestingly non BL Lac objects such as the Seyfert 1 galaxy  
 113 IRAS 13 244-3809 have revealed a similar behavior [17]. Seyfert 1 galaxies have different char-  
 114 acteristics compare to BL Lac: radio quiet (no jet domination), with emission lines and more gas  
 115 surrounding the central engine, indicating again that the lognormal behavior is disk related.




Figure 3: VED of PKS 2155-304.

116 Moreover, the VED of PKS 2155-304 is showing an interesting feature with a similar variabil-  
 117 ity increase in the X-ray and TeV ranges. Such a shape of the VED could help distinguish between  
 118 different acceleration models in blazars as they won't give the same variability signature.

119 **Acknowledgment:** The support of the Namibian authorities and of the University of Namibia in facilitating the  
 120 construction and operation of H.E.S.S. is gratefully acknowledged, as is the support by the German Ministry for Edu-  
 121 cation and Research (BMBF), the Max Planck Society, the German Research Foundation (DFG), the French Ministry  
 122 for Research, the CNRS-IN2P3 and the Astroparticle Interdisciplinary Programme of the CNRS, the U.K. Science and  
 123 Technology Facilities Council (STFC), the IPNP of the Charles University, the Czech Science Foundation, the Polish  
 124 Ministry of Science and Higher Education, the South African Department of Science and Technology and National Re-  
 125 search Foundation, and by the University of Namibia. We appreciate the excellent work of the technical support staff  
 126 in Berlin, Durham, Hamburg, Heidelberg, Palaiseau, Paris, Saclay, and in Namibia in the construction and operation of  
 127 the equipment. This paper has made use of up-to-date SMARTS optical/near-infrared light curves that are available at  
 128 [www.astro.yale.edu/smarts/glast/home.php](http://www.astro.yale.edu/smarts/glast/home.php).

## 129 References

- 130 [1] Uttley, P., McHardy, I. M. (2001). *The flux-dependent amplitude of broadband noise variability in*  
 131 *X-ray binaries and active galaxies*. Monthly Notices of the Royal Astronomical Society, 323(2),  
 132 L26-L30 [arXiv:astro-ph/0103367].
- 133 [2] McHardy, I., *Explaining X-ray variability in blazars*, *Workshop on Blazar Variability across the*  
 134 *Electromagnetic Spectrum* (2008).
- 135 [3] Giebels, B., & Degrange, B. (2009). *Lognormal variability in BL Lacertae*. Astronomy &  
 136 Astrophysics, 503(3), 797-799 [arXiv:0907.2425].
- 137 [4] Abramowski, A., Acero, F., Aharonian, F., Akhperjanian, A. G., Anton, G., De Almeida, U. B., ... &  
 138 Fegan, S. (2010). *VHE  $\gamma$ -ray emission of PKS 2155-304: spectral and temporal variability*. Astronomy  
 139 & Astrophysics, 520, A83 [arXiv:1005.3702].



- 140 [5] Aharonian, F., Akhperjanian, A. G., Bazer-Bachi, A. R., Beilicke, M., Benbow, W., Berge, D., ... & De  
141 Jager, O. C. (2006). *Observations of the Crab nebula with HESS*. *Astronomy & Astrophysics*, 457(3),  
142 899-915 [arXiv:astro-ph/0607333].
- 143 [6] De Naurois, M., Rolland, L. (2009). *A high performance likelihood reconstruction of  $\gamma$ -rays for  
144 imaging atmospheric Cherenkov telescopes*. *Astroparticle Physics*, 32(5), 231-252  
145 [arXiv:0907.2610].
- 146 [7] Atwood, W. B., Abdo, A. A., Ackermann, M., Althouse, W., Anderson, B., Axelsson, M., ... Ciprini, S.  
147 (2009). *The Large Area Telescope on the Fermi gamma-ray space telescope mission*. *The Astrophysical  
148 Journal*, 697(2), 1071 [arXiv:0902.1089].
- 149 [8] Sanchez, D. A., & Deil, C. (2013), *Enrico : a Python package to simplify Fermi-LAT analysis*  
150 [arXiv:1307.4534].
- 151 [9] Acero, F., Ackermann, M., Ajello, M., Albert, A., Atwood, W. B., Axelsson, M., ... Di Venere, L.  
152 (2015). *Fermi Large Area Telescope Third Source Catalog*. *The Astrophysical Journal Supplement  
153 Series*, 218(2), 23 [arXiv:1501.02003].
- 154 [10] Tatischeff, V., Decourchelle, A., & Maurin, G. (2012). *Nonthermal X-rays from low-energy cosmic  
155 rays: application to the 6.4 keV line emission from the Arches cluster region*. *Astronomy &  
156 Astrophysics*, 546, A88 [arXiv:1210.2108].
- 157 [11] Kalberla, P.M.W., Burton, W.B., Hartmann, Dap, Arnal, E.M., Bajaja, E., Morras, R., Pöppel, W.G.L.  
158 (2005) *The Leiden/Argentine/Bonn (LAB) Survey of Galactic HI, Final data release of the combined  
159 LDS and IAR surveys with improved stray-radiation corrections*, *AA*, 440, 775  
160 [arXiv:astro-ph/0504140]
- 161 [12] Schlafly, E. F., & Finkbeiner, D. P. (2011). *Measuring reddening with Sloan Digital Sky Survey stellar  
162 spectra and recalibrating SFD*. *The Astrophysical Journal*, 737(2), 103 [arXiv:1012.4804].
- 163 [13] Cohen, M., Walker, R. G., Barlow, M. J., Deacon, J. R. (1992). *Spectral irradiance calibration in the  
164 infrared. I-Ground-based and IRAS broadband calibrations*. *The Astronomical Journal*, 104, 1650-1657.
- 165 [14] Gleissner, T., Wilms, J., Pottschmidt, K., Uttley, P., Nowak, M. A., & Staubert, R. (2004). *Long term  
166 variability of Cyg X-1-II. The rms-flux relation*. *Astronomy & Astrophysics*, 414(3), 1091-1104  
167 [arXiv:astro-ph/0311039].
- 168 [15] Vaughan, S., Edelson, R., Warwick, R. S., & Uttley, P. (2003). *On characterizing the variability  
169 properties of X-ray light curves from active galaxies*. *Monthly Notices of the Royal Astronomical  
170 Society*, 345(4), 1271-1284 [arXiv:astro-ph/0307420].
- 171 [16] Chakraborty, N., Cologna, G., Rieger, F., et al. for the H.E.S.S. Collaboration, *Rapid variability at  
172 very high energy in Mrk 501, Proceedings of the 34th International Cosmic Ray Conference (ICRC  
173 2015), The Hague (Netherlands) (2015)*.
- 174 [17] Gaskell, C. M. (2004). *Lognormal X-ray flux variations in an extreme narrow-line Seyfert 1 galaxy*.  
175 *The Astrophysical Journal Letters*, 612(1), L21.