

Measurement of the Radiation Energy in the Radio Signal of Extensive Air Showers as a Universal Estimator of Cosmic-Ray Energy

Alexander Aab, Pedro Abreu, Marco Aglietta, Eun-Joo Ahn, Imen Al Samarai, Ivone Albuquerque, Ingomar Allekotte, Patrick Allison, Alejandro Almela, Jesus Alvarez Castillo, et al.

► **To cite this version:**

Alexander Aab, Pedro Abreu, Marco Aglietta, Eun-Joo Ahn, Imen Al Samarai, et al.. Measurement of the Radiation Energy in the Radio Signal of Extensive Air Showers as a Universal Estimator of Cosmic-Ray Energy. *Physical Review Letters*, American Physical Society, 2016, 116, pp.241101. 10.1103/PhysRevLett.116.241101 . in2p3-01313571

HAL Id: in2p3-01313571

<http://hal.in2p3.fr/in2p3-01313571>

Submitted on 6 Aug 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Measurement of the Radiation Energy in the Radio Signal of Extensive Air Showers as a Universal Estimator of Cosmic-Ray Energy

A. Aab,¹ P. Abreu,² M. Aglietta,³ E. J. Ahn,⁴ I. Al Samarai,⁵ I. F. M. Albuquerque,⁶ I. Allekotte,⁷ P. Allison,⁸ A. Almela,^{9,10} J. Alvarez Castillo,¹¹ J. Alvarez-Muñiz,¹² R. Alves Batista,¹³ M. Ambrosio,¹⁴ A. Aminaei,¹⁵ G. A. Anastasi,¹⁶ L. Anchordoqui,¹⁷ S. Andringa,² C. Aramo,¹⁴ F. Arqueros,¹⁸ N. Arsene,¹⁹ H. Asorey,^{7,20} P. Assis,² J. Aublin,²¹ G. Avila,²² N. Awai,²³ A. M. Badescu,²⁴ C. Baus,²⁵ J. J. Beatty,⁸ K. H. Becker,²⁶ J. A. Bellido,²⁷ C. Berat,²⁸ M. E. Bertaina,³ X. Bertou,⁷ P. L. Biermann,²⁹ P. Billoir,²¹ S. G. Blaess,²⁷ A. Blanco,² M. Blanco,²¹ J. Blazek,³⁰ C. Bleve,³¹ H. Blümer,^{25,32} M. Boháčová,³⁰ D. Boncioli,³³ C. Bonifazi,³⁴ N. Borodai,³⁵ J. Brack,³⁶ I. Brancus,³⁷ T. Bretz,³⁸ A. Bridgeman,³² P. Brogueira,² P. Buchholz,¹ A. Bueno,³⁹ S. Buitink,¹⁵ M. Buscemi,¹⁴ K. S. Caballero-Mora,⁴⁰ B. Caccianiga,⁴¹ L. Caccianiga,²¹ M. Candusso,⁴² L. Caramete,⁴³ R. Caruso,¹⁶ A. Castellina,³ G. Cataldi,³¹ L. Cazon,² R. Cester,⁴⁴ A. G. Chavez,⁴⁵ A. Chiavassa,³ J. A. Chinellato,⁴⁶ J. Chudoba,³⁰ M. Cilmo,¹⁴ R. W. Clay,²⁷ G. Cocciolo,³¹ R. Colalillo,¹⁴ A. Coleman,⁴⁷ L. Collica,⁴¹ M. R. Coluccia,³¹ R. Conceição,² F. Contreras,⁴⁸ M. J. Cooper,²⁷ A. Cordier,⁴⁹ S. Coutu,⁴⁷ C. E. Covault,⁵⁰ J. Cronin,⁵¹ R. Dallier,^{52,53} B. Daniel,⁴⁶ S. Dasso,^{54,55} K. Daumiller,³² B. R. Dawson,²⁷ R. M. de Almeida,⁵⁶ S. J. de Jong,^{15,57} G. De Mauro,¹⁵ J. R. T. de Mello Neto,³⁴ I. De Mitri,³¹ J. de Oliveira,⁵⁶ V. de Souza,⁵⁸ L. del Peral,⁵⁹ O. Deligny,⁵ N. Dhital,⁶⁰ C. Di Giulio,⁴² A. Di Matteo,⁶¹ J. C. Diaz,⁶⁰ M. L. Díaz Castro,⁴⁶ F. Diogo,² C. Dobrigkeit,⁴⁶ W. Docters,⁶² J. C. D'Olivo,¹¹ A. Dorofeev,³⁶ Q. Dorosti Hasankiadeh,³² R. C. dos Anjos,⁵⁸ M. T. Dova,⁶³ J. Ebr,³⁰ R. Engel,³² M. Erdmann,³⁸ M. Erfani,¹ C. O. Escobar,^{4,46} J. Espadanal,² A. Etchegoyen,^{10,9} H. Falcke,^{15,64,57} K. Fang,⁵¹ G. Farrar,²³ A. C. Fauth,⁴⁶ N. Fazzini,⁴ A. P. Ferguson,⁵⁰ B. Fick,⁶⁰ J. M. Figueira,¹⁰ A. Filevich,¹⁰ A. Filipčič,^{65,66} O. Fratu,²⁴ M. M. Freire,⁶⁷ T. Fujii,⁵¹ B. García,⁶⁸ D. Garcia-Gamez,⁴⁹ D. Garcia-Pinto,¹⁸ F. Gate,⁵² H. Gemmeke,⁶⁹ A. Gherghel-Lascu,³⁷ P. L. Ghia,²¹ U. Giaccari,³⁴ M. Giammarchi,⁴¹ M. Giller,⁷⁰ D. Glas,⁷⁰ C. Glaser,³⁸ H. Glass,⁴ G. Golup,⁷ M. Gómez Berisso,⁷ P. F. Gómez Vitale,²² N. González,¹⁰ B. Gookin,³⁶ J. Gordon,⁸ A. Gorgi,³ P. Gorham,⁷¹ P. Gouffon,⁶ N. Griffith,⁸ A. F. Grillo,³³ T. D. Grubb,²⁷ F. Guarino,¹⁴ G. P. Guedes,⁷² M. R. Hampel,¹⁰ P. Hansen,⁶³ D. Harari,⁷ T. A. Harrison,²⁷ S. Hartmann,³⁸ J. L. Harton,³⁶ A. Haungs,³² T. Hebbeker,³⁸ D. Heck,³² P. Heimann,¹ A. E. Herve,³² G. C. Hill,²⁷ C. Hojvat,⁴ N. Hollon,⁵¹ E. Holt,³² P. Homola,²⁶ J. R. Hörandel,^{15,57} P. Horvath,⁷³ M. Hrabovský,^{73,30} D. Huber,²⁵ T. Huege,³² A. Insolia,¹⁶ P. G. Isar,⁴³ I. Jandt,²⁶ S. Jansen,^{15,57} C. Jarne,⁶³ J. A. Johnsen,⁷⁴ M. Josebachuili,¹⁰ A. Kääpä,²⁶ O. Kambeitz,²⁵ K. H. Kampert,²⁶ P. Kasper,⁴ I. Katkov,²⁵ B. Keilhauer,³² E. Kemp,⁴⁶ R. M. Kieckhafer,⁶⁰ H. O. Klages,³² M. Kleifges,⁶⁹ J. Kleinfeller,⁴⁸ R. Krause,³⁸ N. Krohm,²⁶ D. Kuempel,³⁸ G. Kuvec Mezek,⁶⁶ N. Kunka,⁶⁹ A. W. Kuotb Awad,³² D. LaHurd,⁵⁰ L. Latronico,³ R. Lauer,⁷⁵ M. Lauscher,³⁸ P. Lautridou,⁵² S. Le Coz,²⁸ D. Lebrun,²⁸ P. Lebrun,⁴ M. A. Leigui de Oliveira,⁷⁶ A. Letessier-Selvon,²¹ I. Lhenry-Yvon,⁵ K. Link,²⁵ L. Lopes,² R. López,⁷⁷ A. López Casado,¹² K. Louedec,²⁸ A. Lucero,¹⁰ M. Malacari,²⁷ M. Mallamaci,⁴¹ J. Maller,⁵² D. Mandat,³⁰ P. Mantsch,⁴ A. G. Mariazzi,⁶³ V. Marin,⁵² I. C. Mariş,³⁹ G. Marsella,³¹ D. Martello,³¹ H. Martinez,⁷⁸ O. Martínez Bravo,⁷⁷ D. Martraire,⁵ J. J. Masías Meza,⁵⁵ H. J. Mathes,³² S. Mathys,²⁶ J. Matthews,⁷⁹ J. A. J. Matthews,⁷⁵ G. Matthiae,⁴² D. Maurizio,⁸⁰ E. Mayotte,⁷⁴ P. O. Mazur,⁴ C. Medina,⁷⁴ G. Medina-Tanco,¹¹ R. Meissner,³⁸ V. B. B. Mello,³⁴ D. Melo,¹⁰ A. Meshnikov,⁶⁹ S. Messina,⁶² M. I. Micheletti,⁶⁷ L. Middendorf,³⁸ I. A. Minaya,¹⁸ L. Miramonti,⁴¹ B. Mitrica,³⁷ L. Molina-Bueno,³⁹ S. Mollerach,⁷ F. Montanet,²⁸ C. Morello,³ M. Mostafá,⁴⁷ C. A. Moura,⁷⁶ M. A. Muller,^{46,81} G. Müller,³⁸ S. Müller,³² S. Navas,³⁹ P. Necesar,³⁰ L. Nellen,¹¹ A. Nelles,^{15,57} J. Neuser,²⁶ P. H. Nguyen,²⁷ M. Niculescu-Oglinzanu,³⁷ M. Niechciol,¹ L. Niemiets,²⁶ T. Niggemann,³⁸ D. Nitz,⁶⁰ D. Nosek,⁸² V. Novotny,⁸² L. Nožka,⁷³ L. A. Núñez,²⁰ L. Ochilo,¹ F. Oikonomou,⁴⁷ A. Olinto,⁵¹ N. Pacheco,⁵⁹ D. Pakk Selmi-Dei,⁴⁶ M. Palatka,³⁰ J. Pallotta,⁸³ P. Papenbreer,²⁶ G. Parente,¹² A. Parra,⁷⁷ T. Paul,^{17,84} M. Pech,³⁰ J. Pękala,³⁵ R. Pelayo,⁸⁵ I. M. Pepe,⁸⁶ L. Perrone,³¹ E. Petermann,⁸⁷ C. Peters,³⁸ S. Petrera,^{61,88} Y. Petrov,³⁶ J. Phuntsok,⁴⁷ R. Piegaia,⁵⁵ T. Pierog,³² P. Pieroni,⁵⁵ M. Pimenta,² V. Pirronello,¹⁶ M. Platino,¹⁰ M. Plum,³⁸ A. Porcelli,³² C. Porowski,³⁵ R. R. Prado,⁵⁸ P. Privitera,⁵¹ M. Prouza,³⁰ E. J. Quel,⁸³ S. Querchfeld,²⁶ S. Quinn,⁵⁰ J. Rautenberg,²⁶ O. Ravel,⁵² D. Ravignani,¹⁰ D. Reinert,³⁸ B. Revenu,⁵² J. Ridky,³⁰ M. Risse,¹ P. Ristori,⁸³ V. Rizi,⁶¹ W. Rodrigues de Carvalho,¹² J. Rodriguez Rojo,⁴⁸ M. D. Rodríguez-Frías,⁵⁹ D. Rogozin,³² J. Rosado,¹⁸ M. Roth,³² E. Roulet,⁷ A. C. Rovero,⁵⁴ S. J. Saffi,²⁷ A. Saftoiu,³⁷ H. Salazar,⁷⁷ A. Saleh,⁶⁶ F. Salesa Greus,⁴⁷ G. Salina,⁴² J. D. Sanabria Gomez,²⁰ F. Sánchez,¹⁰ P. Sanchez-Lucas,³⁹ E. Santos,⁴⁶ E. M. Santos,⁶ F. Sarazin,⁷⁴ B. Sarkar,²⁶ R. Sarmiento,² C. Sarmiento-Cano,²⁰ R. Sato,⁴⁸ C. Scarso,⁴⁸ M. Schauer,²⁶ V. Scherini,³¹ H. Schieler,³² D. Schmidt,³² O. Scholten,^{62,89} H. Schoorlemmer,⁷¹ P. Schovánek,³⁰ F. G. Schröder,³² A. Schulz,³² J. Schulz,¹⁵ J. Schumacher,³⁸ S. J. Sciutto,⁶³ A. Segreto,⁹⁰ M. Settimo,²¹ A. Shadkam,⁷⁹ R. C. Shellard,⁸⁰ G. Sigl,¹³ O. Sima,¹⁹ A. Śmiałkowski,⁷⁰ R. Šmída,³² G. R. Snow,⁸⁷ P. Sommers,⁴⁷ S. Sonntag,¹

J. Sorokin,²⁷ R. Squartini,⁴⁸ Y. N. Srivastava,⁸⁴ D. Stanca,³⁷ S. Stanič,⁶⁶ J. Stapleton,⁸ J. Stasielak,³⁵ M. Stephan,³⁸ A. Stutz,²⁸ F. Suarez,^{10,9} M. Suarez Durán,²⁰ T. Suomijärvi,⁵ A. D. Supanitsky,⁵⁴ M. S. Sutherland,⁸ J. Swain,⁸⁴ Z. Szadkowski,⁷⁰ O. A. Tabora,⁷ A. Tapia,¹⁰ A. Tepe,¹ V. M. Theodoro,⁴⁶ C. Timmermans,^{57,15} C. J. Todero Peixoto,⁹¹ G. Toma,³⁷ L. Tomankova,³² B. Tomé,² A. Tonachini,⁴⁴ G. Torralba Elipe,¹² D. Torres Machado,³⁴ P. Travnicek,³⁰ M. Trini,⁶⁶ R. Ulrich,³² M. Unger,^{23,32} M. Urban,³⁸ J. F. Valdés Galicia,¹¹ I. Valiño,¹² L. Valore,¹⁴ G. van Aar,¹⁵ P. van Bodegom,²⁷ A. M. van den Berg,⁶² S. van Velzen,¹⁵ A. van Vliet,¹³ E. Varela,⁷⁷ B. Vargas Cárdenas,¹¹ G. Varner,⁷¹ R. Vasquez,³⁴ J. R. Vázquez,¹⁸ R. A. Vázquez,¹² D. Veberič,³² V. Verzi,⁴² J. Vicha,³⁰ M. Videla,¹⁰ L. Villaseñor,⁴⁵ B. Vlcek,⁵⁹ S. Vorobiov,⁶⁶ H. Wahlberg,⁶³ O. Wainberg,^{10,9} D. Walz,³⁸ A. A. Watson,⁹² M. Weber,⁶⁹ K. Weidenhaupt,³⁸ A. Weindl,³² C. Welling,³⁸ F. Werner,²⁵ A. Widom,⁸⁴ L. Wiencke,⁷⁴ H. Wilczyński,³⁵ T. Winchen,²⁶ D. Wittkowski,²⁶ B. Wundheiler,¹⁰ S. Wykes,¹⁵ L. Yang,⁶⁶ T. Yapici,⁶⁰ A. Yushkov,¹ E. Zas,¹² D. Zavrtanik,^{66,65} M. Zavrtanik,^{65,66} A. Zepeda,⁷⁸ B. Zimmermann,⁶⁹ M. Ziolkowski,¹ and F. Zuccarello¹⁶

(Pierre Auger Collaboration)*

¹Universität Siegen, Fachbereich 7 Physik - Experimentelle Teilchenphysik, Siegen, Germany

²Laboratório de Instrumentação e Física Experimental de Partículas - LIP and Instituto Superior Técnico - IST, Universidade de Lisboa - UL, Lisboa, Portugal

³Osservatorio Astrofisico di Torino (INAF), Università di Torino and Sezione INFN, Torino, Italy

⁴Fermilab, Batavia, Illinois, USA

⁵Institut de Physique Nucléaire d'Orsay (IPNO), Université Paris 11, CNRS-IN2P3, Orsay, France

⁶Universidade de São Paulo, Instituto de Física, São Paulo, São Paulo, Brazil

⁷Centro Atómico Bariloche and Instituto Balseiro (CNEA-UNCuyo-CONICET), San Carlos de Bariloche, Argentina

⁸Ohio State University, Columbus, Ohio, USA

⁹Universidad Tecnológica Nacional - Facultad Regional Buenos Aires, Buenos Aires, Argentina

¹⁰Instituto de Tecnologías en Detección y Astropartículas (CNEA, CONICET, UNSAM), Buenos Aires, Argentina

¹¹Universidad Nacional Autónoma de México, México, Distrito Federal, Mexico

¹²Universidad de Santiago de Compostela, Santiago de Compostela, Spain

¹³Universität Hamburg, II. Institut für Theoretische Physik, Hamburg, Germany

¹⁴Università di Napoli "Federico II" and Sezione INFN, Napoli, Italy

¹⁵IMAPP, Radboud University Nijmegen, Nijmegen, Netherlands

¹⁶Università di Catania and Sezione INFN, Catania, Italy

¹⁷Department of Physics and Astronomy, Lehman College, City University of New York, Bronx, New York, USA

¹⁸Universidad Complutense de Madrid, Madrid, Spain

¹⁹University of Bucharest, Physics Department, Bucharest, Romania

²⁰Universidad Industrial de Santander, Bucaramanga, Colombia

²¹Laboratoire de Physique Nucléaire et de Hautes Energies (LPNHE), Universités Paris 6 et Paris 7, CNRS-IN2P3, Paris, France

²²Observatorio Pierre Auger and Comisión Nacional de Energía Atómica, Malargüe, Argentina

²³New York University, New York, New York, USA

²⁴University Politehnica of Bucharest, Bucharest, Romania

²⁵Karlsruhe Institute of Technology, Institut für Experimentelle Kernphysik (IEKP), Karlsruhe, Germany

²⁶Bergische Universität Wuppertal, Fachbereich C - Physik, Wuppertal, Germany

²⁷University of Adelaide, Adelaide, South Australia, Australia

²⁸Laboratoire de Physique Subatomique et de Cosmologie (LPSC), Université Grenoble-Alpes, CNRS/IN2P3, Grenoble, France

²⁹Max-Planck-Institut für Radioastronomie, Bonn, Germany

³⁰Institute of Physics of the Academy of Sciences of the Czech Republic, Prague, Czech Republic

³¹Dipartimento di Matematica e Fisica "E. De Giorgi" dell'Università del Salento and Sezione INFN, Lecce, Italy

³²Karlsruhe Institute of Technology, Institut für Kernphysik, Karlsruhe, Germany

³³INFN, Laboratori Nazionali del Gran Sasso, Assergi (L'Aquila), Italy

³⁴Universidade Federal do Rio de Janeiro, Instituto de Física, Rio de Janeiro, Rio de Janeiro, Brazil

³⁵Institute of Nuclear Physics PAN, Krakow, Poland

³⁶Colorado State University, Fort Collins, Colorado, USA

³⁷"Horia Hulubei" National Institute for Physics and Nuclear Engineering, Bucharest-Magurele, Romania

³⁸RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany

³⁹Universidad de Granada and C.A.F.P.E., Granada, Spain

⁴⁰Universidad Autónoma de Chiapas, Tuxtla Gutiérrez, Chiapas, Mexico

⁴¹Università di Milano and Sezione INFN, Milan, Italy

⁴²Università di Roma II "Tor Vergata" and Sezione INFN, Roma, Italy

⁴³Institute of Space Science, Bucharest-Magurele, Romania

- ⁴⁴Università di Torino and Sezione INFN, Torino, Italy
- ⁴⁵Universidad Michoacana de San Nicolás de Hidalgo, Morelia, Michoacán, Mexico
- ⁴⁶Universidade Estadual de Campinas, IFGW, Campinas, São Paulo, Brazil
- ⁴⁷Pennsylvania State University, University Park, Pennsylvania, USA
- ⁴⁸Observatorio Pierre Auger, Malargüe, Argentina
- ⁴⁹Laboratoire de l'Accélérateur Linéaire (LAL), Université Paris 11, CNRS-IN2P3, Orsay, France
- ⁵⁰Case Western Reserve University, Cleveland, Ohio, USA
- ⁵¹University of Chicago, Enrico Fermi Institute, Chicago, Illinois, USA
- ⁵²SUBATECH, école des Mines de Nantes, CNRS-IN2P3, Université de Nantes, Nantes, France
- ⁵³Station de Radioastronomie de Nançay, Observatoire de Paris, CNRS/INSU, Nançay, France
- ⁵⁴Instituto de Astronomía y Física del Espacio (IAFE, CONICET-UBA), Buenos Aires, Argentina
- ⁵⁵Departamento de Física, FCEyN, Universidad de Buenos Aires and CONICET, Buenos Aires, Argentina
- ⁵⁶Universidade Federal Fluminense, EEIMVR, Volta Redonda, Rio de Janeiro, Brazil
- ⁵⁷Nikhef, Science Park, Amsterdam, Netherlands
- ⁵⁸Universidade de São Paulo, Instituto de Física de São Carlos, São Carlos, São Paulo, Brazil
- ⁵⁹Universidad de Alcalá, Alcalá de Henares, Madrid, Spain
- ⁶⁰Michigan Technological University, Houghton, Michigan, USA
- ⁶¹Dipartimento di Scienze Fisiche e Chimiche dell'Università dell'Aquila and INFN, L'Aquila, Italy
- ⁶²KVI - Center for Advanced Radiation Technology, University of Groningen, Groningen, Netherlands
- ⁶³IFLP, Universidad Nacional de La Plata and CONICET, La Plata, Argentina
- ⁶⁴ASTRON, Dwingeloo, Netherlands
- ⁶⁵Experimental Particle Physics Department, J. Stefan Institute, Ljubljana, Slovenia
- ⁶⁶Laboratory for Astroparticle Physics, University of Nova Gorica, Nova Gorica, Slovenia
- ⁶⁷Instituto de Física de Rosario (IFIR) - CONICET/U.N.R. and Facultad de Ciencias Bioquímicas y Farmacéuticas U.N.R., Rosario, Argentina
- ⁶⁸Instituto de Tecnologías en Detección y Astropartículas (CNEA, CONICET, UNSAM), and Universidad Tecnológica Nacional - Facultad Regional Mendoza (CONICET/CNEA), Mendoza, Argentina
- ⁶⁹Karlsruhe Institute of Technology, Institut für Prozessdatenverarbeitung und Elektronik, Karlsruhe, Germany
- ⁷⁰University of Łódź, Łódź, Poland
- ⁷¹University of Hawaii, Honolulu, Hawaii, USA
- ⁷²Universidade Estadual de Feira de Santana, Feira de Santana, Brazil
- ⁷³Palacky University, RCPTM, Olomouc, Czech Republic
- ⁷⁴Colorado School of Mines, Golden, Colorado, USA
- ⁷⁵University of New Mexico, Albuquerque, New Mexico, USA
- ⁷⁶Universidade Federal do ABC, Santo André, São Paulo, Brazil
- ⁷⁷Benemérita Universidad Autónoma de Puebla, Puebla, Mexico
- ⁷⁸Centro de Investigación y de Estudios Avanzados del IPN (CINVESTAV), México, Distrito Federal, Mexico
- ⁷⁹Louisiana State University, Baton Rouge, Louisiana, USA
- ⁸⁰Centro Brasileiro de Pesquisas Físicas, Rio de Janeiro, Rio de Janeiro, Brazil
- ⁸¹Universidade Federal de Pelotas, Pelotas, Rio Grande do Sul, Brazil
- ⁸²Charles University, Faculty of Mathematics and Physics, Institute of Particle and Nuclear Physics, Prague, Czech Republic
- ⁸³Centro de Investigaciones en Láseres y Aplicaciones, CITEDEF and CONICET, Villa Martelli, Argentina
- ⁸⁴Northeastern University, Boston, Massachusetts, USA
- ⁸⁵Unidad Profesional Interdisciplinaria en Ingeniería y Tecnologías Avanzadas del Instituto Politécnico Nacional (UPIITA- IPN), México, Distrito Federal, Mexico
- ⁸⁶Universidade Federal da Bahia, Salvador, BA, Brazil
- ⁸⁷University of Nebraska, Lincoln, Nebraska, USA
- ⁸⁸Gran Sasso Science Institute (INFN), L'Aquila, Italy
- ⁸⁹Vrije Universiteit Brussel, Brussels, Belgium
- ⁹⁰Istituto di Astrofisica Spaziale e Fisica Cosmica di Palermo (INAF), Palermo, Italy
- ⁹¹Universidade de São Paulo, Escola de Engenharia de Lorena, Lorena, São Paulo, Brazil
- ⁹²School of Physics and Astronomy, University of Leeds, Leeds, United Kingdom

(Received 27 August 2015; revised manuscript received 4 April 2016; published 14 June 2016)

We measure the energy emitted by extensive air showers in the form of radio emission in the frequency range from 30 to 80 MHz. Exploiting the accurate energy scale of the Pierre Auger Observatory, we obtain a radiation energy of $15.8 \pm 0.7(\text{stat}) \pm 6.7(\text{syst})$ MeV for cosmic rays with an energy of 1 EeV arriving perpendicularly to a geomagnetic field of 0.24 G, scaling quadratically with the cosmic-ray energy. A comparison with predictions from state-of-the-art first-principles calculations shows agreement with our measurement. The radiation energy provides direct access to the calorimetric energy in the

electromagnetic cascade of extensive air showers. Comparison with our result thus allows the direct calibration of any cosmic-ray radio detector against the well-established energy scale of the Pierre Auger Observatory.

DOI: [10.1103/PhysRevLett.116.241101](https://doi.org/10.1103/PhysRevLett.116.241101)

In this work, we address one of the most important challenges in cosmic-ray physics: the accurate determination of the absolute energy scale of cosmic rays. Measurements with surface particle detector arrays rely on assumptions about cosmic-ray composition and on extrapolations of our knowledge about hadronic interactions to energies beyond the reach of the Large Hadron Collider. Consequently, their determination of the absolute cosmic-ray energy suffers from significant uncertainties [1]. Fluorescence detectors measure the calorimetric energy in the electromagnetic cascade of air showers, which allows an accurate determination of the energy of the primary particle [2]. However, fluorescence light detection is possible only at sites with good atmospheric conditions, and precise quantification of scattering and absorption of fluorescence light under changing atmospheric conditions requires extensive atmospheric monitoring efforts [3–6].

An attractive option to determine the energy scale of cosmic-ray particles is given by the detection of radio signals. Radio detection of extensive air showers can be performed at any site not overwhelmed by anthropogenic radio signals, requiring only detector arrays of moderate size and complexity. It has been known since the 1960s that air showers emit measurable radio pulses [7]. The physics of the radio emission from extensive air showers is by now well understood (see [8] for an overview). The radiation dominantly arises from geomagnetically induced, time-varying transverse currents [9,10] and is strongly forward beamed in a cone of a few degree opening angle due to the relativistic speed of the emitting particles. The atmosphere is transparent for radio waves at the relevant frequencies; i.e., scattering and absorption are negligible. As the emission is generally coherent at frequencies below 100 MHz, the amplitude of the electric field scales linearly with the number of electrons and positrons in the air-shower cascade, which in turn scales linearly with the primary cosmic-ray energy.

Several analyses exploiting this calorimetric property of the radio emission for the determination of the energy of cosmic-ray particles have previously been published [11–14]. All of these approaches used the radio-signal strength at a characteristic lateral distance from the shower axis as an estimator for the cosmic-ray energy. While this method has long been known to provide good precision [15], it has the marked disadvantage that the corresponding energy estimator cannot be directly compared across different experiments. Asymmetries arising from the charge-excess contribution [16–18] can be corrected for, and the air-shower zenith angle can be normalized out. The systematic influence of the observation altitude on the

lateral signal distribution, however, poses a fundamental problem for such comparisons. In a simulation study, we have quantified the difference between radio amplitudes at the characteristic lateral distance measured for the same showers at sea level (altitude of LOFAR [19]) and at 1560 m above sea level (altitude of the radio detector array of the Pierre Auger Observatory [20]). We observe differences between -11% and $+23\%$ with an average deviation of 11% . These deviations in the measured amplitude arise from the fact that the lateral radio-signal distribution flattens systematically with the increasing distance of the radio antennas to the air-shower maximum. Furthermore, the optimal lateral distance at which to make the measurement also varies with observation altitude [21]. While absolute values for the amplitudes measured at a characteristic lateral distance as a function of cosmic-ray energy have been published by several experiments [13,14,22], no direct comparison between the energy scales of these cosmic-ray radio detectors has therefore been performed to date. (Most experiments obtain their energy scale based on surface detector arrays and thus incur uncertainties from hadronic interaction models).

Here, we make an important conceptual step forward in using radio signals from extensive air showers for the absolute calibration of the energy scale of cosmic-ray detectors. We use the total energy radiated by extensive air showers in the form of radio emission in the frequency range from 30 to 80 MHz, hereafter called *radiation energy*, as an estimator of the cosmic-ray energy. Because of conservation of energy and the absence of absorption in the atmosphere, the radiation energy measured at different observation altitudes is virtually identical. In the above-mentioned simulation study, the radiation energy was shown to vary less than 0.5% between an observation altitude of 1560 m above sea level and sea level itself. (This scatter arises from slight clipping effects of the air-shower evolution at an observation altitude of 1560 m above sea level and from statistical uncertainties in the determination of the radiation energy from the simulated radio-emission footprint.) The radiation energy directly reflects the calorimetric energy in the electromagnetic cascade of an extensive air shower, akin to an integral of the Gaisser-Hillas profile measured with fluorescence detectors. It constitutes a universal, well-defined quantity that can be measured with radio detectors worldwide and can thus be compared directly between different experiments, as well as with theoretical predictions.

In this work, we measure the absolute value of the radiation energy with the Auger Engineering Radio Array

(AERA) [23], an array of radio detectors in the Pierre Auger Observatory [20]. We then cross-calibrate our measurement with data taken with the baseline detectors of the Auger Observatory. The Observatory includes an array of water-Cherenkov particle detectors covering an area of 3000 km². The atmosphere above the surface detector is monitored by fluorescence telescopes which provide an absolute calibration of the cosmic-ray energy scale [24] with a systematic uncertainty of 16% at 10^{17.5} eV and 14% at energies $\geq 10^{18}$ eV [2], reflecting the state of the art in the determination of the absolute energy scale achieved to date. We thus use the accurate calibration of the energy scale of the Pierre Auger Observatory to relate the radiation energy to the cosmic-ray energy. The radiation energy can in turn be used to calibrate cosmic-ray radio detectors worldwide against the Auger energy scale. Finally, we provide a first comparison with predictions from first-principles calculations.

Details of the analysis presented here can be found in an accompanying publication [25].

The energy content in the radio signal.—With the radio antennas of AERA, we continuously sample voltage traces arising from the measurement of the local electric field with antennas oriented along the geomagnetic north-south and east-west directions. Upon a trigger from coincident radio pulses or external trigger information from other Auger detectors, the voltage traces are read out for off-line analysis [26]. From these voltage traces, we reconstruct the electric field vector at the location of each radio detector as a function of time. Detector effects are carefully unfolded [25]. The uncertainty of the electric field amplitude between different measurements is dominated by temperature variations (4%) and uncertainties of the antenna response pattern (5%) and amounts to a total of 6.4%. The uncertainty of the absolute amplitude scale is dominated by the antenna response (12.5% [22,27]) and the analog signal chain (6%) and amounts to a total of 14%.

After digital processing (involving noise cleaning, up-sampling, and enveloping), we identify radio pulses exceeding a suitable signal-to-noise threshold. We calculate the instantaneous Poynting flux at each radio detector and integrate it over a time window of 200 ns which is centered on the pulse maximum. The contribution of noise to the integral is estimated from data recorded before the arrival of the extensive air shower and is subtracted from the integrated signal. The result of the time integration corresponds to the energy deposited per area by air-shower radio signals at the locations of the individual radio detectors. We measure this *energy fluence* in units of eV/m². Typical values are in the range of dozens of eV/m². The energy of a photon at our center-of-band frequency of 55 MHz corresponds to 2.27×10^{-7} eV. The number of received photons is thus very high, illustrating that uncertainties from photon statistics are negligible in the radio detection of extensive air showers.

The area illuminated by radio signals has a limited extent due to the forward-beamed nature of the radio emission. The local energy fluence at the radio detectors with an identified signal is fitted with a two-dimensional distribution function of the signal [28], adapted to the observation altitude of AERA, which takes into account azimuthal asymmetries arising from the superposition of geomagnetic and charge-excess [16–18] effects as well as ring-shaped areas of enhanced emission caused by Cherenkov-like time compression due to the refractive index in the atmosphere [29,30]. During the fit procedure, spurious radio pulses not related to the extensive air shower are flagged and rejected by means of the signal polarization. In rare cases, the flagging of spurious radio pulses can lead to the rejection of a complete event. An example for the resulting fit is illustrated in Fig. 1. For radio events detected in three or four radio detectors, the impact point of the shower axis used for the fit is fixed to the one reconstructed with the Auger surface detector. For radio events with signals in five or more radio detectors, the impact point is determined during the fit of the two-dimensional signal distribution function.

After a successful fit of the signal distribution function, we analytically integrate it over the plane perpendicular to the shower axis. The result is the total energy measured in the radio signal $E_{30-80 \text{ MHz}}^{\text{Auger}}$ (in units of eV), the *radiation energy*. This quantity does not depend on any characteristics of the detector except the finite measurement bandwidth from 30 to 80 MHz. The superscript “Auger” emphasizes that this quantity applies to the geomagnetic-field strength as present at the site of the Pierre Auger Observatory in southern Argentina.

Cross-calibration with the Auger energy scale.—To establish the relation between the radiation energy and the absolute energy scale of cosmic rays, we analyzed data from the first stage of AERA taken between April 2011 and March 2013, when the array consisted of 24 radio detectors equipped with logarithmic-periodic dipole antennas [27]. The signal distribution fit was applied to data preselected with standard Auger quality cuts for surface detector events measured with the 750 m grid of the array. We allowed a maximum zenith angle of 55° and required an energy of at least 10¹⁷ eV. This resulted in a data set with 126 events.

For each of these events, the cosmic-ray energy E_{CR} as reconstructed with the Auger surface detector [31] is available. We stress that the energy reconstruction of the surface detector has been calibrated with the calorimetric energy measurement of the fluorescence detector using a subset of events measured with both detectors simultaneously. Because of the dominance of geomagnetic radio emission [11,18,32] and the scaling of its amplitude with the magnitude of the Lorentz force, the radiation energy scales with $\sin^2(\alpha)$, where α denotes the angle between the air-shower axis and the geomagnetic-field axis. We thus normalize the radiation energy for perpendicular incidence




FIG. 1. Top: Energy fluence for an extensive air shower with an energy of 4.4×10^{17} eV and a zenith angle of 25° as measured in individual AERA radio detectors (circles filled with color corresponding to the measured value) and fitted with the azimuthally asymmetric, two-dimensional signal distribution function (background color). Both radio detectors with a detected signal (*data*) and those below the detection threshold (*subthreshold*) participate in the fit. The fit is performed in the plane perpendicular to the shower axis, with the x axis oriented along the direction of the Lorentz force for charged particles propagating along the shower axis \vec{v} in the geomagnetic field \vec{B} . The best-fitting impact point of the air shower is at the origin of the plot, slightly offset from the one reconstructed with the Auger surface detector [*core (SD)*]. Bottom: Representation of the same data and fitted two-dimensional signal distribution as a function of distance from the shower axis. The colored and black squares denote the energy fluence measurements, and gray squares represent radio detectors with signal below threshold. For the three data points with the highest energy fluence, the one-dimensional projection of the two-dimensional signal distribution fit onto lines connecting the best-fitting impact point of the air shower with the corresponding radio detector positions is illustrated with colored lines. This demonstrates the azimuthal asymmetry and complexity of the two-dimensional signal distribution function. The inset illustrates the polar angles of the three projections. The distribution of the residuals (data versus fit) is shown as well.




FIG. 2. Correlation between the normalized radiation energy and the cosmic-ray energy E_{CR} as determined by the Auger surface detector. Open circles represent air showers with radio signals detected in three or four radio detectors. Solid circles denote showers with five or more detected radio signals.

with respect to the geomagnetic field by dividing it by $\sin^2(\alpha)$. This normalization is valid for all incoming directions of cosmic rays except for a small region around the geomagnetic-field axis. In particular, it is valid for all events in the data set presented here.

In Fig. 2, the value of $E_{30-80 \text{ MHz}}^{\text{Auger}} / \sin^2(\alpha)$ for each measured air shower is plotted as a function of the cosmic-ray energy measured with the Auger surface detector. A log-likelihood fit taking into account threshold effects, measurement uncertainties, and the steeply falling cosmic-ray energy spectrum [33] shows that the data can be described well with the power law

$$E_{30-80 \text{ MHz}}^{\text{Auger}} / \sin^2(\alpha) = A \times 10^7 \text{ eV} (E_{\text{CR}} / 10^{18} \text{ eV})^B. \quad (1)$$

The result of the fit yields $A = 1.58 \pm 0.07$ and $B = 1.98 \pm 0.04$. For a cosmic ray with an energy of 1 EeV arriving perpendicularly to Earth's magnetic field at the Pierre Auger Observatory, the radiation energy thus amounts to 15.8 MeV, a minute fraction of the energy of the primary particle. The observed quadratic scaling is expected for coherent radio emission, for which amplitudes scale linearly and thus the radiated energy scales quadratically.

Taking into account the energy- and zenith-dependent uncertainty of E_{CR} , the resolution of $E_{30-80 \text{ MHz}}^{\text{Auger}} / \sin^2(\alpha)$ is determined from the scatter of points in Fig. 2. It amounts to 22% for the full data set. Performing this analysis for the high-quality subset of events with a successful radio

detection in at least five radio detectors yields a resolution of 17%.

The value of A reported here applies for a cosmic-ray shower with an energy of 1 EeV evolving in a geomagnetic field with a strength of 0.24 G, as present at the site of the Pierre Auger Observatory. With dedicated simulations we confirmed that the radiation energy is only marginally influenced by the charge-excess contribution (at the level of 2% for showers arriving perpendicular to the magnetic field at the Pierre Auger site and less for stronger geomagnetic fields). Hence, a normalization with the field strength of the geomagnetic field is possible and yields

$$E_{30-80 \text{ MHz}} = [15.8 \pm 0.7(\text{stat}) \pm 6.7(\text{syst})] \text{ MeV} \\ \times \left(\sin \alpha \frac{E_{\text{CR}}}{10^{18} \text{ eV}} \frac{B_{\text{Earth}}}{0.24 \text{ G}} \right)^2. \quad (2)$$

$E_{30-80 \text{ MHz}}$ can be used by radio detectors worldwide for cross-calibration of the energy scale, except for experiments deployed at a high altitude where part of the radio emission is clipped when the shower reaches the ground before radiating the bulk of its radio emission. The frequency window from ~ 30 to ~ 80 MHz is shared by many radio detectors [11,34–36]: Below 30 MHz, atmospheric noise and transmitters in the short-wave band dominate; above 80 MHz, coherence diminishes and the FM band interferes with the measurement. Possible second-order effects arising in the determination of $E_{30-80 \text{ MHz}}$, e.g., due to shower geometry, should be addressed in a follow-up analysis, because they could lead to further improvements. The systematic uncertainty of $E_{30-80 \text{ MHz}}$ quoted here arises from the quadratic sum of the systematic uncertainty on the energy scale of the Pierre Auger Observatory (16% at $10^{17.5}$ eV, propagated from the fluorescence detector to the surface detector) and the uncertainty on the radio-electric field amplitude measurement (14%). These two contributions amount to uncertainties of 5.1 and 4.4 MeV in the measurement of the radiation energy at 1 EeV, respectively. We note that the systematic uncertainty in the determination of the cosmic-ray energy from radio measurements is half of that of $E_{30-80 \text{ MHz}}$, as the cosmic-ray energy scales with the square root of the radiation energy.

Comparison with first-principles calculations.—In addition to a *cross-calibration* of techniques and experiments against each other, the radiation energy can also be used for an *independent determination* of the absolute energy scale of cosmic-ray observatories. Sophisticated Monte Carlo simulations [30,37,38] provide a quantitative prediction of the radiation energy based on first-principles calculations combining classical electrodynamics with the well-established properties of the electromagnetic cascade in extensive air showers. A direct comparison of the predicted and measured radiation energies can thus be used for an absolute determination of the energy scale of cosmic-ray detectors.

We have evaluated the radiation energy at a cosmic-ray energy of 1 EeV using the typical zenith angle of our event sample of 37° and a geomagnetic field strength of 0.24 G with the two available full Monte Carlo simulation codes CoREAS [37] and ZHAireS [30]. The predicted values for the radiation energy amount to 11.9 and 11.3 MeV, respectively. Both predictions are thus in agreement with our measurement within the quoted uncertainties. Further work will be undertaken to better understand and minimize experimental and theoretical systematic uncertainties.

Conclusions.—We have measured the radiation energy of extensive air showers and have used it as an energy estimator directly reflecting the calorimetric energy in the electromagnetic cascade. Its value is $15.8 \pm 0.7(\text{stat}) \pm 6.7(\text{syst})$ MeV in the frequency band from 30 to 80 MHz for a cosmic ray with an energy of 10^{18} eV arriving perpendicularly to a magnetic field with a strength of 0.24 G. The radiation energy can be measured at any location that does not suffer from strong anthropogenic noise using moderately sized radio detector arrays. It can thus be used for an efficient cross-calibration of the energy scales of different experiments and detection techniques against each other, in particular, against the well-established energy scale of the Pierre Auger Observatory. Our measurement is in agreement with predictions from first-principles calculations.

The successful installation, commissioning, and operation of the Pierre Auger Observatory would not have been possible without the strong commitment and effort from the technical and administrative staff in Malargüe. We are very grateful to the following agencies and organizations for financial support: Comisión Nacional de Energía Atómica, Agencia Nacional de Promoción Científica y Tecnológica (ANPCyT), Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Gobierno de la Provincia de Mendoza, Municipalidad de Malargüe, NDM Holdings and Valle Las Leñas, in gratitude for their continuing cooperation over land access, Argentina; the Australian Research Council (DP150101622); Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), Financiadora de Estudos e Projetos (FINEP), Fundação de Amparo à Pesquisa do Estado de Rio de Janeiro (FAPERJ), São Paulo Research Foundation (FAPESP) Grants No. 2010/07359-6 and No. 1999/05404-3, Ministério de Ciência e Tecnologia (MCT), Brazil; Grants No. MSMT-CR LG13007, No. 7AMB14AR005, and the Czech Science Foundation Grant No. 14-17501S, Czech Republic; Centre de Calcul IN2P3/CNRS, Centre National de la Recherche Scientifique (CNRS), Conseil Régional Ile-de-France, Département Physique Nucléaire et Corpusculaire (PNC-IN2P3/CNRS), Département Sciences de l'Univers (SDU-INSU/CNRS), Institut Lagrange de Paris (ILP) Grant No. LABEX ANR-10-LABX-63, within the Investissements d'Avenir Programme Grant No. ANR-11-IDEX-0004-02, France; Bundesministerium für Bildung und Forschung

(BMBF), Deutsche Forschungsgemeinschaft (DFG), Finanzministerium Baden-Württemberg, Helmholtz Alliance for Astroparticle Physics (HAP), Helmholtz-Gemeinschaft Deutscher Forschungszentren (HGF), Ministerium für Wissenschaft und Forschung, Nordrhein Westfalen, Ministerium für Wissenschaft, Forschung und Kunst, Baden-Württemberg, Germany; Istituto Nazionale di Fisica Nucleare (INFN), Istituto Nazionale di Astrofisica (INAF), Ministero dell'Istruzione, dell'Università e della Ricerca (MIUR), Gran Sasso Center for Astroparticle Physics (CFA), CETEMPS Center of Excellence, Ministero degli Affari Esteri (MAE), Italy; Consejo Nacional de Ciencia y Tecnología (CONACYT), Mexico; Ministerie van Onderwijs, Cultuur en Wetenschap, Nederlandse Organisatie voor Wetenschappelijk Onderzoek (NWO), Stichting voor Fundamenteel Onderzoek der Materie (FOM), Netherlands; National Centre for Research and Development, Grants No. ERA-NET-ASPERA/01/11 and No. ERA-NET-ASPERA/02/11, National Science Centre, Grants No. 2013/08/M/ST9/00322, No. 2013/08/M/ST9/00728 and No. HARMONIA 5 - 2013/10/M/ST9/00062, Poland; Portuguese national funds and FEDER funds within Programa Operacional Factores de Competitividade through Fundação para a Ciência e a Tecnologia (COMPETE), Portugal; Romanian Authority for Scientific Research ANCS, CNDI-UEFISCDI partnership projects Grants No. 20/2012 and No. 194/2012, Grants No. 1/ASPERA2/2012 ERA-NET, No. PN-II-RU-PD-2011-3-0145-17, and No. PN-II-RU-PD-2011-3-0062, the Minister of National Education, Programme Space Technology and Advanced Research (STAR), Grant No. 83/2013, Romania; Slovenian Research Agency, Slovenia; Comunidad de Madrid, FEDER funds, Ministerio de Educación y Ciencia, Xunta de Galicia, European Community 7th Framework Program, Grant No. FP7-PEOPLE-2012-IEF-328826, Spain; Science and Technology Facilities Council, United Kingdom; Department of Energy, Contracts No. DE-AC02-07CH11359, No. DE-FR02-04ER41300, No. DE-FG02-99ER41107, and No. DE-SC0011689, National Science Foundation, Grant No. 0450696, The Grainger Foundation, USA; NAFOSTED, Vietnam; Marie Curie-IRSES/EPLANET, European Particle Physics Latin American Network, European Union 7th Framework Program, Grant No. PIRSES-2009-GA-246806; and UNESCO.

* auger_spokespersons@fnal.gov

- [1] R. Engel, D. Heck, and T. Pierog, *Annu. Rev. Nucl. Part. Sci.* **61**, 467 (2011).
- [2] V. Verzi for the Pierre Auger Collaboration, in *Proceedings of the 33rd International Cosmic Ray Conference, Rio de Janeiro, Brazil, 2013*, p. 0928, <http://www.cbpf.br/~icrc2013/papers/icrc2013-0928.pdf>.
- [3] P. Abreu *et al.* (Pierre Auger Collaboration), *J. Instrum.* **7**, P09001 (2012).
- [4] J. Abraham *et al.* (Pierre Auger Collaboration), *Astropart. Phys.* **33**, 108 (2010).
- [5] J. Abraham *et al.* (Pierre Auger Collaboration), *Astropart. Phys.* **32**, 89 (2009).
- [6] P. Abreu *et al.* (Pierre Auger Collaboration), *J. Instrum.* **8**, P04009 (2013).
- [7] H. R. Allan, in *Progress in Particle and Nuclear Physics: Cosmic Ray Physics* (North-Holland Publishing Company, Amsterdam, 1971), Vol. 10, p. 169.
- [8] T. Huege, *Phys. Rep.* **620**, 1 (2016).
- [9] F. D. Kahn and I. Lerche, *Proc. R. Soc. A* **289**, 206 (1966).
- [10] K. Werner and O. Scholten, *Astropart. Phys.* **29**, 393 (2008).
- [11] H. Falcke *et al.* (LOPES Collaboration), *Nature* **435**, 313 (2005).
- [12] C. Glaser for the Pierre Auger Collaboration, *AIP Conf. Proc.* **1535**, 68 (2013).
- [13] W. Apel *et al.* (LOPES Collaboration), *Phys. Rev. D* **90**, 062001 (2014).
- [14] P. A. Bezyazeev *et al.* (Tunka-Rex Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **802**, 89 (2015).
- [15] T. Huege, R. Ulrich, and R. Engel, *Astropart. Phys.* **30**, 96 (2008).
- [16] G. A. Askaryan, *Sov. Phys. JETP* **14**, 441 (1962).
- [17] V. Marin for the Codalema Collaboration, in *Proceedings of the 32nd International Cosmic Ray Conference, Beijing, China* (Curran Associates, Inc., Red Hook, New York, 2014), Vol. 1, p. 291.
- [18] A. Aab *et al.* (Pierre Auger Collaboration), *Phys. Rev. D* **89**, 052002 (2014).
- [19] M. P. van Haarlem *et al.*, *Astron. Astrophys.* **556**, A2 (2013).
- [20] A. Aab *et al.* (Pierre Auger Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **798**, 172 (2015).
- [21] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevLett.116.241101> for a more comprehensive presentation.
- [22] K. Weidenhaupt, Ph.D. thesis, RWTH Aachen University, 2014.
- [23] J. Schulz for the Pierre Auger Collaboration, in *Proceedings of Science, the 34th International Cosmic Ray Conference, The Hague, Netherlands, 2015*, 615, <http://pos.sissa.it/cgi-bin/reader/conf.cgi?confid=236>.
- [24] J. Abraham *et al.* (Pierre Auger Collaboration), *Phys. Lett. B* **685**, 239 (2010).
- [25] A. Aab *et al.* (Pierre Auger Collaboration), *Phys. Rev. D* **93**, 122005 (2016).
- [26] P. Abreu *et al.* (Pierre Auger Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **635**, 92 (2011).
- [27] P. Abreu *et al.* (Pierre Auger Collaboration), *J. Instrum.* **7**, P10011 (2012).
- [28] A. Nelles, S. Buitink, H. Falcke, J. R. Hörandel, T. Huege, and P. Schellart, *Astropart. Phys.* **60**, 13 (2015).
- [29] K. D. de Vries, A. M. van den Berg, O. Scholten, and K. Werner, *Phys. Rev. Lett.* **107**, 061101 (2011).
- [30] J. Alvarez-Muñiz, W. R. Carvalho, and E. Zas, *Astropart. Phys.* **35**, 325 (2012).
- [31] I. Mariş for the Pierre Auger Collaboration, in *Proceedings of the 32nd International Cosmic Ray Conference, Beijing,*

- China* (Curran Associates, Inc., Red Hook, New York, 2014), Vol. 1, p. 267.
- [32] D. Ardouin *et al.* (Codalema Collaboration), *Astropart. Phys.* **31**, 192 (2009).
- [33] H. P. Dembinski, B. Kégl, I. C. Mariş, M. Roth, and D. Veberič, *Astropart. Phys.* **73**, 44 (2016).
- [34] D. Ardouin *et al.* (Codalema Collaboration), *Astropart. Phys.* **26**, 341 (2006).
- [35] A. Nelles *et al.*, *J. Cosmol. Astropart. Phys.* **05** (2015) 018.
- [36] D. Kostunin *et al.*, *Nucl. Instrum. Methods Phys. Res., Sect. A* **742**, 89 (2014).
- [37] T. Huege, M. Ludwig, and C. W. James, *AIP Conf. Proc.* **1535**, 128 (2013).
- [38] V. Marin and B. Revenu, *Astropart. Phys.* **35**, 733 (2012).