

Competition of β -delayed protons and β -delayed γ rays in ^{56}Zn and the exotic β -delayed γ -proton decay

S.E.A. Orrigo, B. Rubio, Y. Fujita, B. Blank, W. Gelletly, J. Agramunt, A. Algora, P. Ascher, B. Bilgier, L. Caceres, et al.

► To cite this version:

S.E.A. Orrigo, B. Rubio, Y. Fujita, B. Blank, W. Gelletly, et al.. Competition of β -delayed protons and β -delayed γ rays in ^{56}Zn and the exotic β -delayed γ -proton decay. V. Greco, M. La Cognata, S. Pirrone, F. Rizzo, C. Spitaleri. 12th International Conference on Nucleus - Nucleus Collisions, Jun 2015, Catania, Italy. EDP Sciences, EPJ Web of Conferences, 117, pp.06019, 2016, Proceedings of the 12th International Conference on Nucleus-Nucleus Collisions 2015 <www.lns.infn.it/link/NN2015>. <10.1051/epjconf/201611706019>. <in2p3-01316683>

HAL Id: in2p3-01316683

<http://hal.in2p3.fr/in2p3-01316683>

Submitted on 4 Jul 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Competition of β -delayed protons and β -delayed γ rays in ^{56}Zn and the exotic β -delayed γ -proton decay

S. E. A. ORRIGO^{1*}, B. RUBIO¹, Y. FUJITA², B. BLANK³,
W. GELLETLY⁴, J. AGRAMUNT¹, A. ALGORA^{1,5}, P. ASCHER³,
B. BILGIER⁶, L. CÁCERES⁷, R. B. ÇAKIRLI⁶, H. FUJITA⁸,
E. GANIOĞLU⁶, M. GERBAUX³, J. GIOVINAZZO³, S. GRÉVY³,
O. KAMALOU⁷, H. C. KOZER⁶, L. KUCUK⁶, T. KURTUKIAN-NIETO³,
F. MOLINA^{1,9}, L. POPESCU¹⁰, A. M. ROGERS¹¹, G. SUSOY⁶,
C. STODEL⁷, T. SUZUKI⁸, A. TAMII⁸ and J. C. THOMAS⁷

¹Instituto de Física Corpuscular, CSIC-Universidad de Valencia, E-46071
Valencia, Spain

²Department of Physics, Osaka University, Toyonaka, Osaka 560-0043, Japan

³Centre d'Etudes Nucléaires de Bordeaux Gradignan, CNRS/IN2P3 - Université
Bordeaux 1, 33175 Gradignan Cedex, France

⁴Department of Physics, University of Surrey, Guildford GU2 7XH, Surrey, UK

⁵Inst. of Nuclear Research of the Hung. Acad. of Sciences, Debrecen, H-4026,
Hungary

⁶Department of Physics, Istanbul University, Istanbul, 34134, Turkey

⁷Grand Accélérateur National d'Ions Lourds, BP 55027, F-14076 Caen, France

⁸Research Center for Nuclear Physics, Osaka University, Ibaraki, Osaka 567-0047,
Japan

⁹Comisión Chilena de Energía Nuclear, Casilla 188-D, Santiago, Chile

¹⁰SCK.CEN, Boeretang 200, 2400 Mol, Belgium

¹¹Physics Division, Argonne National Laboratory, Argonne, IL 60439, USA

Abstract

Remarkable results have been published recently on the β decay of ^{56}Zn . In particular, the rare and exotic β -delayed γ -proton emission has been detected for the first time in the fp shell. Here we focus the

*e-mail: sonja.orrigo@ific.uv.es

discussion on this exotic decay mode and on the observed competition between β -delayed protons and β -delayed γ rays from the Isobaric Analogue State.

1 Introduction

Decay spectroscopy is a powerful tool for exploring the structure of nuclei at the drip-lines. β -decay studies, in particular, provide direct access to the absolute values of the Fermi and Gamow-Teller transition strengths, $B(F)$ and $B(GT)$, respectively.

The proton-rich ^{56}Zn nucleus was observed for the first time at GANIL in 1999 [1]. ^{56}Zn is a weakly-bound nucleus lying very close to the proton drip-line. It has a quite small proton separation energy, $S_p = 560(140)$ keV [2], and third component of the isospin quantum number $T_z = -2$.

The first study of the β decay of ^{56}Zn was reported in *ref.* [3]. More recently, some interesting results on ^{56}Zn decay have been reported in *ref.* [4]. Among them the discovery of a rare and exotic decay mode, β -delayed γ -proton decay, which has been seen for the first time in the fp shell. The consequences of this rare decay sequence for the determination of the Gamow-Teller (GT) strength have also been analyzed.

2 The experiment

The experimental study of ^{56}Zn decay was performed at GANIL in 2010. The experiment used a primary beam of $^{58}\text{Ni}^{26+}$ to produce ^{56}Zn . The ^{58}Ni beam, of $3.7\text{ e}\mu\text{A}$ and accelerated to 74.5 MeV/nucleon , was fragmented on a natural Ni target, $200\text{ }\mu\text{m}$ thick. The fragments were selected by the LISE3 separator and implanted into a Double-Sided Silicon Strip Detector (DSSSD). The detection set-up comprised the aforementioned DSSSD detector, $300\text{ }\mu\text{m}$ thick, a silicon ΔE detector located 28 cm upstream, and four EXOGAM Ge clovers surrounding the DSSSD.

The EXOGAM clovers were used to detect β -delayed γ rays. The purpose of the DSSSD was the detection of both the implanted fragments and the subsequent charged-particle decays, *i.e.*, β particles and β -delayed protons. An implantation event was defined by simultaneous signals in both the ΔE and DSSSD detectors. A decay event was defined by a signal above threshold ($50\text{-}90\text{ keV}$) in the DSSSD and no coincident signal in the ΔE .

The implanted ions were identified and selected by putting a gate in a two-dimensional identification matrix, obtained by combining the energy

loss signal from the ΔE detector and the Time-of-Flight. The latter was defined as the time difference between the cyclotron radio-frequency and ΔE signal.

3 Results on the β decay of ^{56}Zn

The results on the β decay of ^{56}Zn [4] are summarized in the decay scheme in *fig. 1* and in table 1, and discussed below.

A half-life of $T_{1/2} = 32.9(8)$ ms was obtained for ^{56}Zn , in agreement with *ref.* [3]. To determine $T_{1/2}$, a decay-time spectrum has been constructed from the time correlations between a decay event in a given pixel of the DSSSD (with a total of 256 pixels) and any implantation signal that occurred before and after it in the same pixel, satisfying the identification condition required to select ^{56}Zn .




Figure 1: Scheme of the β decay of ^{56}Zn . The solid lines indicate observed proton or γ transitions, while the dashed lines correspond to transitions observed in the mirror ^{56}Co nucleus.

The analysis of the charged-particle spectrum measured in the DSSSD has provided new spectroscopic information on the energy levels populated in the ^{56}Cu nucleus, the β -daughter of ^{56}Zn . These levels are shown in *fig. 1*. The comparison of this level spectrum with that of the mirror ^{56}Co , obtained by the $^{56}\text{Fe}(^3\text{He},t)$ charge exchange reaction [5], has been very fruitful.

The analysis of the γ spectrum measured in the EXOGAM clovers and γ -proton coincidences have identified three γ rays at 309, 861 and 1835 keV.

Absolute $B(F)$ and $B(GT)$ strengths have been determined (table 1).

Table 1: β feedings, Fermi and Gamow Teller transition strengths to the ^{56}Cu levels populated in the β^+ decay of ^{56}Zn .

$E_X(\text{keV})$	$I_\beta(\%)$	$B(F)$	$B(GT)$
3508(140)*	43(5)	2.7(5)	
3423(140)	21(1)	1.3(5)	≤ 0.32
2661(140)	14(1)		0.34(6)
2537(140)	0		0
1691(140)	22(6)		0.30(9)
1391(140)	0		0

*Main component of the IAS.

3.1 Competition of β -delayed protons and β -delayed γ rays

In the first study of the ^{56}Zn β decay [3], the emission of β -delayed protons was observed but no β -delayed γ rays were seen. This was not a surprise because, in general, in proton-rich nuclei the proton decay is expected to dominate for states well above (>1 MeV) the proton separation energy S_p . The consequence is that normally the β feeding is directly inferred from the measured intensities of the proton peaks. However, cases where there is a competition between β -delayed proton emission and β -delayed γ de-excitation have also been observed, *e.g.*, in *refs.* [3, 6].

In the $T_z = -2 \rightarrow -1$, β^+ decay of ^{56}Zn to ^{56}Cu , the ^{56}Zn ground state decays with a Fermi transition to its Isobaric Analogue State (IAS) in ^{56}Cu . It should be noted that the de-excitation of this $T = 2$, $J^\pi = 0^+$ IAS via proton decay to the ground state of ^{55}Ni ($T = 1/2$, $J^\pi = 7/2^-$) is isospin forbidden. Therefore the proton emission that we observe can only happen through a $T = 1$ isospin impurity present in the IAS. Moreover in general, when the proton emission is isospin forbidden, the competitive emission of de-exciting γ rays from the IAS also becomes possible and can be observed even from IAS lying at an excitation energy well above S_p [3, 6].

The competition between β -delayed protons and γ rays has indeed been observed in ^{56}Zn . The γ decays represent 56(6)% of the total decays from the 3508 keV IAS. Thus one has to take into account the intensities of both the proton and γ peaks to determine the Fermi strength correctly.

We have also found evidence for the fragmentation of $B(F)$ due to a strong isospin mixing with a 0^+ state at 3423 keV [4], which is important in terms of the mass evaluation [7]. The isospin impurity in the ^{56}Cu IAS, $\alpha^2 = 33(10)\%$ (defined as in *ref.* [5]), and the off-diagonal matrix element of the charge-dependent part of the Hamiltonian, $\langle H_c \rangle = 40(23)$ keV, which is responsible for the isospin mixing of the 3508 keV IAS ($T = 2$, $J^\pi = 0^+$) and the 0^+ part of the 3423 keV level ($T = 1$), are similar to the values obtained in the mirror ^{56}Co nucleus [5].

Thus, the proton decay of the IAS proceeds thanks to the $T = 1$ component. However, considering the quite large isospin mixing in ^{56}Cu , the much faster proton decay ($t_{1/2} \sim 10^{-18}$ s) should dominate on the γ de-excitation ($t_{1/2} \sim 10^{-14}$ s in the mirror). This is not the case since we are still observing the γ decay of the IAS in competition with it.

The knowledge on the nuclear structure of the three nuclei involved in the decay, *i.e.*, ^{56}Zn , ^{56}Cu and ^{55}Ni , can provide us with a possible explanation for the hindrance of the proton decay. Shell model calculations are in progress to clarify this point.

3.2 The β -delayed γ -proton decay

Besides the competition between β -delayed proton emission and γ decay, the exotic sequence of β -delayed γ -proton decay has been detected. Indeed ^{56}Zn does β decay to its IAS in ^{56}Cu and from there we observe the emission of two γ rays of 861 and 1835 keV, populating the ^{56}Cu levels at 2661 and 1691 keV, respectively. Due to the low S_p , these levels are still proton-unbound and thereafter they decay by proton emission. Consequently the rare and exotic β -delayed γ -proton decay has been observed. In addition to these two branches, there is a third case. The 1691 keV level emits a γ ray of 309 keV, going to the level at 1391 keV that is again proton-unbound and then it de-excites by proton emission.

The β -delayed γ -proton decay has been observed here for the first time in the fp shell. This rare decay mode was seen only once before, in the sd shell in ^{32}Ar [6], but the consequences for the determination of $B(GT)$ were not addressed in *ref* [6].

The observation of this special decay mode is very important because it does affect the conventional way to determine $B(GT)$ near the proton drip-

line. For a proper determination of $B(\text{GT})$, indeed, it is crucial to correct the intensity of the proton transitions for the amount of indirect feeding coming from the γ de-excitation. This finding indicates that it is important to employ γ detectors in such studies. This decay mode is expected to be significant in heavier proton-rich nuclei with $T_z \leq -3/2$ under study at RIKEN.

Acknowledgements

This work was supported by the Spanish MICINN grants FPA2008-06419-C02-01, FPA2011-24553; Centro de Excelencia Severo Ochoa del IFIC SEV-2014-0398; CPAN Consolider-Ingenio 2010 Programme CSD2007-00042; *Junta para la Ampliación de Estudios* Programme (CSIC JAE-Doc contract) co-financed by FSE; MEXT, Japan 18540270 and 22540310; Japan-Spain coll. program of JSPS and CSIC; Istanbul University Scientific Research Projects, Num. 5808; UK Science and Technology Facilities Council (STFC) Grant No. ST/F012012/1; Region of Aquitaine. R.B.C. acknowledges support by the Alexander von Humboldt foundation and the Max-Planck-Partner Group. We acknowledge the EXOGAM collaboration for the use of their clover detectors.

References

- [1] J. Giovinazzo et al., *Eur. Phys. J. A*, **11** (2001) 247.
- [2] G. Audi et al., *Nucl. Phys. A*, **729** (2003) 1.
- [3] C. Dossat et al., *Nucl. Phys. A*, **792** (2007) 18.
- [4] S.E.A. Orrigo et al., *Phys. Rev. Lett.*, **112** (2014) 222501.
- [5] H. Fujita et al., *Phys. Rev. C*, **88** (2013) 054329.
- [6] M. Bhattacharya et al., *Phys. Rev. C*, **77** (2008) 065503.
- [7] M. MacCormick and G. Audi, *Nucl. Phys. A*, **925** (2014) 61.