



HAL
open science

Current LHC constraints on minimal universal extra dimensions

N. Deutschmann, T. Flacke, Jong Soo Kim

► **To cite this version:**

N. Deutschmann, T. Flacke, Jong Soo Kim. Current LHC constraints on minimal universal extra dimensions. *Physics Letters B*, 2017, 771, pp.515-520. 10.1016/j.physletb.2017.06.004. in2p3-01454835

HAL Id: in2p3-01454835

<https://hal.in2p3.fr/in2p3-01454835>

Submitted on 14 Dec 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Current LHC constraints on minimal universal extra dimensions



Nicolas Deutschmann^{a,b}, Thomas Flacke^{c,*}, Jong Soo Kim^c

^a Univ Lyon, Université Lyon 1, CNRS/IN2P3, IPNL, F-69622, Villeurbanne, France

^b Centre for Cosmology, Particle Physics and Phenomenology (CP3), Université catholique de Louvain, Chemin du Cyclotron 2, B-1348 Louvain-la-Neuve, Belgium

^c Center for Theoretical Physics of the Universe, Institute for Basic Science (IBS), Daejeon, 34051, Republic of Korea

ARTICLE INFO

Article history:

Received 8 February 2017

Received in revised form 2 May 2017

Accepted 2 June 2017

Available online 7 June 2017

Editor: G.F. Giudice

ABSTRACT

In this letter, we present LHC limits on the minimal universal extra dimension (MUED) model from LHC Run 1 data and current limits from searches of the ongoing Run 2. Typical collider signals of the Kaluza–Klein (KK) states mimic generic degenerate supersymmetry (SUSY) missing transverse momentum signatures since the KK particles cascade decay into jets, leptons and the lightest KK particle which is stable due to KK parity and evades detection. We test the parameter space against a large number of supersymmetry based missing energy searches implemented in the public code `CheckMATE`. We demonstrate the complementarity of employing various searches which target a large number of final state signatures, and we derive the most up to date limits on the MUED parameter space from 13 TeV SUSY searches.

© 2017 The Author(s). Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>). Funded by SCOAP³.

1. Universal extra dimensions – introduction and review

Models with universal extra dimensions (UED) [1] represent a simple extension of the Standard Model which include a dark matter candidate and are testable at the LHC.¹ The extra-dimensions are universal in the sense that all Standard Model (SM) fields are promoted to fields which propagate on the full space–time $\mathcal{M} \times X$, where \mathcal{M} is the flat four dimensional (4D) Minkowski space and X is a compact space. As X is compact, the momenta along the extra-dimensions are discretized. In the 4D effective theory, each extra-dimensional field yields a 4D field without extra-dimensional momentum (the zero-mode which is to be identified with the 4D SM field) as well as a Kaluza–Klein (KK) tower of excitations which are heavy partner states with the same quantum numbers as the zero-mode. The KK mass spectrum is determined by the inverse size and the geometry of the extra-dimensions. In the simplest case of only one extra dimension, – which we focus on in this letter – compactification on the orbifold $X = S^1/Z_2$ allows to have chiral zero-modes of fermions and A_μ zero-mode for gauge fields without an additional A_5 mode.²

The 5D UED model appears to be a very simple and predictive model as it seems to have only one parameter beyond the Standard Model (BSM), the compactification radius R . However, as a 5D theory, the model is inherently non-renormalizable and can only be considered as an effective theory, valid below a cutoff scale Λ , which introduces an additional parameter into the model. Naive dimensional analysis [2–5] and bounds from unitarity violation in gluon KK mode scattering [6] suggest that the cutoff is rather low: $\Lambda R \lesssim \mathcal{O}(10 - 50)$. As a consequence, higher-dimensional operators at the cutoff scale can be phenomenologically relevant.³ In the 5D minimal UED (MUED) model [7], all higher-dimensional operators are assumed to be absent at the cutoff scale Λ , and they are only induced at lower energies due to renormalization group running, thus keeping the model a simple BSM scenario with only two parameters: the inverse compactification radius R^{-1} which sets the mass scale of the first KK excitations, *i.e.* of the lightest partners of the SM fields, and ΛR , which controls the number of KK modes present in the spectrum below the cutoff, and determines how much the KK mode masses and couplings are effected by one-loop running.

* Corresponding author.

E-mail addresses: n.deutschmann@ipnl.in2p3.fr (N. Deutschmann), flacke@ibs.re.kr (T. Flacke), jongsoo.kim@tu-dortmund.de (J.S. Kim).

¹ Cf. Refs. [60,61] for earlier proposals of TeV scale extra dimensions. For Reviews on UED models and their phenomenology cf. [62,63].

² For more than one extra-dimension, the compact space is not unique. Cf. Ref. [64] for a classification of flat 2D orbifolds, Ref. [65,66] for realizations. Models with spherical orbifolds have also been studied [67,68].

³ The least irrelevant operators are boundary localized kinetic terms and other SM-like operators. They are induced by renormalization group running and thus generically present. Their inclusion yields non-minimal UED models [69] with a much larger parameter space. Another UED extension – split UED [70] – includes fermion bulk mass terms. Fermion bulk mass terms are not radiatively induced and could thus be absent consistently.

The phenomenology of the MUED model resembles the phenomenology of the minimal supersymmetric standard model (MSSM) in many ways. Each SM particle is accompanied by a partner particle at the first KK mode level (but in the case of UED, the partners have the same spin as the SM particle). Also, the MUED model possesses a geometric parity (“KK parity”), which is respected by loop corrections and corresponds to the reflection of the orbifold S^1/Z_2 at its midpoint.⁴ The lightest partner state (which in MUED is the partner of the $U(1)_Y$ gauge boson [7]) represents a dark matter candidate [8–11] which reproduces the observed dark matter relic density if $1.25 \text{ TeV} \lesssim R^{-1} \lesssim 1.5 \text{ TeV}$ [11], while for larger R^{-1} , the Universe would be over-closed. Via loop corrections, the KK resonances contribute to electroweak precision observables [1,12,13] and flavor physics [14–16] which impose a bound of $R^{-1} \gtrsim 750 \text{ GeV}$ and $R^{-1} > 600 \text{ GeV}$.⁵

At the LHC, two main signal classes (non-SUSY-like and SUSY-like) allow to test UED models. MUED predicts the existence of a whole tower of partner states of which the first KK level states are KK parity odd, while the second KK level states are the lightest KK parity even BSM states. As loop-induced couplings violate KK number (whilst conserving KK parity), second KK mode states can be resonantly produced at the LHC and searched for in Z' , W' and colored resonance searches [17–19]. The currently strongest known bound on MUED from this signal class amounts to a bound on the second KK photon mass of $m_{A(2)} \gtrsim 1.4 \text{ TeV}$, which corresponds to $R^{-1} \gtrsim 715 \text{ GeV}$ [20] and was obtained from a recast of the CMS search for a di-lepton resonance at 8 TeV with 20.6 fb^{-1} [21].

Complementarily, SUSY searches also provide a high discovery- and exclusion potential for the MUED model. The first KK mode partners are produced in pairs and then cascade decay to the lightest state at the first KK level which itself only leaves a missing energy/momentum signature [1,22]. ATLAS and CMS performed many SUSY searches at 8 TeV and 13 TeV which are suitable to constrain the MUED parameter space. The ATLAS searches Refs. [23,24] provide explicit MUED limits, and [23] reports the currently strongest bound of $R^{-1} \gtrsim 900 - 950 \text{ GeV}$ (depending on the value of ΔR).

In this letter, we provide the MUED bounds obtained from recasts of a large number of SUSY searches at 8 and 13 TeV, performed with CheckMATE.

2. Constraining MUED with existing SUSY searches

We focus on the LHC phenomenology of the SM and the first KK level excitations. The KK zero mode level contains precisely the SM field content, including one Higgs doublet. At the first KK level, each chiral SM fermion (Q_i, u_i, d_i, L_i, e_i) (i is the SM family index) has one a Dirac-fermion partner ($Q_i^{(1)}, u_i^{(1)}, d_i^{(1)}, L_i^{(1)}, e_i^{(1)}$), each SM gauge boson g_μ, W_μ, B_μ has a massive gauge boson partner $g_\mu^{(1)}, W_\mu^{(1)}, B_\mu^{(1)}$, and the Higgs partner sector contains a scalar, a pseudo-scalar and a charged partner ($h^{(1)}, A_0^{(1)}, H_\pm^{(1)}$).⁶ The masses of the SM partners are at tree-level given by $m_n = \sqrt{(n/R)^2 + m_{SM}^2}$, suggesting a very compressed mass spectrum. However, at loop level, the near-mass-degeneracy is partially lifted (with increas-




Fig. 1. Loop corrected KK mass spectrum in MUED for $R^{-1}=960 \text{ GeV}$, $\Delta R=30$ and $m_h=125 \text{ GeV}$.

ing splittings for increased ΔR), making the KK gluon the heaviest state and the KK partner of the $U(1)_Y$ gauge boson the lightest state at each KK level [7,22]. A sample spectrum of the first KK resonances is shown in Fig. 1.⁷ As can be seen, the mass differences can reach up to several hundreds of GeV.

In this letter, we restrict ourselves to the strong production of colored KK modes such as KK gluons and KK quarks,

$$pp \rightarrow g^{(1)} g^{(1)}, \quad pp \rightarrow Q_i^{(1)} Q_j^{(1)}, \quad pp \rightarrow g^{(1)} Q_j^{(1)}, \quad (1)$$

where $Q^{(1)} = Q^{(1)}, q^{(1)}$ and $Q^{(1)}$ denotes the $SU(2)$ doublet quark partners (or their anti-particles) and $q^{(1)} = u^{(1)}, d^{(1)}$ are the $SU(2)$ singlet quark partners (or their anti-particles) taking into account all three generations. The total production cross section is fully determined via QCD interactions and varies between 1(8) pb and 20(480) fb for $R^{-1} = 800 \text{ GeV}$ and $R^{-1} = 1200 \text{ GeV}$ with $\Delta R = 40$ at 8 (13) TeV, respectively.⁸

The decay chain can become very complicated and in the following we briefly discuss typical decay modes of the KK states.⁹ The KK gluon is the heaviest particle, and it decays into KK doublet and singlet quarks with roughly equal branching ratio. The KK quark decay modes mainly depend on its $SU(2)$ charge. The $SU(2)$ singlet KK quark directly decays to the $U(1)_Y$ KK gauge boson γ_1 which is stable. The top KK mode also has sizable decays into the $W_\pm^{(1)}$ and $H_\pm^{(1)}$ as the KK top $SU(2)$ doublet and singlet mix. The $SU(2)$ doublet KK quarks mainly decay into the $SU(2)$ KK gauge bosons $W_\pm^{(1)}$ and $Z^{(1)}$. The KK W and Z bosons are lighter than the KK quarks and thus mostly decay into leptonic KK states which themselves decay further into the lightest stable KK particle $\gamma^{(1)}$. As a result, typical events have a relatively large lepton multiplicity, multiple jets and missing transverse momentum in the final state configuration although all decay products will be relatively soft due to the relatively compressed spectrum.

The masses and decay branching ratios are calculated with the multi purpose Monte Carlo (MC) event generator Herwig++2.7.1 [25] which is also used to generate the fully hadronized MC events employing the default parton distribution function (PDF) set MRST [26]. We work with the default settings

⁴ In UED extensions with boundary terms and bulk fermion masses [71], KK parity is conserved if boundary terms are chosen symmetric on both boundaries and if bulk fermion masses are chosen KK parity odd.

⁵ Bounds from other precision observables, like the muon $g-2$ [72,73], $Zb\bar{b}$ [74], or modifications of Higgs couplings [75–77] are weaker.

⁶ The KK gauge bosons acquire their masses dominantly from the scalar KK modes $g_5^{(1)}, W_5^{(1)}, B_5^{(1)}$, which weakly mix with the first KK Higgs modes due to electroweak symmetry breaking. Thus at each KK level, five scalar degrees of freedom are “eaten” while four physical scalar degrees remain in the electroweak sector.

⁷ Cf. Refs. [7,22] for full mass expressions.

⁸ We clearly see that the cross section increase from 8 to 13 TeV is quite significant for heavy KK masses and thus we expect a large increase in sensitivity from 8 to 13 TeV.

⁹ Cf. Refs. [7,22] for expressions of the relevant zero- and first KK mode couplings at one-loop level, and an illustration of possible decay modes.

Table 1

8 TeV analyses used in our study. Articles are shown by their arXiv number. The middle column denotes the target final state, and the third column shows the total integrated luminosity.

Reference	Final state	\mathcal{L} [fb $^{-1}$]
1403.4853 (ATLAS) [36]	$2\ell+\cancel{E}_T$	20.3
1404.2500 (ATLAS) [37]	SS 2ℓ or 3ℓ	20.3
1405.7875 (ATLAS) [38]	jets + \cancel{E}_T	20.3
1407.0583 (ATLAS) [39]	$1\ell+(b)$ jets+ \cancel{E}_T	20.0
1407.0608 (ATLAS) [40]	monojet+ \cancel{E}_T	20.3
1402.7029 (ATLAS) [41]	$3\ell+\cancel{E}_T$	20.3
1501.03555 (ATLAS) [23]	$1\ell+\text{jets}+\cancel{E}_T$	20.3
1303.2985 (CMS) [42]	α_T+b jets	11.7
1405.7570 (CMS) [43]	$1,SS\text{-}OS2,3,4\ell+\cancel{E}_T$	20.3

Table 2

13 TeV analyses used in our study. The *CONF* papers are only published as conference proceedings. The middle column denotes the target final state, and the third column shows the total integrated luminosity.

Reference	Final state	\mathcal{L} [fb $^{-1}$]
1605.03814 (ATLAS) [44]	jets+ \cancel{E}_T	3.2
1605.04285 (ATLAS) [45]	$1\ell+\text{jets}+\cancel{E}_T$	3.2
ATLAS-CONF-2016-054 [46]	$1\ell+\text{jets}+\cancel{E}_T$	14.8
ATLAS-CONF-2016-076 [47]	2ℓ (stop search)	13
ATLAS-CONF-2016-078 [48]	jets+ \cancel{E}_T	13.3

in all collider tools. We feed the truth level events to CheckMATE2.0.2 [27–29] which is based on the fast detector simulation Delphes 3.4.0 [30]. Fastjet3.2.1 is used for the jet clustering [31]. CheckMATE allows for easy testing of whether a model point is excluded or not at 95% confidence level against current ATLAS and CMS searches at the LHC. CheckMATE requires an event file and the corresponding production cross section as input. For the study presented here, we take the leading order cross section from Herwig++, which yields a very conservative estimate of the constraints as we do not re-scale the leading order estimate with a \mathcal{K} factor. In principle, MC events should be generated with at least one additional parton at matrix element level and matched with the Herwig++ parton shower since the MUED particle spectrum is relatively degenerate [32–34]. However, this is beyond the scope of the current work. Our simple approach will certainly introduce a non-negligible systematic uncertainty in our predictions for mass splitting smaller than 100 GeV, which in particular occurs in the low $\Delta R \leq 5$ region.

We have performed a grid scan in the $R^{-1}-\Delta R$ plane with the SM Higgs mass fixed to $m_h = 125$ GeV. For each grid point, 10^5 events have been generated. We test each grid point against all ATLAS and CMS searches implemented into CheckMATE. However, not every search is sensitive to MUED and we only show the relevant 8 and 13 TeV studies in Table 1 and 2. Most searches are validated and the validation notes can be found on the official webpage.^{10,11} Each analysis typically contains a large number of signal regions which target different mass hierarchies and final state multiplicities. For a given search, the *best signal region* for each point in the parameter space is defined by CheckMATE as that with the largest *expected* exclusion potential. This criterium is then repeated to select the *best search* which is defined as the search whose *best signal region* has the largest expected limit. This




Fig. 2. 95% confidence level limit on the $R^{-1}-\Delta R$ plane from ATLAS and CMS searches performed at the center of mass energy $\sqrt{s} = 8$ TeV. The solid and dashed line denotes the observed and expected 95% confidence level limit, respectively from the searches employed in our paper. The total integrated luminosity used by the various searches are summarized in Table 1. The black solid line denotes the best limit excluding Ref. [23] and the shaded area corresponds to our theoretical uncertainty. We fixed the SM Higgs mass to $m_h = 125$ GeV.

means that the best observed limit is not always used but it ensures that the result is less sensitive to downward fluctuations in the data that are bound to be present when scanning over many searches and many signal regions. Once a *best search* is selected for a given point in parameter space, the signal yield in our model is then compared to the observed limit at 95% confidence level [35],

$$r = \frac{S - 1.96 \cdot \Delta S}{S_{\text{exp}}^{95}} \quad (2)$$

where S , ΔS , and S_{exp}^{95} denotes the number of signal events, its theoretical uncertainty and the experimentally determined 95% confidence level limit on the number of signal events S , respectively. We only consider the statistical uncertainty due to the finite Monte Carlo sample with $\Delta S = \sqrt{S}$. The quantity $S - 1.96 \cdot \Delta S$ corresponds to the 95% lower bound on our prediction for the number of signal event, which ensures that the limits we set are conservative [29]. The r value is only calculated for the expected best signal region. CheckMATE does not combine signal regions nor analyses in order to optimize exclusion since this would require knowledge from the experimental collaborations. We consider a model point as excluded if r is larger than one. However, due to theoretical uncertainties we cannot evaluate directly such as missing next to leading order calculations, PDF uncertainties, details of the parton shower and the finite MC event sample, we assign a conservative uncertainty to our signal prediction. As a result we define a point as definitively allowed for $r < 2/3$ and definitively excluded for $r > 3/2$, which should be large enough to account for the aforementioned effects.

3. Results

Our numerical results are shown in Fig. 2 and 3. We first give a brief summary of the 8 TeV results. Then we present the most up to date limits on the two dimensional MUED parameter space from 13 TeV data.

Run 1 was very successful and both LHC experiments, ATLAS and CMS published a large number of searches sensitive to a multitude of final state signatures. We want to investigate the impact of taking into account all relevant 8 TeV searches on MUED since only a few dedicated studies have been considered for MUED so far. In Fig. 2, we fix the SM Higgs mass to 125 GeV and we present the excluded region at 95% confidence level in the $R^{-1}-\Delta R$ plane.

¹⁰ <https://checkmate.hepforge.org/>.

¹¹ The implementation of the search [24] has been only partially validated and we did not include it in our search. The search [23] presents limits on the MUED parameter space and we plot their limits in Fig. 2. The conference notes [46,48] did not provide cutflows. However, both searches are based on [44,45] with minor modifications and the two latter searches are validated.




Fig. 3. 95% confidence level limit on the R^{-1} – ΔR plane from ATLAS and CMS searches performed at the center of mass energy $\sqrt{s} = 13$ TeV. The solid and dashed line denotes the observed and expected 95% confidence level limit, respectively from the searches employed in our paper. The total integrated luminosity used by the various searches are summarised in Table 2. The black solid line denotes the best limit and the shaded area corresponds to our theoretical uncertainty. We fixed the SM Higgs mass to $m_h = 125$ GeV.

Ref. [23] has an explicit MUED search included, and we show their results for reference. As can be seen, the search Ref. [23] provides the best sensitivity to the MUED parameter space. In particular its dilepton signal regions targeting soft as well as hard leptons provide strong constraints. It should be pointed out that the observed limit of [23] is actually much better than the expected one due to mild down fluctuations in the observed data. As a consequence, a compactification scale up to 950 GeV for $\Delta R = 30$ is excluded. Now, we want to investigate how the other SUSY searches perform in comparison and we explore the sensitivity on MUED excluding Ref. [23]. The limits of other SUSY searches, as obtained with CheckMATE are shown in green, turquoise and brown. We only show results of searches out of the list provided in Table 1 which yield the best expected bound (shown as dashed lines) on R^{-1} for some value of ΔR . The recast observed bounds of the individual searches are shown as solid lines. The black solid curve denotes the 95% confidence level bound as determined from Eq. (2) for $r = 1$,¹² and the grey shaded area is our estimate for the uncertainty and denotes the region with $0.67 < r < 1.5$. Considering searches other than Ref. [23], the limit is mainly driven by [38,41,43]. [38] is the ‘vanilla’ ATLAS multijet and large missing transverse momentum search which performs well for sizable ΔR . For small $\Delta R \sim 5$ GeV, the mass difference between the KK gluon g_1 and the lightest KK mode γ_1 diminishes quite rapidly with a splitting of ~ 100 GeV. In this regime, monojet searches can become competitive. However, since we did not generate matched event samples, the systematic uncertainty on the exclusion limit in this region of parameter space would be too large and thus the results should be interpreted with great care. Ref. [41] and [43] are dedicated multi-lepton and missing transverse momentum searches. Multiple leptons in the final state are very handy in suppressing the SM background. It is clear that for small ΔR values, the leptons become too soft and thus the multi-lepton searches lose their sensitivity very quickly. Again, dedicated studies targeting soft leptons such as Ref. [23] and pos-

sibly monojet searches will improve sensitivity in this degenerate region.

To summarize, we have demonstrated that a large number of searches can provide complementary constraints on the MUED parameter space. The strongest constraints from 8 TeV searches are obtained from the dedicated MUED motivated signal region of Ref. [23], but signal regions of other SUSY searches yield bounds which are not substantially weaker. Other 8 TeV searches such as [49] could be sensitive to the signals we consider, but are currently not implemented or validated in CheckMATE.

Now, we turn our attention to LHC searches at $\sqrt{s} = 13$ TeV. So far, ATLAS and CMS did not present dedicated MUED results from Run 2. In Fig. 3, we display the excluded region from the ongoing Run 2 as obtained from our recast study with CheckMATE. We include two published searches [44,45] from the early Run 2 data and three conference notes with roughly 14 fb^{-1} of collected data [48,47,46]. Other searches such as [50–58] could also be sensitive to the signals we consider, but are not yet implemented or validated in CheckMATE. We observe a clear improvement of the limits as compared to Run 1. The mass scale up to $R^{-1} = 1500$ GeV can be excluded at 95% confidence level. The search [46] requiring one isolated lepton performs better than the corresponding search with a lepton veto [48]. For both searches, the limits become much weaker for increasing ΔR . This might be quite surprising since naively one would expect that for increasing mass splitting, the efficiency in the signal regions would increase. However, increasing ΔR raises also the overall mass scale of the hard process which results in a large reduction of the hadronic cross section. In addition, the best signal region of Ref. [46] is GG2J which targets relatively heavy and compressed spectra. Sensitivity is enhanced for events with relatively large jet recoils against the pair produced KK partons which results in large missing transverse momentum. However, for increasing mass splitting, the missing transverse energy is reduced and thus we observe that the best limits are derived for relatively small ΔR . On the other hand, the dilepton stop search [47] performs very well even for large ΔR values. Here, a large mass splitting results in more energetic leptons which improves the sensitivity. However, the response degrades very quickly for very small ΔR as in this region, the leptons tend to be too soft to be detected.

4. Conclusions

We have determined the current limits on the MUED parameter space from LHC supersymmetry searches at 8 and 13 TeV. After the end of Run 1, a large number of supersymmetry searches targeting a vast number of final state topologies have been published. We demonstrated the complementarity of using all available searches in order to constrain MUED. Run 2 is ongoing and the number of searches is quite restricted compared to Run 1. However, the large gain in parton luminosity allows for impressive improvement in the exclusion of parameter space. The dilepton stop search performs very well, and we have derived the most stringent limits on the MUED parameter space so far with $R^{-1} \approx 1400$ GeV excluded for $\Delta R \sim 10$. This limit is conservative in the sense that it does not include and \mathcal{K} factor. For a \mathcal{K} factor of 1.5, the limit would be increased to $R^{-1} \approx 1500$ GeV. MUED yields the observed dark matter relic density if co-annihilation and second KK mode resonant (co-) annihilation are taken into account if $1.25 \text{ TeV} \lesssim R^{-1} \lesssim 1.5 \text{ TeV}$ [11] while larger R^{-1} predicts a too large dark matter relic density. Thus the results from ATLAS and CMS Run 2 recasts of SUSY searches presented in this letter show that the MUED model (*i.e.* the simplest UED benchmark model) is getting in tension with observation.

¹² The apparent discrepancy between the limit shown by this line and trying to follow the individual limits is a result of the interpolation we performed to produce this figure. In particular, the displayed expected limit for the CMS search is in reality briefly the best search around $\Delta R = 20$, which is not displayed because the interpolation smoothed the curve. This explains the outward bulge of the limit in that region.

5. Note added

While we were finishing this letter, we became aware of the complementary study Ref. [59], which studies MUED bounds from SUSY searches with an MUED implementation in `Pythia 6` and `Pythia 8` for event generation and `CheckMATE`. We thank the authors of Ref. [59] for communication and for making a preliminary draft of their paper available to us. In the overlapping regions of the two studies, the results were found to be in reasonable agreement.

Acknowledgements

We thank Kyoungchul Kong and Seong Chan Park for their contribution and collaboration in the beginning of this work as well as for many helpful comments. We also thank Simon Platzer and Peter Richardson for information and details on the MUED implementation in `Herwig`. The work of J.S.K and T.F. was supported by IBS under the project code, IBS-R018-D1. N.D. and T.F. acknowledge the Partenariat Hubert Curien (PHC) STAR project no.34299VE, and partial support from the CNRS LIA FKPLP.

References

- [1] T. Appelquist, H.-C. Cheng, B.A. Dobrescu, Bounds on universal extra dimensions, *Phys. Rev. D* 64 (2001) 035002, arXiv:hep-ph/0012100.
- [2] R. Barbieri, A. Strumia, What is the limit on the Higgs mass?, *Phys. Lett. B* 462 (1999) 144–149, arXiv:hep-ph/9905281.
- [3] M. Papucci, NDA and perturbativity in Higgsless models, arXiv:hep-ph/0408058.
- [4] Z. Chacko, M.A. Luty, E. Ponton, Massive higher dimensional gauge fields as messengers of supersymmetry breaking, *J. High Energy Phys.* 07 (2000) 036, arXiv:hep-ph/9909248.
- [5] G. Bhattacharyya, A. Datta, S.K. Majee, A. Raychaudhuri, Power law blitzkrieg in universal extra dimension scenarios, *Nucl. Phys. B* 760 (2007) 117–127, arXiv:hep-ph/0608208.
- [6] R.S. Chivukula, D.A. Dicus, H.-J. He, Unitarity of compactified five-dimensional Yang–Mills theory, *Phys. Lett. B* 525 (2002) 175–182, arXiv:hep-ph/0111016.
- [7] H.-C. Cheng, K.T. Matchev, M. Schmaltz, Radiative corrections to Kaluza–Klein masses, *Phys. Rev. D* 66 (2002) 036005, arXiv:hep-ph/0204342.
- [8] G. Servant, T.M.P. Tait, Is the lightest Kaluza–Klein particle a viable dark matter candidate?, *Nucl. Phys. B* 650 (2003) 391–419, arXiv:hep-ph/0206071.
- [9] K. Kong, K.T. Matchev, Precise calculation of the relic density of Kaluza–Klein dark matter in universal extra dimensions, *J. High Energy Phys.* 01 (2006) 038, arXiv:hep-ph/0509119.
- [10] F. Burnell, G.D. Kribs, The abundance of Kaluza–Klein dark matter with coannihilation, *Phys. Rev. D* 73 (2006) 015001, arXiv:hep-ph/0509118.
- [11] G. Belanger, M. Kakizaki, A. Pukhov, Dark matter in UED: the role of the second KK level, *J. Cosmol. Astropart. Phys.* 1102 (2011) 009, arXiv:1012.2577 [hep-ph].
- [12] T. Appelquist, H.-U. Yee, Universal extra dimensions and the Higgs boson mass, *Phys. Rev. D* 67 (2003) 055002, arXiv:hep-ph/0211023.
- [13] M. Baak, M. Goebel, J. Haller, A. Hoecker, D. Ludwig, K. Moenig, M. Schott, J. Stelzer, Updated status of the global electroweak fit and constraints on new physics, *Eur. Phys. J. C* 72 (2012) 2003, arXiv:1107.0975 [hep-ph].
- [14] A.J. Buras, M. Spranger, A. Weiler, The impact of universal extra dimensions on the unitarity triangle and rare K and B decays, *Nucl. Phys. B* 660 (2003) 225–268, arXiv:hep-ph/0212143.
- [15] A.J. Buras, A. Poschenrieder, M. Spranger, A. Weiler, The impact of universal extra dimensions on $B \rightarrow X_s \gamma$, $B \rightarrow X_s g$, $B \rightarrow X_s \mu^+ \mu^-$, $K_L \rightarrow \pi_0 e^+ e^-$ and $e^+ e^-$, *Nucl. Phys. B* 678 (2004) 455–490, arXiv:hep-ph/0306158.
- [16] U. Haisch, A. Weiler, Bound on minimal universal extra dimensions from $\bar{B} \rightarrow X_s \gamma$, *Phys. Rev. D* 76 (2007) 034014, arXiv:hep-ph/0703064.
- [17] A. Datta, K. Kong, K.T. Matchev, Discrimination of supersymmetry and universal extra dimensions at hadron colliders, *Phys. Rev. D* 72 (2005) 096006, arXiv:hep-ph/0509246, *Phys. Rev. D* 72 (2005) 119901 (Erratum).
- [18] T. Flacke, A. Menon, Z. Sullivan, Constraints on UED from W' searches, *Phys. Rev. D* 86 (2012) 093006, arXiv:1207.4472 [hep-ph].
- [19] S. Chang, S.Y. Shim, J. Song, Direct bound on the minimal universal extra dimension model from the $t\bar{t}$ resonance search at the tevatron, *Phys. Rev. D* 86 (2012) 117503, arXiv:1207.6876 [hep-ph].
- [20] L. Edelhauser, T. Flacke, M. Kramer, Constraints on models with universal extra dimensions from dilepton searches at the LHC, *J. High Energy Phys.* 08 (2013) 091, arXiv:1302.6076 [hep-ph].
- [21] CMS Collaboration, Search for resonances in the dilepton mass distribution in pp collisions at $\sqrt{s} = 8$ TeV, CMS-PAS-EXO-12-061.
- [22] H.-C. Cheng, K.T. Matchev, M. Schmaltz, Bosonic supersymmetry? Getting fooled at the CERN LHC, *Phys. Rev. D* 66 (2002) 056006, arXiv:hep-ph/0205314.
- [23] ATLAS Collaboration, G. Aad, et al., Search for squarks and gluinos in events with isolated leptons, jets and missing transverse momentum at $\sqrt{s} = 8$ TeV with the ATLAS detector, *J. High Energy Phys.* 04 (2015) 116, arXiv:1501.03555 [hep-ex].
- [24] ATLAS Collaboration, Search for squarks and gluinos in events with isolated leptons, jets and missing transverse momentum at $\sqrt{s} = 8$ TeV with the ATLAS detector, Tech. Rep. ATLAS-CONF-2013-062, CERN, Geneva, Jun 2013, <https://cds.cern.ch/record/1557779>.
- [25] J. Bellm, et al., `Herwig++` 2.7 release note, arXiv:1310.6877 [hep-ph].
- [26] A.D. Martin, R.G. Roberts, W.J. Stirling, R.S. Thorne, NNLO global parton analysis, *Phys. Lett. B* 531 (2002) 216–224, arXiv:hep-ph/0201127.
- [27] D. Dercks, N. Desai, J.S. Kim, K. Rolbiecki, J. Tattersall, T. Weber, `CheckMATE 2`: from the model to the limit, arXiv:1611.09856 [hep-ph].
- [28] J.S. Kim, D. Schmeier, J. Tattersall, K. Rolbiecki, A framework to create customised LHC analyses within `CheckMATE`, *Comput. Phys. Commun.* 196 (2015) 535–562, arXiv:1503.01123 [hep-ph].
- [29] M. Drees, H. Dreiner, D. Schmeier, J. Tattersall, J.S. Kim, `CheckMATE`: confronting your favourite new physics model with LHC data, *Comput. Phys. Commun.* 187 (2015) 227–265, arXiv:1312.2591 [hep-ph].
- [30] DELPHES 3 Collaboration, J. de Favereau, C. Delaere, P. Demin, A. Giammanco, V. Lemaitre, A. Mertens, M. Selvaggi, DELPHES 3, a modular framework for fast simulation of a generic collider experiment, *J. High Energy Phys.* 02 (2014) 057, arXiv:1307.6346 [hep-ex].
- [31] M. Cacciari, G.P. Salam, G. Soyez, `FastJet` user manual, *Eur. Phys. J. C* 72 (2012) 1896, arXiv:1111.6097 [hep-ph].
- [32] H.K. Dreiner, M. Kramer, J. Tattersall, How low can SUSY go? Matching, mono-jets and compressed spectra, *Europhys. Lett.* 99 (2012) 61001, arXiv:1207.1613 [hep-ph].
- [33] H. Dreiner, M. Krämer, J. Tattersall, Exploring QCD uncertainties when setting limits on compressed supersymmetric spectra, *Phys. Rev. D* 87 (3) (2013) 035006, arXiv:1211.4981 [hep-ph].
- [34] M. Drees, M. Hanussek, J.S. Kim, Light stop searches at the LHC with monojet events, *Phys. Rev. D* 86 (2012) 035024, arXiv:1201.5714 [hep-ph].
- [35] A.L. Read, Presentation of search results: the CL(s) technique, *J. Phys. G* 28 (2002) 2693–2704.
- [36] ATLAS Collaboration, G. Aad, et al., Search for direct top-squark pair production in final states with two leptons in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector, *J. High Energy Phys.* 06 (2014) 124, arXiv:1403.4853 [hep-ex].
- [37] ATLAS Collaboration, G. Aad, et al., Search for supersymmetry at $\sqrt{s} = 8$ TeV in final states with jets and two same-sign leptons or three leptons with the ATLAS detector, *J. High Energy Phys.* 06 (2014) 035, arXiv:1404.2500 [hep-ex].
- [38] ATLAS Collaboration, G. Aad, et al., Search for squarks and gluinos with the ATLAS detector in final states with jets and missing transverse momentum using $\sqrt{s} = 8$ TeV proton–proton collision data, *J. High Energy Phys.* 09 (2014) 176, arXiv:1405.7875 [hep-ex].
- [39] ATLAS Collaboration, G. Aad, et al., Search for top squark pair production in final states with one isolated lepton, jets, and missing transverse momentum in $\sqrt{s} = 8$ TeV pp collisions with the ATLAS detector, *J. High Energy Phys.* 11 (2014) 118, arXiv:1407.0583 [hep-ex].
- [40] ATLAS Collaboration, G. Aad, et al., Search for pair-produced third-generation squarks decaying via charm quarks or in compressed supersymmetric scenarios in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector, *Phys. Rev. D* 90 (5) (2014) 052008, arXiv:1407.0608 [hep-ex].
- [41] ATLAS Collaboration, G. Aad, et al., Search for direct production of charginos and neutralinos in events with three leptons and missing transverse momentum in $\sqrt{s} = 8$ TeV pp collisions with the ATLAS detector, *J. High Energy Phys.* 04 (2014) 169, arXiv:1402.7029 [hep-ex].
- [42] CMS Collaboration, S. Chatrchyan, et al., Search for supersymmetry in hadronic final states with missing transverse energy using the variables α_T and b-quark multiplicity in pp collisions at $\sqrt{s} = 8$ TeV, *Eur. Phys. J. C* 73 (9) (2013) 2568, arXiv:1303.2985 [hep-ex].
- [43] CMS Collaboration, V. Khachatryan, et al., Searches for electroweak production of charginos, neutralinos, and sleptons decaying to leptons and W , Z , and Higgs bosons in pp collisions at 8 TeV, *Eur. Phys. J. C* 74 (9) (2014) 3036, arXiv:1405.7570 [hep-ex].
- [44] ATLAS Collaboration, M. Aaboud, et al., Search for squarks and gluinos in final states with jets and missing transverse momentum at $\sqrt{s} = 13$ TeV with the ATLAS detector, *Eur. Phys. J. C* 76 (7) (2016) 392, arXiv:1605.03814 [hep-ex].
- [45] ATLAS Collaboration, G. Aad, et al., Search for gluinos in events with an isolated lepton, jets and missing transverse momentum at $\sqrt{s} = 13$ TeV with the ATLAS detector, *Eur. Phys. J. C* 76 (10) (2016) 565, arXiv:1605.04285 [hep-ex].
- [46] ATLAS Collaboration, Search for squarks and gluinos in events with an isolated lepton, jets and missing transverse momentum at $\sqrt{s} = 13$ TeV with the ATLAS detector, Tech. Rep. ATLAS-CONF-2016-054, CERN, Geneva, Aug 2016, <https://cds.cern.ch/record/2206136>.

- [47] ATLAS Collaboration, Search for direct top squark pair production and dark matter production in final states with two leptons in $\sqrt{s} = 13$ TeV pp collisions using 13.3 fb^{-1} of ATLAS data, Tech. Rep. ATLAS-CONF-2016-076, CERN, Geneva, Aug 2016, <http://cds.cern.ch/record/2206249>.
- [48] ATLAS Collaboration, Further searches for squarks and gluinos in final states with jets and missing transverse momentum at $\sqrt{s} = 13$ TeV with the ATLAS detector, Tech. Rep. ATLAS-CONF-2016-078, CERN, Geneva, Aug 2016, <http://cds.cern.ch/record/2206252>.
- [49] ATLAS Collaboration, Searches for direct scalar top pair production in final states with two leptons using the transverse mass variable and a multivariate analysis technique in $\sqrt{s} = 8$ TeV pp collisions using 20.3 fb^{-1} of ATLAS data, Tech. Rep. ATLAS-CONF-2013-065, Geneva, <https://cds.cern.ch/record/1562840>, 2013.
- [50] ATLAS Collaboration, M. Aaboud, et al., Search for new phenomena in events containing a same-flavour opposite-sign dilepton pair, jets, and large missing transverse momentum in $\sqrt{s} = 13$ pp collisions with the ATLAS detector, Eur. Phys. J. C 77 (3) (2017) 144, arXiv:1611.05791 [hep-ex].
- [51] ATLAS Collaboration, Search for supersymmetry with two same-sign leptons or three leptons using 13.2 fb^{-1} of $\sqrt{s} = 13$ TeV pp collision data collected by the ATLAS detector, Tech. Rep. ATLAS-CONF-2016-037, Geneva, <https://cds.cern.ch/record/2205745>, 2016.
- [52] ATLAS Collaboration, Search for squarks and gluinos in final states with jets and missing transverse momentum using 36 fb^{-1} of $\sqrt{s} = 13$ TeV pp collision data with the ATLAS detector, Tech. Rep. ATLAS-CONF-2017-022, Geneva, <https://cds.cern.ch/record/2258145>, 2017.
- [53] CMS Collaboration, V. Khachatryan, et al., Search for supersymmetry with multiple charged leptons in proton–proton collisions at $\sqrt{s} = 13$ TeV, arXiv:1701.06940 [hep-ex].
- [54] CMS Collaboration, A.M. Sirunyan, et al., Searches for pair production for third-generation squarks in $\sqrt{s} = 13$ TeV pp collisions, arXiv:1612.03877 [hep-ex].
- [55] CMS Collaboration, V. Khachatryan, et al., Search for new physics in final states with two opposite-sign, same-flavor leptons, jets, and missing transverse momentum in pp collisions at $\sqrt{s} = 13$ TeV, J. High Energy Phys. 12 (2016) 013, arXiv:1607.00915 [hep-ex].
- [56] CMS Collaboration, V. Khachatryan, et al., Search for supersymmetry in the multijet and missing transverse momentum final state in pp collisions at 13 TeV, Phys. Lett. B 758 (2016) 152–180, arXiv:1602.06581 [hep-ex].
- [57] CMS Collaboration, Search for supersymmetry in pp collisions at $\sqrt{s} = 13$ TeV in the single-lepton final state using the sum of masses of large-radius jets, Tech. Rep. CMS-PAS-SUS-16-037, Geneva, <https://cds.cern.ch/record/2256652>, 2017.
- [58] CMS Collaboration, Search for new physics with multileptons and jets in 35.9 fb^{-1} of pp collision data at $\sqrt{s} = 13$ TeV, Tech. Rep. CMS-PAS-SUS-16-041, Geneva, <https://cds.cern.ch/record/2256435>, 2017.
- [59] J. Beuria, A. Datta, D. Debnath, K.T. Matchev, LHC collider phenomenology of minimal universal extra dimensions, arXiv:1702.00413 [hep-ph].
- [60] I. Antoniadis, A possible new dimension at a few TeV, Phys. Lett. B 246 (1990) 377–384.
- [61] I. Antoniadis, N. Arkani-Hamed, S. Dimopoulos, G.R. Dvali, New dimensions at a millimeter to a Fermi and superstrings at a TeV, Phys. Lett. B 436 (1998) 257–263, arXiv:hep-ph/9804398.
- [62] D. Hooper, S. Profumo, Dark matter and collider phenomenology of universal extra dimensions, Phys. Rep. 453 (2007) 29–115, arXiv:hep-ph/0701197.
- [63] G. Servant, Status report on universal extra dimensions after LHC8, Mod. Phys. Lett. A 30 (15) (2015) 1540011, arXiv:1401.4176 [hep-ph].
- [64] L. Nilse, Classification of 1D and 2D orbifolds, AIP Conf. Proc. 903 (2007) 411–414, arXiv:hep-ph/0601015.
- [65] B.A. Dobrescu, E. Ponton, Chiral compactification on a square, J. High Energy Phys. 03 (2004) 071, arXiv:hep-th/0401032.
- [66] G. Cacciapaglia, A. Deandrea, J. Llodra-Perez, A dark matter candidate from Lorentz invariance in 6D, J. High Energy Phys. 03 (2010) 083, arXiv:0907.4993 [hep-ph].
- [67] G. Cacciapaglia, A. Deandrea, N. Deutschmann, Dark matter and localised fermions from spherical orbifolds?, J. High Energy Phys. 04 (2016) 083, arXiv:1601.00081 [hep-ph].
- [68] N. Maru, T. Nomura, J. Sato, M. Yamanaka, The universal extra dimensional model with S^2/Z_2 extra-space, Nucl. Phys. B 830 (2010) 414–433, arXiv:0904.1909 [hep-ph].
- [69] T. Flacke, A. Menon, D.J. Phalen, Non-minimal universal extra dimensions, Phys. Rev. D 79 (2009) 056009, arXiv:0811.1598 [hep-ph].
- [70] S.C. Park, J. Shu, Split universal extra dimensions and dark matter, Phys. Rev. D 79 (2009) 091702, arXiv:0901.0720 [hep-ph].
- [71] T. Flacke, K. Kong, S.C. Park, Phenomenology of universal extra dimensions with bulk-masses and brane-localized terms, J. High Energy Phys. 05 (2013) 111, arXiv:1303.0872 [hep-ph].
- [72] P. Nath, M. Yamaguchi, Effects of Kaluza–Klein excitations on $(g_\mu - 2)$, Phys. Rev. D 60 (1999) 116006, arXiv:hep-ph/9903298.
- [73] K. Agashe, N.G. Deshpande, G.H. Wu, Can extra dimensions accessible to the SM explain the recent measurement of anomalous magnetic moment of the muon?, Phys. Lett. B 511 (2001) 85–91, arXiv:hep-ph/0103235.
- [74] J.F. Oliver, J. Papavassiliou, A. Santamaria, Universal extra dimensions and $Z \rightarrow b\bar{b}$, Phys. Rev. D 67 (2003) 056002, arXiv:hep-ph/0212391.
- [75] U.K. Dey, T.S. Ray, Constraining minimal and nonminimal universal extra dimension models with Higgs couplings, Phys. Rev. D 88 (5) (2013) 056016, arXiv:1305.1016 [hep-ph].
- [76] T. Kakuda, K. Nishiwaki, K.-y. Oda, R. Watanabe, Universal extra dimensions after Higgs discovery, Phys. Rev. D 88 (2013) 035007, arXiv:1305.1686 [hep-ph].
- [77] T. Flacke, K. Kong, S.C. Park, 126 GeV Higgs in next-to-minimal universal extra dimensions, Phys. Lett. B 728 (2014) 262–267, arXiv:1309.7077 [hep-ph].