



HAL
open science

Virtual compton scattering at low energy and the generalized polarizabilities of the nucleon

H. Fonvieille

► **To cite this version:**

H. Fonvieille. Virtual compton scattering at low energy and the generalized polarizabilities of the nucleon. International Conference on Nuclear and Particle Physics with CEBAF at Jefferson Laboratory 2 NAPP03, May 2003, Dubrovnik, Croatia. pp.209-216. in2p3-00014006

HAL Id: in2p3-00014006

<https://hal.in2p3.fr/in2p3-00014006>

Submitted on 22 Sep 2003

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

VIRTUAL COMPTON SCATTERING
AT LOW ENERGY AND THE
GENERALIZED POLARIZABILITIES OF THE NUCLEON

H. FONVIEILLE

*for the Jefferson Lab Hall A Collaboration
and the VCS Collaboration*

*Laboratoire de Physique Corpusculaire IN2P3/CNRS
Université Blaise Pascal - Clermont II
63170 AUBIERE Cedex, FRANCE*

Abstract

Virtual Compton Scattering (VCS) $\gamma^*p \rightarrow \gamma p$ at low CM energy gives access to the Generalized Polarizabilities of the nucleon. These observables generalize the concept of electromagnetic polarizabilities to the case of a virtual photon. Dedicated VCS experiments have been performed at MAMI, Jefferson Lab and MIT-Bates. The experimental status is reviewed, including analysis methods and physics results. The measurement of absolute ($ep \rightarrow ep\gamma$) cross sections allows the extraction of the two unpolarized VCS structure functions $P_{LL} - P_{TT}/\epsilon$ and P_{LT} , which are combinations of the Generalized Polarizabilities of the proton. Future prospects in the field of VCS at low energy are also presented.

The study of Virtual Compton Scattering (VCS) off the proton: $\gamma^*p \rightarrow \gamma p$ provides new insights into the nucleon structure. It is related to the “two-photon physics” that was mentioned at this Conference when discussing the proton form factor measurements (see the talk of M.Jones). My talk is about the part of VCS that is devoted to the investigation of Generalized Polarizabilities (GPs).

1 Real and Virtual Compton Scattering and the concept of Generalized Polarizabilities

We recall that the electric and magnetic polarizabilities α_E and β_M that are studied in Real Compton Scattering (RCS) [1] are a measure of the deformation of the nucleon structure under an applied EM field. Their smallness indicates that the proton is a very rigid object. The smallness of β_M relative to α_E is usually interpreted in terms of a cancellation between a diamagnetic (pion cloud) and a paramagnetic (quark core) contribution.

The concept of Generalized Polarizabilities was first established by P.Guichon *et al.* in 1995 [2], introducing observables that are defined not only at $Q^2 = 0$ but at any photon virtuality Q^2 . Thus the GPs measure the EM deformation locally inside the nucleon,

with a scale given by Q^2 , and they can be seen as “deformed form factors”. The full formalism [3, 4] leads to six independent GPs at lowest order: the two scalar ones $\alpha_E(Q^2)$ and $\beta_M(Q^2)$ (giving the above mentioned RCS polarizabilities at $Q^2 = 0$) and four spin GPs.

VCS can be accessed via photon electroproduction $ep \rightarrow ep\gamma$. The amplitude of this process is decomposed into the coherent sum of the Bethe-Heitler (BH), the VCS Born, and the VCS non-Born amplitudes. The (BH+Born) cross section is known and depends only on the proton EM form factors. The VCS non-Born term contains the unknown physics, i.e. the GPs which are the goal of the measurements.

2 Analysis methods

Up to now, our knowledge of GPs comes from the measurement of cross section for the (unpolarized) $ep \rightarrow ep\gamma$ process. The main kinematic variables for physics analysis are the four-momentum transfer squared Q^2 , the available energy \sqrt{s} in the (γp) center-of-mass (CM), and the CM variables of the outgoing photon: its momentum q' and its spherical angles θ and φ w.r.t. the virtual photon momentum vector \vec{q} . There are presently two methods to extract GPs, depending on the range of CM energy. A short description of these methods is given below.

2.1 The Low Energy Theorem

Below pion production threshold ($\sqrt{s} \leq (m_p + m_\pi)$) one can use the Low Energy Theorem (LET), first applied to VCS by P.Guichon *et al.* [2]. In this context one obtains the following expression for the ($ep \rightarrow ep\gamma$) cross section:

$$d^5\sigma^{EXP} = d^5\sigma^{BH+Born} + (\phi q') \times [v_1(P_{LL} - \frac{1}{\epsilon}P_{TT}) + v_2P_{LT}] + \mathcal{O}(q'^2) \quad (1)$$

where $d^5\sigma^{BH+Born}$ represents the Bethe-Heitler+Born cross section, $(\phi q')$ a phase space factor, and v_1 and v_2 are known kinematic factors [3]. $P_{LL} - \frac{1}{\epsilon}P_{TT}$ and P_{LT} are the structure functions to be measured, and ϵ is the virtual photon polarization. These structure functions are combinations of the GPs (see [3] for details); they contain the electric and magnetic Generalized Polarizabilities $\alpha_E(Q^2)$ and $\beta_M(Q^2)$ under the form:

$$\begin{aligned} P_{LL} &\sim \alpha_E(Q^2) \\ P_{LT} &\sim \beta_M(Q^2) + \text{spin GPs.} \end{aligned} \quad (2)$$

To extract the structure functions from the data at fixed $|\vec{q}'|$ and fixed ϵ , one basically makes a fit of the quantity $(d^5\sigma^{EXP} - d^5\sigma^{BH+Born})/(\phi q')$ to the expression of the bracketed term in eq. 1, thus adjusting two free parameters. The main difficulty of the method is that the effect of GPs is small below pion threshold; it amounts to no more than 10-15% of the cross section. So systematics of the experiments have to be well under control.

2.2 The approach of Dispersion Relations

Both below and above pion threshold and up to the $\pi\pi N$ threshold, one can use the Dispersion Relation formalism (DR) developed by B.Pasquini *et al.* for Real and Virtual Compton Scattering [5]. This allows VCS measurements in the $\Delta(1232)$ resonance region to be interpreted in terms of GPs. In this model the GPs $\alpha_E(Q^2)$ and $\beta_M(Q^2)$ contain

free parameters; experimentally we fit their value at a given Q^2 . The advantage of this method is that the $(ep\gamma)$ cross section is more sensitive to the GPs in the resonance region than below pion threshold.

3 Experiments and results

Table 1 summarizes the experiments performed as of today in the $(ep \rightarrow ep\gamma)$ channel to investigate the GPs. All of them use magnetic spectrometers to detect the final electron and proton, and they identify the single photon electroproduction reaction by the missing mass technique. These difficult experiments usually have to meet the following requirements: *i*) a good spectrometer resolution to separate the nearby missing mass peak due to $(ep \rightarrow ep\pi^0)$; *ii*) measurements of small absolute cross sections with good accuracy, implying a precise calculation of the solid angle; *iii*) a careful study of experimental cuts (data are not background free).

Table 1: Status of performed VCS experiments.

ref.	Lab	Q^2 (GeV ²)	CM energy	data taking	status
[6]	MAMI (I)	0.33	$< (m_N + m_\pi)$	1995+97	published
[7]	JLab E93-050	0.9, 1.8	up to 1.9 GeV	1998	final stage
[8]	Bates E97-03	0.05	$< (m_N + m_\pi)$	2000	analysis
[9]	Bates E97-05	0.12	at $\Delta(1232)$	2001	analysis
[10]	MAMI, \vec{e} (II)	0.33	at $\Delta(1232)$	2002-2003	analysis

Figure 1 shows the fit of the structure functions using the Low Energy Theorem approach, in the experiments performed at MAMI [6] and in the Hall A of the Thomas Jefferson National Accelerator Facility (JLab) [11]. The reasonable goodness of the fit confirms the validity of the LET even at high Q^2 (~ 0.9 and 1.8 GeV² at JLab). Figure 2 shows $(ep \rightarrow ep\gamma)$ cross sections from the JLab experiment [12], scanning the resonance region for the first time. The data are well reproduced by the DR model in the region of the $\Delta(1232)$ resonance. These data are used to fit the free parameters of the model and subsequently extract the structure functions $P_{LL} - \frac{1}{\epsilon}P_{TT}$ and P_{LT} at $Q^2 = 1$ GeV².

Figure 3 gives a representation of the structure functions measured by the various experiments as a function of Q^2 . On the left plot the structure function P_{LL} is obtained from the measurement of $(P_{LL} - P_{TT}/\epsilon)$ by adding the value of P_{TT}/ϵ calculated in the DR model, at the value of Q^2 and ϵ of each experiment ¹. After dividing by the proton electric form factor G_E^p , one gets the Q^2 -behavior of the Generalized Polarizability $\alpha_E(Q^2)$, up to a constant factor. On the right plot, P_{LT}/G_E^p is proportional to the Generalized Polarizability $\beta_M(Q^2)$, neglecting a small term containing the spin GPs. The JLab data [13] are in good agreement with the DR model (and parameter values of the dashed curve). At lower Q^2 , the Chiral Perturbation Theory calculation to order p^3 [14] is in good agreement with the MAMI data for P_{LT} and $P_{LL} - P_{TT}/\epsilon$. Measurements tend to confirm the turn-over of P_{LT} that is predicted by most models in the low Q^2 region. This turn-over reflects the competing effects of para- and dia-magnetism in $\beta_M(Q^2)$, and the expected Bates measurement at $Q^2 = 0.05$ GeV² [8] is of great interest in this regard.

¹This is a model-dependent operation, since P_{TT} has not been measured yet.




Figure 1: A graphical representation of the polarizability fit in the LET method. The ordinate is the quantity $(d^5\sigma^{EXP} - d^5\sigma^{BH+Born})/(\phi q')/v_2$ extrapolated to $q' = 0$ in each bin in (θ, φ) . In abscissa is the ratio of the kinematic factors v_1 and v_2 . The fitted slope yields $P_{LL} - P_{TT}/\epsilon$ and the intercept yields the P_{LT} structure function.

4 Future prospects and conclusions

Table 2 summarizes the foreseen experiments in the field of VCS at low energy. Most projects aim at a deeper disentangling of the individual GPs, by polarization measurements or Rosenbluth-type separations.

Table 2: Experimental prospects of VCS at low energies. *SSA* (*DSA*)= single (double) spin asymmetry.

type of experiment	Lab	meas. \rightarrow structure functions	\sqrt{s}	Q^2 (GeV ²)
double polarization	MAMI & Bates	<i>DSA</i> \rightarrow separate the six GPs	$< \pi N$	0.33
		<i>DSA</i>	$< \pi N$	0.20
polarized beam	JLab	$d^5\sigma \rightarrow P_{LL} - P_{TT}/\epsilon$ and P_{LT} <i>SSA</i> \rightarrow test inputs to DR model	Δ	≤ 4
unpolarized	MAMI & Bates	$d^5\sigma \rightarrow \epsilon$ separation $\rightarrow P_{TT}, P_{LL}$	$< \pi N$	0.33
			$< \pi N$	0.20

Virtual Compton Scattering is an active field of research; Generalized Polarizabilities are new observables providing an original way to study nucleon structure, and there is an ongoing effort to learn more about them both experimentally and theoretically. Experiments make use of low and high energy machines (Bates, MAMI, JLab), of polarization degrees of freedom, and they exploit the versatility of methods to extract GPs at low and high Q^2 . These observables are also predicted by many theoretical approaches: chiral perturbation theory [14, 15], quark models [16], linear σ -model [17], dispersion relations [5], effective lagrangian model [18], ..., most calculations being valid at rather low Q^2 . In that view the results of the JLab VCS E93-050 experiment should stimulate new calculations of the GPs at high Q^2 .

Acknowledgements





Figure 2: Photon electroproduction cross section $d^5\sigma/dk'_{lab}d[\Omega_e]_{lab}d[\Omega_p]_{CM}$ measured in the JLab experiment [12] for six bins in φ , at $Q^2 = 1 \text{ GeV}^2$ and $\cos\theta_{\gamma^*\gamma_{CM}} = -0.975$. The solid curve is the DR model prediction, and the dashed curve is the (BH+Born) calculation.

This work was supported by DOE, NSF, by contract DE-AC05-84ER40150 under which the Southeastern Universities Research Association (SURA) operates the Thomas Jefferson National Accelerator Facility for DOE, by the CEA, the CNRS-IN2P3 and the Conseil Régional d'Auvergne (France), the FWO-Flanders and the BOF-Gent University (Belgium) and by the European Commission ERB FMRX-CT96-0008. I thank all my VCS colleagues for stimulating work.

References

- [1] V. Olmos de León *et al.*, Eur. Phys. J. A 10 (2001) 207 and references therein.
- [2] P.A.M. Guichon, G.Q. Liu and A.W. Thomas, Nucl. Phys. A 591 (1995) 606.
- [3] P.A.M. Guichon and M. Vanderhaeghen, Prog. Part. Nucl. Phys. 41 (1998) 125.
- [4] D. Drechsel *et al.*, Phys. Rev. C 55 (1997) 424; Phys. Rev. C 57 (1998) 941.
- [5] B. Pasquini *et al.*, Eur. Phys. J. A 11 (2001) 185; D. Drechsel, B. Pasquini and M. Vanderhaeghen, Phys. Rept 378 (2003) 99.
- [6] J. Roche *et al.*, Phys. Rev. Lett. 85 (2000) 708.
- [7] P.Y. Bertin, C. Hyde-Wright, P. Guichon *et al.*, CEBAF Proposal PR93050 (1993). <http://hallaweb.jlab.org/experiment/E93-050/vcs.html>



proton EM Form factors = parametrization of Brash et al., Phys Rev C 65 (2002) 051001

Figure 3: World results on VCS structure functions (statistical errors only). The data points at $Q^2 = 0$ are deduced from the TAPS experiment [1], the ones at $Q^2 = 0.33$ GeV^2 are from MAMI [6] and the other ones are from the JLab VCS E93-050 experiment including both analysis methods [11, 12]. Some points have been shifted in abscissa for clarity. The dash-dotted curve is the calculation of Chiral Perturbation Theory [14], the solid and dashed curves are the DR model predictions for different values of the free parameters (Λ_α and Λ_β , related to the GPs $\alpha_E(Q^2)$ and $\beta_M(Q^2)$; see [5] for more details).

- [8] J. Shaw, R. Miskimen *et al.*, MIT-Bates Proposal E97-03 (1997).
- [9] N. Kaloskamis, C. Papanicolas *et al.*, MIT-Bates Proposal E97-05 (1997).
- [10] H. Merkel, N. D'Hose, spokespersons, MAMI Proposal (2000).
- [11] S. Jaminion *et al.*, in preparation.
- [12] G. Laveissière *et al.*, in preparation.
- [13] H. Fonvieille, JLab E93-050 preliminary results in Proc. Baryons 2002 Conf. (World Scientific, 2003), hep-ex/0206035.
- [14] T. Hemmert *et al.*, Phys. Rev. D 62 (2000) 014013.
- [15] C.W. Kao and M. Vanderhaeghen, Phys. Rev. Lett. 89 (2002) 272002.
- [16] G.Q. Liu, A.W. Thomas and P.A.M. Guichon, Austral. J. Phys. 49 (1996) 905; B. Pasquini, S. Scherer and D. Drechsel, Phys. Rev. C 63 (2001) 025205.
- [17] A. Metz and D. Drechsel, Z. Phys. A 356 (1996) 351; Z. Phys. A 359 (1997) 165.
- [18] M. Vanderhaeghen, Phys. Lett. B 368 (1996) 13.