



Parametrization of the deuteron wave function of the paris n-n potential

M. Lacombe, B. Loiseau, R. Vinh Mau, J. Cote, P. Pires, R. de Tournell

► To cite this version:

M. Lacombe, B. Loiseau, R. Vinh Mau, J. Cote, P. Pires, et al.. Parametrization of the deuteron wave function of the paris n-n potential. Physics Letters B, 1981, 101, pp.139-140. 10.1016/0370-2693(81)90659-6 . in2p3-00017487

HAL Id: in2p3-00017487

<https://hal.in2p3.fr/in2p3-00017487>

Submitted on 25 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

PARAMETRIZATION OF THE DEUTERON WAVE FUNCTION OF THE PARIS N–N POTENTIAL

M. LACOMBE¹, B. LOISEAU and R. VINH MAU

*Division de Physique Théorique², Institut de Physique Nucléaire, Orsay 91406, France
and LPTPE, Université Pierre et Marie Curie, Paris 75230, France*

J. CÔTÉ and P. PIRÉS

Division de Physique Théorique, Institut de Physique Nucléaire, Orsay 91406, France

and

R. de TOURREIL

Laboratoire GANIL, F-14201 Caen, France

Received 22 January 1981

We present a convenient analytical parametrization, in both configuration and momentum spaces, of the deuteron wavefunction calculated with the Paris potential.

The deuteron wavefunction of the recently published Paris potential [1] reproduces quite well the known low energy properties and the electromagnetic form factor $A(q^2)$ of the deuteron. In particular the predicted value of the asymptotic D- to S-wave ratio η , 0.02608, is in excellent agreement with the recent high precision measurements of ref. [2], 0.0259 ± 0.0007 , and ref. [3], 0.02649 ± 0.00043 . It has been shown furthermore by Klarsfeld et al. [4] that for internucleon distances $r \gtrsim 0.6$ fm, this wavefunction satisfies rigorous bounds imposed by the experimental deuteron data and now improved by the new experimental values of η [2,3].

In view of the use of this wavefunction in low and medium energy nuclear reaction problems, numerical values have been given in table VI of ref. [1]. We have also found that a cubic spline interpolation of this table reproduces very accurately the original wavefunction. The numerical values and the cubic spline interpolation subroutine can be obtained on request. However, since in some cases an analytical form is use-

ful, we present here an algebraic parametrization of the Paris-potential deuteron wavefunction, in both configuration and momentum spaces.

It is appropriate to parametrize the radial wavefunction components $U(r)/r$ and $W(r)/r$ as a discrete superposition of Yukawa-type terms since this was done for the potential itself. Therefore $U(r)$ and $W(r)$ of table VI of ref. [1] are parametrized as follows

$$U_a(r) = \sum_{j=1}^n C_j \exp(-m_j r), \quad (1)$$

$$W_a(r) = \sum_{j=1}^n D_j \exp(-m_j r) (1 + 3/m_j r + 3/m_j^2 r^2),$$

where the subscript a refers to "analytical".

The corresponding momentum space wavefunction components are

$$U_a(p)/p = (2/\pi)^{1/2} \sum_{j=1}^n C_j / (p^2 + m_j^2), \quad (2)$$

$$W_a(p)/p = (2/\pi)^{1/2} \sum_{j=1}^n D_j / (p^2 + m_j^2).$$

¹ Faculté des Sciences de Reims, 51-Reims, France.

² Laboratoire associé au CNRS.

The boundary conditions $U_a(r) \rightarrow r$ and $W_a(r) \rightarrow r^3$ as $r \rightarrow 0$ imply

$$\sum_{J=1}^n C_J = 0, \quad \sum_{J=1}^n D_J/m_J^2 = \sum_{J=1}^n D_J = \sum_{J=1}^n D_J m_J^2 = 0. \quad (3)$$

In order to satisfy these constraints with accuracy the last coefficient of $U_a(r)$ and the last three of $W_a(r)$ are to be computed with the following formulae

$$C_n = - \sum_{J=1}^{n-1} C_J,$$

$$D_{n-2} = \frac{m_{n-2}^2}{(m_n^2 - m_{n-2}^2)(m_{n-1}^2 - m_{n-2}^2)} \times \left(-m_{n-1}^2 m_n^2 \sum_{J=1}^{n-3} \frac{D_J}{m_J^2} + (m_{n-1}^2 + m_n^2) \right. \\ \left. \times \sum_{J=1}^{n-3} D_J - \sum_{J=1}^{n-3} D_J m_J^2 \right) \quad (4)$$

and two other relations deduced by circular permutation of $n-2$, $n-1$ and n . The masses m_J are chosen to be $m_J = \alpha + (J-1)m_0$, with $m_0 = 1 \text{ fm}^{-1}$ and

$$\alpha = (2m_R |E_D|)^{1/2} / \hbar = 0.23162461 \text{ fm}^{-1},$$

where m_R and E_D are the neutron proton reduced mass and the deuteron binding energy of ref. [1]. This choice ensures the correct asymptotic behavior.

The values of the parameters C_J and D_J are listed in table 1. The accuracy of this parametrization is illustrated by the magnitude of

$$I_S = \left(\int_0^\infty dr [U(r) - U_a(r)]^2 \right)^{1/2} = 3.5 \times 10^{-4}$$

Table 1

Coefficients of the parametrized deuteron wavefunction components. The last C_J and the last three D_J are to be computed from eq. (4). The notation $a \pm n$ stands for $a \times 10^{\pm n}$.

$C_J (\text{fm}^{-1/2})$	$D_J (\text{fm}^{-1/2})$
0.88688076 + 00	0.23135193 - 01
-0.34717093 + 00	-0.85604572 + 00
-0.30502380 + 01	0.56068193 + 01
0.56207766 + 02	-0.69462922 + 02
-0.74957334 + 03	0.41631118 + 03
0.53365279 + 04	-0.12546621 + 04
-0.22706863 + 05	0.12387830 + 04
0.60434469 + 05	0.33739172 + 04
-0.10292058 + 06	-0.13041151 + 05
0.11223357 + 06	0.19512524 + 05
-0.75925226 + 05	see eq. (4)
0.29059715 + 05	see eq. (4)
see eq. (4)	see eq. (4)

and

$$I_D = \left(\int_0^\infty dr [W(r) - W_a(r)]^2 \right)^{1/2} = 4.0 \times 10^{-4}.$$

We have also checked that this parametrization reproduces very well the deuteron parameters and form factor obtained in ref. [1].

References

- [1] M. Lacombe et al., Phys. Rev. C21 (1980) 861.
- [2] W. Grüebler, V. König, P.A. Schmelzbach, B. Jenny and F. Sperisen, Phys. Lett. 92B (1980) 279.
- [3] K. Stephenson and W. Haeberli, Phys. Rev. Lett. 45 (1980) 520.
- [4] S. Klarsfeld, J. Martorell and D.W.L. Sprung, Nucl. Phys. A352 (1981) 113;
S. Klarsfeld, private communication.