



**HAL**  
open science

## Competition between collective and noncollective excitation modes at high spin in $^{124}\text{Ba}$

A. Al-Khatib, A.K. Singh, H. Hubel, P. Bringel, A. Burger, J. Domscheit, A. Neusser-Neffgen, G. Schoenwasser, G.B. Hagemann, C.R. Hansen, et al.

► **To cite this version:**

A. Al-Khatib, A.K. Singh, H. Hubel, P. Bringel, A. Burger, et al.. Competition between collective and noncollective excitation modes at high spin in  $^{124}\text{Ba}$ . *Physical Review C*, 2006, 74, pp.014305. 10.1103/PhysRevC.74.014305 . in2p3-00090258

**HAL Id: in2p3-00090258**

**<https://hal.in2p3.fr/in2p3-00090258>**

Submitted on 4 Jan 2007

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Competition between collective and noncollective excitation modes at high spin in $^{124}\text{Ba}$

A. Al-Khatib, A.K. Singh\*, H. Hübel, P. Bringel, A. Bürger,  
J. Domscheit, A. Neußer-Neffgen, and G. Schönwaßer  
*Helmholtz-Institut für Strahlen- und Kernphysik,  
Universität Bonn, Nussallee 14-16, D-53115 Bonn, Germany*

G.B. Hagemann, C. Ronn Hansen, B. Herskind, G. Sletten, and J.N. Wilson†  
*Niels Bohr Institute, Blegdamsvej 17, DK-2100 Copenhagen Ø, Denmark*

J. Timár, A. Algora, Zs. Dombrádi, J. Gál, G. Kalinka,  
J. Molnár, B.M. Nyakó, D. Sohler, and L. Zolnai  
*Institute of Nuclear Research of the Hungarian  
Academy of Sciences, H-4001 Debrecen, Hungary*

R. Clark, M. Cromaz, P. Fallon, I.Y. Lee, A.O. Macchiavelli, and D. Ward  
*Nuclear Science Division, Lawrence Berkeley National Laboratory, Berkeley CA 94720, USA*

H. Amro‡ and W.C. Ma  
*Department of Physics, Mississippi State University, Mississippi State, MS 39762, USA*

M. Kmiecik, A. Maj, J. Styczen, and K. Zuber  
*Niewodniczanski Institute of Nuclear Physics,  
Polish Academy of Sciences, PL-31342 Krakow, Poland*

K. Hauschild, A. Korichi, A. Lopez-Martens, J. Rocaaz, and S. Siem§  
*CSNSM Orsay, IN2P3/CNRS, F-91405 Orsay, France*

F. Hannachi and J.N. Scheurer  
*Centre d'Etudes Nucléaires de Bordeaux-Gradignan, F-33175 Gradignan, France*

P. Bednarczyk¶, Th. Byrski, D. Curien, O. Dorvaux, G.

---

\* Present address: Department of Physics, IIT Kharagpur, Kharagpur - 721302, India

† Present address: LPSC/IN2P3, F-38026 Grenoble Cedex, France

‡ Present address: Department of Physics, Univ. of Notre Dame, Notre Dame, IN 46556-5670, USA

§ Permanent address: Department of Physics, University of Oslo, N-0316 Oslo, Norway

¶ Permanent address: Niewodniczanski Institute of Nuclear Physics, Polish Academy of Sciences, PL-31342 Krakow, Poland. Present address: GSI, D-64291 Darmstadt, Germany

Duchêne, B. Gall, F. Khalfallah, I. Piqueras, and J. Robin  
*Institut de Recherches Subatomiques,  
CNRS-IN2P3, F-67037 Strasbourg, France*

A. Görgen  
*DAPNIA/SPhN, CEA-Saclay, F-91191 Gif-sur-Yvette, France*

K. Juhász  
*Department of Information Technology, Faculty of Informatics,  
University of Debrecen, H-4032 Debrecen, Hungary*

S.B. Patel  
*Department of Physics, University of Mumbai, Mumbai, India*

A.O. Evans and G. Rainovski  
*Oliver Lodge Laboratory, University of Liverpool, Liverpool L69 7ZE, UK*

G. Benzoni, A. Bracco, F. Camera, S. Leoni, P. Mason,  
B. Million, A. Paleni, R. Sacchi, and O. Wieland  
*Dipartimento di Fisica, Università degli Studi di Milano and INFN,  
Sezione di Milano, via Celoria 16, 20133 Milano, Italy*

C.M. Petrache and D. Petrache  
*Dipartimento di Fisica, Università di Camerino and INFN,  
Sezione di Perugia, I-62032 Camerino, Italy*

G. La Rana and R. Moro  
*Physical Science Department and INFN,  
Complesso Universitario di Monte S. Angelo, via Cinthia, I-80126 Napoli, Italy*

G. De Angelis  
*Istituto Nazionale di Fisica Nucleare, Laboratori Nazionali de Legnaro,  
via Romea 4, I-35020 Legnaro, Italy*

J.C. Lisle  
*Schuster Laboratory, University of Manchester,  
Brunswick Street, Manchester M13 9PL, UK*

B. Cederwall and K. Lagergen

*Department of Physics, Royal Institute of Technology, SE-10691 Stockholm, Sweden*

R.M. Lieder, E. Podsvirova, W. Gast, and H. Jäger

*Institut für Kernphysik, Forschungszentrum Jülich, D-52425 Jülich, Germany*

N. Redon

*IPN Lyon, IN2P3/CNRS, Université de Lyon-1, F-69622 Villeurbanne, France*

(Dated: February 9, 2006)

## Abstract

High-spin states in  $^{124}\text{Ba}$  were investigated in two experiments using the  $^{64}\text{Ni}(^{64}\text{Ni},4n)^{124}\text{Ba}$  reaction at three different beam energies. In-beam  $\gamma$ -ray coincidences were measured with the Euroball and Gammasphere detector arrays, respectively. In the experiment with Euroball, the CsI detector array Diamant was employed to discriminate against charged-particle channels. Six new rotational bands were observed in  $^{124}\text{Ba}$  and previously known bands were extended to higher spins and in the low-spin region. One of the bands shows a transition from collective to non-collective behavior at high spins. Configuration assignments are suggested on the basis of comparison with cranked shell model and cranked Nilsson-Strutinsky calculations.

PACS numbers: 23.20.Lv, 23.20.En, 27.60.+j, 21.60.Ev

## I. INTRODUCTION

Transitional nuclei between spherical and strongly deformed regions of the nuclear chart are usually soft with respect to deformation changes. Shape changes, including triaxiality, can be induced by the excitation of nucleons into specific deformation-driving orbitals. In the mass region around  $A = 125$ , both protons and neutrons can occupy the unique-parity  $h_{11/2}$  intruder orbitals which play an important role in driving the nuclear shape. The proton Fermi surface lies in the lower part of the  $h_{11/2}$  subshell, which favors prolate shape, whereas the neutron Fermi surface lies in the middle or upper part of the  $h_{11/2}$  subshell, which favors oblate shape. Thus, a coexistence of different shapes is expected in these nuclei due to opposite shape-driving forces of protons and neutrons in  $h_{11/2}$  orbitals.

Several studies of the development of the shape and collectivity as a function of neutron and proton number have been made in this mass region, e.g. for Xe, Ba and Ce nuclei [1–18]. It was found that the deformation increases with increasing proton number towards  $Z = 66$  whereas it decreases with increasing neutron number towards the  $N = 82$  shell closure. Total routhian surface (TRS) calculations show that after the proton  $h_{11/2}$  alignment the nuclei remain prolate ( $\gamma \simeq 0^\circ$ ) whereas the alignment of  $h_{11/2}$  neutrons favors a triaxial shape with  $\gamma \simeq -30^\circ$  [11, 14].

For nuclei with a small number of particles outside the semi-closed core with  $Z = 50$  and  $N = 64$ , the alignment of these particles along the rotation axis at high spin can polarize the nuclei towards oblate shape. In this spin range the rotational bands loose collectivity and single-particle alignments are favored as an efficient way to generate angular momentum. The collective bands terminate when all the single-particle angular momenta involved are aligned along the rotation axis. Higher-spin states can then be generated only by breaking the core.

Transitions from prolate to oblate shape have been observed to take place in two different ways, as smooth band termination, e.g. in the  $A \simeq 110$  region [19], and as sudden band termination, e.g. in the  $A \simeq 160$  region [20]. In the  $A = 125$  region the low-energy level structure is dominated by collective rotational bands with a moderate prolate deformation ( $\varepsilon_2 \simeq 0.25$  and  $\gamma \simeq 0^\circ$ ). However, non-collective excitations have been found to compete with collective rotation at high spin, e.g. in Xe [5] and Cs [21] isotopes, and may become yrast when the proton number increases. One of the aims of the present work was to explore such features in  $^{124}\text{Ba}$ . Eight rotational bands were previously known in this nucleus [12]. In this work, they were extended to much higher spins and partly also in the low-spin region. In addition, six new bands were discovered and a transition to non-collective excitations was observed in the yrast band. A preliminary report on this work has been given at a recent

conference [22].

In the following section the experimental details and the data analysis are described, followed by a presentation of the experimental results and the level scheme in section III. In section IV the configurations of the observed structures are discussed and a brief summary is given in the last section.

## II. EXPERIMENTAL DETAILS AND DATA ANALYSIS

Two experiments were performed in which high-spin states in  $^{124}\text{Ba}$  were populated using the  $^{64}\text{Ni}(^{64}\text{Ni},4n)$  reaction at beam energies of 255, 261 and 265 MeV, respectively. The higher beam energies were chosen to enhance the population of very high-spin structures since the primary goal of both experiments was to search for hyperdeformation. The search for hyperdeformation is still in progress [23]. A number of different reaction channels are open at these beam energies, among them the  $4n$  channel leading to  $^{124}\text{Ba}$  which is the strongest of the neutron-evaporation channels. In this work, we focus on the normal-deformed states in this nucleus.

The first experiment was carried out at Lawrence Berkeley National Laboratory. The  $^{64}\text{Ni}$  beam of 265 MeV was provided by the 88-Inch cyclotron. Gamma-ray coincidences were measured using the Gammasphere spectrometer with 100 Compton-suppressed Ge detectors [24]. The target consisted of a  $^{64}\text{Ni}$  foil of  $476 \mu\text{g}/\text{cm}^2$  thickness, enriched to 96.5 %. Data were recorded with a trigger condition of six or more Ge detectors after Compton suppression and 15 or more 'modules' showing a signal. A module is defined as a Ge detector and the BGO scintillator of the suppression shields. After presorting and setting a prompt time window, a total of  $1.2 \cdot 10^9$  events with Ge fold  $\geq 4$  were obtained.

The second experiment was performed at the Institut de Recherches Subatomiques at Strasbourg. The  $^{64}\text{Ni}$  beams with energies of 255 and 261 MeV were provided by the Vivitron Tandem accelerator. A self supporting foil of  $\simeq 500 \mu\text{g}/\text{cm}^2$  thickness was used as a target. Gamma-ray coincidences were measured using the Euroball spectrometer [25, 26]. It consisted of 30 Compton-suppressed conventional (tapered) Ge detectors and 41 composite Ge detectors. Of the latter, the 26 Clover detectors are composed of four Ge crystals each and the 15 Cluster detectors consist of seven Ge crystals each. At the time of the experiment, 230 individual Ge crystals out of the total of 239 of the full array could be used. An 'inner ball' of 210 BGO scintillation detectors was used as a multiplicity filter to enhance the selection of high-spin events. In addition, the charged-particle array Diamant [27, 28] consisting of 84 CsI scintillation detectors covering a geometrical solid angle of 95 % was placed inside

the target chamber. Each crystal has a size of  $4.5 \times 4.5 \text{ mm}^2$  and a thickness of 3 mm. The CsI detectors were wrapped in  $1.5 \text{ }\mu\text{m}$  thick aluminized mylar foil for light reflection. In front of the forward detectors (up to  $67.5^\circ$ ), Ta foils with thicknesses between 5 and  $15 \text{ }\mu\text{m}$  were mounted to reduce the background from scattered beam particles and delta electrons. Coincidence events were recorded with the requirement that at least three Ge crystals and eleven BGO detectors of the inner ball were in coincidence.

In the off-line analysis, a presorting of the data which includes Compton suppression, pile-up rejection, gain matching and calibration of the Ge detectors was made. For the investigation of  $^{124}\text{Ba}$  the information from the Diamant detectors was used to reject events in which charged particles were detected. After setting a wide gate on the prompt time peaks,  $\sim 3 \cdot 10^9$  events of three- and higher-fold Ge-detector coincidences remained. The coincidence events from both experiments were unpacked and sorted off-line into two-, three- and four-dimensional arrays (matrices, cubes and hypercubes, respectively) using the RADWARE program package [29].

To obtain information on the multipole order of  $\gamma$ -ray transitions, an asymmetric matrix was sorted with events registered in the detectors at forward (average angle  $35^\circ$ ) and backward (average angle  $156^\circ$ ) angles on one axis and those detected near- $90^\circ$  on the other axis. Directional correlation (DCO) ratios, defined as  $R_{DCO} = I(\gamma_2^{fb}, \gamma_1^{90^\circ})/I(\gamma_2^{90^\circ}, \gamma_1^{fb})$ , where  $I(\gamma_2, \gamma_1)$  denotes the intensity of  $\gamma_2$  in the spectrum gated on  $\gamma_1$ , were determined by setting gates on stretched E2 transitions. For a multipolarity determination of low-intensity transitions two angular correlation matrices were sorted. The first one contained events detected at forward and backward angles on one axis and those registered in all detectors on the other axis. Similarly, the second matrix contained events detected around  $90^\circ$  on one axis and those of all detectors on the other axis. Gates were set in these matrices on the axis with events detected in all detectors. The intensity ratio  $R_A = I(\gamma_2^{fb}, \gamma_1^{all})/I(\gamma_2^{90^\circ}, \gamma_1^{all})$  was used to distinguish between stretched dipole and stretched quadrupole transitions. Values of the DCO and the angular correlation ratios were around 0.6 and 1.0 for stretched dipole and stretched quadrupole transitions, respectively.

Linear-polarization measurements of  $\gamma$  rays linking the various bands to levels of known spin and parity were made to determine the parity of the bands. The Clover detectors of the Euroball spectrometer were used for this purpose. The reference plane of each  $\gamma$  ray is spanned by the beam axis and the emission direction of the  $\gamma$  ray. Compton-scattered events in neighboring pairs of Ge crystals of the Clover detectors, scattered horizontally (h) and vertically (v) to the reference plane, were analyzed separately. Two pairs of  $E_\gamma$ - $E_\gamma$  matrices were created with either horizontally or vertically scattered  $\gamma$  rays incremented on one axis and  $\gamma$  rays from all the detectors on the other axis. Such pairs of matrices were sorted with

gates on uncontaminated  $\gamma$ -ray transitions from the bands. The linear-polarization asymmetry ratio,  $A(E_\gamma) = [I_v(E_\gamma) - I_h(E_\gamma)]/[I_v(E_\gamma) + I_h(E_\gamma)]$ , was obtained by setting gates on  $\gamma$ -ray transitions in the two matrices. Here,  $I_v(E_\gamma)$  and  $I_h(E_\gamma)$  are the intensities of vertically and horizontally scattered  $\gamma$  rays of energy  $E_\gamma$ , respectively. The linear polarization,  $P$ , was calculated from this asymmetry ratio by  $P = A(E_\gamma) Q(E_\gamma)$ , where  $Q(E_\gamma)$  is the polarization sensitivity of the polarimeter. For point-like detectors this quantity can be calculated using the Klein-Nishina formula [30]. For finite crystal-size detectors it was obtained by introducing a scaling factor. The scaling factor has been measured for the Euroball array in several experiments. We have adopted the linear-polarization sensitivity obtained in ref. [31] for our analysis. The linear-polarization asymmetry for pure stretched magnetic transitions is negative, whereas for pure stretched electric transitions it is positive.

### III. RESULTS AND LEVEL SCHEME

The energy-level scheme of  $^{124}\text{Ba}$  is presented in Fig. 1. It is based on the results of the present work except for the lower-spin part which has been adopted from previous work [12, 18, 32]. Eight of the fourteen rotational bands were known previously. They were partly extended at low and high spins. The level scheme was constructed on the basis of coincidence relationships and relative  $\gamma$ -ray intensities. Spin assignments are based on DCO and angular correlation ratios. For transitions between low-spin states, no angular-correlation information could be obtained due to the loss of the spin alignment from the reaction. While the alignment stays approximately constant at high spins, it is reduced at low spins by the interaction between the nuclear moments and the fields produced by the atomic electrons. Transition energies and their relative intensities, DCO ratios, multipolarities and placements in the level scheme are listed in Table I. In several cases, in particular for Bands 12 and 13, it was not possible to obtain reliable intensities since connecting transitions to bands with well established intensities were too weak. However, in some of these cases DCO or angular correlation ratios could be determined, as they are derived from intensity ratios.

Band 1 was already established in previous work [12] up to the  $I^\pi = 34^+$  state. Above this level, we observe several new transitions resulting in an irregular level pattern. Fig. 2 shows two  $\gamma$ -ray coincidence spectra which demonstrate the forking of Band 1 above the 1444.3 keV transition and into the 1583.4 - 1610.9 keV sequence. The DCO ratio of the 1444.3 keV transition is compatible with a stretched quadrupole, presumably E2, transition. The multipolarity of the other new transitions could not be determined. Hence, the spin assignments to the high-spin levels are tentative and, therefore, given in parenthesis in Fig. 1.





Band 2, which was known up to the  $I^\pi = 24^+$  state [12], is extended by six transitions to higher spins. For the new transitions stretched quadrupole, probably E2, multipolarity was established up to 1261.2 keV. However, due to the regularity of the band, E2 multipolarity may be assumed also for the other high-spin transitions. Several new inter-band transitions connecting Band 2 with Band 1 have also been observed.




FIG. 2: Summed triple-gated  $\gamma$ -ray coincidence spectra of Band 1. The upper spectrum was produced by setting two gates on transitions (689.4 to 1444.3 keV) of Band 1 and one gate on the 1454.8 keV transition. The lower spectrum was obtained by setting two gates on transitions (689.4 to 1454.8 keV) of Band 1 and one gate on the 1444.3 keV transition.

TABLE I: Energies, relative intensities, DCO ratios, multipolarity and spin assignments of  $\gamma$ -ray transitions of  $^{124}\text{Ba}$

Excitation Energy <sup>a</sup> $E_i(\text{keV})$	Intensity <sup>b</sup> $E_\gamma(\text{keV})$	$I_\gamma$	$R_{DCO}^c$ ratio	$Band_i \rightarrow Band_f$	Assignment $J_i^\pi \rightarrow J_f^\pi$	Multipolarity
230	229.7	536(31)	-	$gsb \rightarrow gsb$	$2^+ \rightarrow 0^+$	E2
652	421.1	524(28)	-	$gsb \rightarrow gsb$	$4^+ \rightarrow 2^+$	E2
873	643.4	-	-	$8 \rightarrow gsb$	$2^+ \rightarrow 2^+$	M1
873	873.3	-	-	$8 \rightarrow gsb$	$2^+ \rightarrow 0^+$	E2
1162	510.0	-	-	$7 \rightarrow gsb$	$3^+ \rightarrow 4^+$	M1
1162	932.8	-	-	$7 \rightarrow gsb$	$3^+ \rightarrow 2^+$	M1
1228	576.5	500	0.90	$gsb \rightarrow gsb$	$6^+ \rightarrow 4^+$	E2
1325	451.7	-	-	$8 \rightarrow 8$	$4^+ \rightarrow 2^+$	E2
1325	673.1	-	-	$8 \rightarrow gsb$	$4^+ \rightarrow 4^+$	M1
1325	1094.5	-	-	$8 \rightarrow gsb$	$4^+ \rightarrow 2^+$	E2
1672	444.4	-	-	$7 \rightarrow gsb$	$5^+ \rightarrow 6^+$	M1
1672	510.0	-	-	$7 \rightarrow 7$	$5^+ \rightarrow 3^+$	E2
1672	1020.8	-	-	$7 \rightarrow gsb$	$5^+ \rightarrow 4^+$	M1
1722	1491.6	-	-	$4a \rightarrow gsb$	$(3^-) \rightarrow 2^+$	E1
1858	533.4	-	-	$8 \rightarrow 8$	$6^+ \rightarrow 4^+$	E2
1858	629.7	-	-	$8 \rightarrow gsb$	$6^+ \rightarrow 6^+$	M1
1913	684.9	-	-	$3 \rightarrow gsb$	$5^- \rightarrow 6^+$	E1
1913	1260.8	23(5)	0.74	$3 \rightarrow gsb$	$5^- \rightarrow 4^+$	E1
1923	694.7	348(22)	1.00	$gsb \rightarrow gsb$	$8^+ \rightarrow 6^+$	E2
2034	312.0	-	0.45	$4 \rightarrow 4a$	$4^- \rightarrow (3^-)$	M1
2034	1381.9	-	1.11	$4 \rightarrow gsb$	$4^- \rightarrow 4^+$	E1
2262	338.4	2(1)	-	$3 \rightarrow gsb$	$7^- \rightarrow 8^+$	E1
2262	348.4	6(1)	-	$3 \rightarrow 3$	$7^- \rightarrow 5^-$	E2

TABLE I: (continued)

Excitation Energy <sup>a</sup> $E_i$ (keV)	Intensity <sup>b</sup> $E_\gamma$ (keV)	$R_{DCO}^c$ $I_\gamma$	$Band_i \rightarrow Band_f$ ratio	Assignment $J_i^\pi \rightarrow J_f^\pi$	Multipolarity	
2262	1033.7	67(16)	0.66	$3 \rightarrow gsb$	$7^- \rightarrow 6^+$	E1
2267	354.0	-	-	$5a \rightarrow 3$	$5^- \rightarrow 5^-$	M1
2267	942.4	-	-	$5a \rightarrow 8$	$5^- \rightarrow 4^+$	E1
2267	1038.6	-	-	$5a \rightarrow gsb$	$5^- \rightarrow 6^+$	E1
2267	1615.3	6(2)	-	$5a \rightarrow gsb$	$5^- \rightarrow 4^+$	E1
2285	612.7	-	-	$7 \rightarrow 7$	$7^+ \rightarrow 5^+$	E2
2285	1057.0	-	-	$7 \rightarrow gsb$	$7^+ \rightarrow 6^+$	M1
2359	325.5	13(2)	1.32	$4 \rightarrow 4$	$6^- \rightarrow 4^-$	E2
2359	446.3	22(4)	0.35	$4 \rightarrow 3$	$6^- \rightarrow 5^-$	M1
2359	1130.0	31(7)	0.99	$4 \rightarrow gsb$	$6^- \rightarrow 6^+$	E1
2479	555.7	-	-	$8 \rightarrow gsb$	$8^+ \rightarrow 8^+$	M1
2479	620.9	-	-	$8 \rightarrow 8$	$8^+ \rightarrow 6^+$	E2
2498	230.5	64(12)	0.58	$6 \rightarrow 5a$	$6^- \rightarrow 5^-$	M1
2498	824.9	-	-	$6 \rightarrow 7$	$6^- \rightarrow 5^+$	E1
2647	288.0	2(1)	-	$5a \rightarrow 4$	$7^- \rightarrow 6^-$	M1
2647	380.4	5(2)	1.05	$5a \rightarrow 5a$	$7^- \rightarrow 5^-$	E2
2647	385.7	7(2)	0.67	$5a \rightarrow 3$	$7^- \rightarrow 7^-$	M1
2647	789.3	8(3)	-	$5a \rightarrow 8$	$7^- \rightarrow 6^+$	E1
2688	764.4	213(32)	0.97	$gsb \rightarrow gsb$	$10^+ \rightarrow 8^+$	E2
2691	193.0	39(14)	0.41	$5 \rightarrow 6$	$7^- \rightarrow 6^-$	M1
2705	345.2	46(8)	0.99	$4 \rightarrow 4$	$8^- \rightarrow 6^-$	E2
2705	442.7	6(3)	0.48	$4 \rightarrow 3$	$8^- \rightarrow 7^-$	M1
2705	781.0	29(6)	0.75	$4 \rightarrow gsb$	$8^- \rightarrow 8^+$	E1
2722	459.8	43(9)	1.13	$3 \rightarrow 3$	$9^- \rightarrow 7^-$	E2

TABLE I: (continued)

Excitation $E_i$ (keV)	Energy <sup>a</sup> $E_\gamma$ (keV)	Intensity <sup>b</sup> $I_\gamma$	$R_{DCO}^c$ ratio	$Band_i \rightarrow Band_f$	Assignment $J_i^\pi \rightarrow J_f^\pi$	Multipolarity
2722	798.0	75(8)	0.50	$3 \rightarrow gsb$	$9^- \rightarrow 8^+$	E1
2906	215.9	23(11)	0.35	$6 \rightarrow 5$	$8^- \rightarrow 7^-$	M1
2906	408.7	6(4)	-	$6 \rightarrow 6$	$8^- \rightarrow 6^-$	E2
2975	689.8	-	-	$7 \rightarrow 7$	$9^+ \rightarrow 7^+$	E2
2975	1053.0	-	-	$7 \rightarrow gsb$	$9^+ \rightarrow 8^+$	M1
3110	203.2	-	-	$5 \rightarrow 6$	$9^- \rightarrow 8^-$	M1
3110	404.9	6(2)	0.77	$5 \rightarrow 4$	$9^- \rightarrow 8^-$	M1
3110	419.2	15(3)	-	$5 \rightarrow 5$	$9^- \rightarrow 7^-$	E2
3110	462.0	14(5)	0.90	$5 \rightarrow 5a$	$9^- \rightarrow 7^-$	E2
3157	434.7	3(1)	-	$4 \rightarrow 3$	$10^- \rightarrow 9^-$	M1
3157	452.0	61(17)	1.02	$4 \rightarrow 4$	$10^- \rightarrow 8^-$	E2
3178	698.7	-	-	$8 \rightarrow 8$	$10^+ \rightarrow 8^+$	E2
3287	564.9	101(22)	1.09	$3 \rightarrow 3$	$11^- \rightarrow 9^-$	E2
3287	599.8	-	-	$3 \rightarrow gsb$	$11^- \rightarrow 10^+$	E1
3335	225.8	13(3)	-	$6 \rightarrow 5$	$10^- \rightarrow 9^-$	M1
3335	429.4	28(6)	1.01	$6 \rightarrow 6$	$10^- \rightarrow 8^-$	E2
3437	748.3	196(13)	0.97	$1 \rightarrow gsb$	$12^+ \rightarrow 10^+$	E2
3591	256.0	13(5)	-	$5 \rightarrow 6$	$11^- \rightarrow 10^-$	M1
3591	482.2	39(8)	0.95	$5 \rightarrow 5$	$11^- \rightarrow 9^-$	E2
3692	255.6	6(3)	0.74	$2 \rightarrow 1$	$12^+ \rightarrow 12^+$	M1
3692	1004.0	16(5)	1.17	$2 \rightarrow gsb$	$12^+ \rightarrow 10^+$	E2
3694	718.8	-	-	$7 \rightarrow 7$	$11^+ \rightarrow 9^+$	E2
3772	486.1	-	-	$4 \rightarrow 3$	$12^- \rightarrow 11^-$	M1
3772	615.5	56(16)	1.03	$4 \rightarrow 4$	$12^- \rightarrow 10^-$	E2

TABLE I: (continued)

Excitation Energy <sup>a</sup>	Intensity <sup>b</sup>	$R_{DCO}^c$	$Band_i \rightarrow Band_f$	Assignment	Multipolarity	
$E_i(\text{keV})$	$E_\gamma(\text{keV})$	$I_\gamma$	ratio	$J_i^\pi \rightarrow J_f^\pi$		
3891	299.9	16(4)	-	6 $\rightarrow$ 5	12 <sup>-</sup> $\rightarrow$ 11 <sup>-</sup>	M1
3891	556.4	13(5)	0.97	6 $\rightarrow$ 6	12 <sup>-</sup> $\rightarrow$ 10 <sup>-</sup>	E2
3968	681.0	94(25)	1.15	3 $\rightarrow$ 3	13 <sup>-</sup> $\rightarrow$ 11 <sup>-</sup>	E2
4127	434.2	-	-	1 $\rightarrow$ 2	14 <sup>+</sup> $\rightarrow$ 12 <sup>+</sup>	E2
4127	689.4	173(13)	1.03	1 $\rightarrow$ 1	14 <sup>+</sup> $\rightarrow$ 12 <sup>+</sup>	E2
4228	336.6	7(2)	-	5 $\rightarrow$ 6	13 <sup>-</sup> $\rightarrow$ 12 <sup>-</sup>	M1
4228	636.3	20(4)	1.15	5 $\rightarrow$ 5	13 <sup>-</sup> $\rightarrow$ 11 <sup>-</sup>	E2
4382	689.9	-	-	12 $\rightarrow$ 2	(11 <sup>+</sup> ) $\rightarrow$ 12 <sup>+</sup>	M1
4382	945.4	-	-	12 $\rightarrow$ 1	(11 <sup>+</sup> ) $\rightarrow$ 12 <sup>+</sup>	M1
4382	1407.0	-	-	12 $\rightarrow$ 7	(11 <sup>+</sup> ) $\rightarrow$ 9 <sup>+</sup>	E2
4408	281.3	3(1)	0.53	2 $\rightarrow$ 1	14 <sup>+</sup> $\rightarrow$ 14 <sup>+</sup>	M1
4408	715.5	14(4)	1.22	2 $\rightarrow$ 2	14 <sup>+</sup> $\rightarrow$ 12 <sup>+</sup>	E2
4408	971.1	24(6)	1.38	2 $\rightarrow$ 1	14 <sup>+</sup> $\rightarrow$ 12 <sup>+</sup>	E2
4534	566.2	-	-	4 $\rightarrow$ 3	14 <sup>-</sup> $\rightarrow$ 13 <sup>-</sup>	M1
4534	762.1	48(16)	1.03	4 $\rightarrow$ 4	14 <sup>-</sup> $\rightarrow$ 12 <sup>-</sup>	E2
4552	170.0	-	-	13 $\rightarrow$ 12	(12 <sup>+</sup> ) $\rightarrow$ (11 <sup>+</sup> )	M1
4552	722.0	-	0.74	13 $\rightarrow$ 13a	(12 <sup>+</sup> ) $\rightarrow$ (11)	-
4552	858.2	-	-	13 $\rightarrow$ 7	(12 <sup>+</sup> ) $\rightarrow$ 11 <sup>+</sup>	M1
4604	375.6	5(2)	-	6 $\rightarrow$ 5	14 <sup>-</sup> $\rightarrow$ 13 <sup>-</sup>	M1
4604	711.2	13(6)	1.00	6 $\rightarrow$ 6	14 <sup>-</sup> $\rightarrow$ 12 <sup>-</sup>	E2
4762	793.8	61(9)	1.03	3 $\rightarrow$ 3	15 <sup>-</sup> $\rightarrow$ 13 <sup>-</sup>	E2
4767	215.0	-	-	12 $\rightarrow$ 13	(13 <sup>+</sup> ) $\rightarrow$ (12 <sup>+</sup> )	M1
4767	384.2	-	1.14	12 $\rightarrow$ 12	(13 <sup>+</sup> ) $\rightarrow$ (11 <sup>+</sup> )	E2
4767	1073.2	-	-	12 $\rightarrow$ 7	(13 <sup>+</sup> ) $\rightarrow$ 11 <sup>+</sup>	E2

TABLE I: (continued)

Excitation $E_i$ (keV)	Energy <sup>a</sup> $E_\gamma$ (keV)	Intensity <sup>b</sup> $I_\gamma$	$R_{DCO}^c$ ratio	$Band_i \rightarrow Band_f$	Assignment $J_i^\pi \rightarrow J_f^\pi$	Multipolarity
4893	766.1	156(13)	1.05	1 $\rightarrow$ 1	16 <sup>+</sup> $\rightarrow$ 14 <sup>+</sup>	E2
5010	405.6	3(1)	-	5 $\rightarrow$ 6	15 <sup>-</sup> $\rightarrow$ 14 <sup>-</sup>	M1
5010	780.9	20(3)	0.92	5 $\rightarrow$ 5	15 <sup>-</sup> $\rightarrow$ 13 <sup>-</sup>	E2
5028	261.0	-	-	13 $\rightarrow$ 12	(14 <sup>+</sup> ) $\rightarrow$ (13 <sup>+</sup> )	M1
5028	476.0	-	1.05	13 $\rightarrow$ 13	(14 <sup>+</sup> ) $\rightarrow$ (12 <sup>+</sup> )	E2
5216	323.2	3(2)	0.78	2 $\rightarrow$ 1	16 <sup>+</sup> $\rightarrow$ 16 <sup>+</sup>	M1
5216	808.5	42(8)	1.35	2 $\rightarrow$ 2	16 <sup>+</sup> $\rightarrow$ 14 <sup>+</sup>	E2
5216	1090.0	11(5)	-	2 $\rightarrow$ 1	16 <sup>+</sup> $\rightarrow$ 14 <sup>+</sup>	E2
5330	302.0	-	-	12 $\rightarrow$ 13	(15 <sup>+</sup> ) $\rightarrow$ (14 <sup>+</sup> )	M1
5330	562.5	-	1.03	12 $\rightarrow$ 12	(15 <sup>+</sup> ) $\rightarrow$ (13 <sup>+</sup> )	E2
5330	1203.7	-	-	12 $\rightarrow$ 1	(15 <sup>+</sup> ) $\rightarrow$ 14 <sup>+</sup>	M1
5392	630.7	-	-	4 $\rightarrow$ 3	16 <sup>-</sup> $\rightarrow$ 15 <sup>-</sup>	M1
5392	858.3	31(11)	0.94	4 $\rightarrow$ 4	16 <sup>-</sup> $\rightarrow$ 14 <sup>-</sup>	E2
5446	436.3	2(1)	-	6 $\rightarrow$ 5	16 <sup>-</sup> $\rightarrow$ 15 <sup>-</sup>	M1
5446	842.6	9(2)	0.85	6 $\rightarrow$ 6	16 <sup>-</sup> $\rightarrow$ 14 <sup>-</sup>	E2
5639	878.0	51(12)	1.00	3 $\rightarrow$ 3	17 <sup>-</sup> $\rightarrow$ 15 <sup>-</sup>	E2
5669	339.0	-	-	13 $\rightarrow$ 12	(16 <sup>+</sup> ) $\rightarrow$ (15 <sup>+</sup> )	M1
5669	640.9	-	1.00	13 $\rightarrow$ 13	(16 <sup>+</sup> ) $\rightarrow$ (14 <sup>+</sup> )	E2
5726	832.0	48(9)	0.27	9 $\rightarrow$ 1	17 <sup>+</sup> $\rightarrow$ 16 <sup>+</sup>	M1
5764	871.6	84(15)	1.10	1 $\rightarrow$ 1	18 <sup>+</sup> $\rightarrow$ 16 <sup>+</sup>	E2
5906	459.8	-	-	5 $\rightarrow$ 6	17 <sup>-</sup> $\rightarrow$ 16 <sup>-</sup>	M1
5906	895.7	18(3)	1.12	5 $\rightarrow$ 5	17 <sup>-</sup> $\rightarrow$ 15 <sup>-</sup>	E2
6046	377.0	-	-	12 $\rightarrow$ 13	(17 <sup>+</sup> ) $\rightarrow$ (16 <sup>+</sup> )	M1
6046	715.5	-	1.01	12 $\rightarrow$ 12	(17 <sup>+</sup> ) $\rightarrow$ (15 <sup>+</sup> )	E2

TABLE I: (continued)

Excitation $E_i$ (keV)	Energy <sup>a</sup> $E_\gamma$ (keV)	Intensity <sup>b</sup> $I_\gamma$	$R_{DCO}^c$ ratio	$Band_i \rightarrow Band_f$	Assignment $J_i^\pi \rightarrow J_f^\pi$	Multipolarity
6081	317.2	-	-	2 $\rightarrow$ 1	18 <sup>+</sup> $\rightarrow$ 18 <sup>+</sup>	M1
6081	864.7	37(9)	0.93	2 $\rightarrow$ 2	18 <sup>+</sup> $\rightarrow$ 16 <sup>+</sup>	E2
6081	1187.9	-	-	2 $\rightarrow$ 1	18 <sup>+</sup> $\rightarrow$ 16 <sup>+</sup>	E2
6192	1299.0	-	-	10 $\rightarrow$ 1	18 <sup>+</sup> $\rightarrow$ 16 <sup>+</sup>	E2
6290	896.6	20(8)	1.21	4 $\rightarrow$ 4	18 <sup>-</sup> $\rightarrow$ 16 <sup>-</sup>	E2
6383	477.2	-	-	6 $\rightarrow$ 5	18 <sup>-</sup> $\rightarrow$ 17 <sup>-</sup>	M1
6383	938.1	9(2)	-	6 $\rightarrow$ 6	18 <sup>-</sup> $\rightarrow$ 16 <sup>-</sup>	E2
6454	408.0	-	-	13 $\rightarrow$ 12	(18 <sup>+</sup> ) $\rightarrow$ (17 <sup>+</sup> )	M1
6454	784.9	-	1.04	13 $\rightarrow$ 13	(18 <sup>+</sup> ) $\rightarrow$ (16 <sup>+</sup> )	E2
6556	917.3	29(6)	1.15	3 $\rightarrow$ 3	19 <sup>-</sup> $\rightarrow$ 17 <sup>-</sup>	E2
6583	819.1	7(2)	-	9 $\rightarrow$ 1	19 <sup>+</sup> $\rightarrow$ 18 <sup>+</sup>	M1
6583	857.7	22(9)	0.95	9 $\rightarrow$ 9	19 <sup>+</sup> $\rightarrow$ 17 <sup>+</sup>	E2
6705	1066.0	-	-	3a $\rightarrow$ 3	(18) $\rightarrow$ 17 <sup>-</sup>	-
6712	948.6	59(13)	1.06	1 $\rightarrow$ 1	20 <sup>+</sup> $\rightarrow$ 18 <sup>+</sup>	E2
6871	487.8	-	-	5 $\rightarrow$ 6	19 <sup>-</sup> $\rightarrow$ 18 <sup>-</sup>	M1
6871	965.0	7(2)	0.96	5 $\rightarrow$ 5	19 <sup>-</sup> $\rightarrow$ 17 <sup>-</sup>	E2
6898	444.0	-	-	12 $\rightarrow$ 13	(19 <sup>+</sup> ) $\rightarrow$ (18 <sup>+</sup> )	M1
6898	852.0	-	1.06	12 $\rightarrow$ 12	(19 <sup>+</sup> ) $\rightarrow$ (17 <sup>+</sup> )	E2
7000	919.4	25(6)	1.08	2 $\rightarrow$ 2	20 <sup>+</sup> $\rightarrow$ 18 <sup>+</sup>	E2
7000	1236.2	-	-	2 $\rightarrow$ 1	20 <sup>+</sup> $\rightarrow$ 18 <sup>+</sup>	E2
7084	891.9	20(10)	0.91	10 $\rightarrow$ 10	20 <sup>+</sup> $\rightarrow$ 18 <sup>+</sup>	E2
7084	1319.7	-	0.92	10 $\rightarrow$ 1	20 <sup>+</sup> $\rightarrow$ 18 <sup>+</sup>	E2
7230	940.5	13(6)	0.90	4 $\rightarrow$ 4	20 <sup>-</sup> $\rightarrow$ 18 <sup>-</sup>	E2
7365	910.0	-	1.08	13 $\rightarrow$ 13	(20 <sup>+</sup> ) $\rightarrow$ (18 <sup>+</sup> )	E2



TABLE I: (continued)

Excitation Energy <sup>a</sup>	Intensity <sup>b</sup>	$R_{DCO}^c$	$Band_i \rightarrow Band_f$	Assignment	Multipolarity	
$E_i(\text{keV})$	$E_\gamma(\text{keV})$	$I_\gamma$	ratio	$J_i^\pi \rightarrow J_f^\pi$		
7366	495.2	-	-	6 $\rightarrow$ 5	20 <sup>-</sup> $\rightarrow$ 19 <sup>-</sup>	M1
7366	982.6	4(2)	-	6 $\rightarrow$ 6	20 <sup>-</sup> $\rightarrow$ 18 <sup>-</sup>	E2
7502	945.3	16(4)	1.26	3 $\rightarrow$ 3	21 <sup>-</sup> $\rightarrow$ 19 <sup>-</sup>	E2
7503	791.3	-	-	9 $\rightarrow$ 1	21 <sup>+</sup> $\rightarrow$ 20 <sup>+</sup>	M1
7503	918.0	12(4)	0.87	9 $\rightarrow$ 9	21 <sup>+</sup> $\rightarrow$ 19 <sup>+</sup>	E2
7717	1004.9	41(10)	1.08	1 $\rightarrow$ 1	22 <sup>+</sup> $\rightarrow$ 20 <sup>+</sup>	E2
7865	966.9	-	1.16	12 $\rightarrow$ 12	(21 <sup>+</sup> ) $\rightarrow$ (19 <sup>+</sup> )	E2
7878	1007.4	-	-	5 $\rightarrow$ 5	21 <sup>-</sup> $\rightarrow$ 19 <sup>-</sup>	E2
7984	983.7	18(4)	1.09	2 $\rightarrow$ 2	22 <sup>+</sup> $\rightarrow$ 20 <sup>+</sup>	E2
7984	1272.3	-	-	2 $\rightarrow$ 1	22 <sup>+</sup> $\rightarrow$ 20 <sup>+</sup>	E2
8101	1016.8	11(5)	1.08	10 $\rightarrow$ 10	22 <sup>+</sup> $\rightarrow$ 20 <sup>+</sup>	E2
8101	1389.0	-	-	10 $\rightarrow$ 1	22 <sup>+</sup> $\rightarrow$ 20 <sup>+</sup>	E2
8263	1034.4	11(5)	0.91	4 $\rightarrow$ 4	22 <sup>-</sup> $\rightarrow$ 20 <sup>-</sup>	E2
8371	1006.3	-	1.06	13 $\rightarrow$ 13	(22 <sup>+</sup> ) $\rightarrow$ (20 <sup>+</sup> )	E2
8411	1044.0	-	-	6 $\rightarrow$ 6	22 <sup>-</sup> $\rightarrow$ 20 <sup>-</sup>	E2
8486	769.2	-	-	9 $\rightarrow$ 1	23 <sup>+</sup> $\rightarrow$ 22 <sup>+</sup>	M1
8486	983.4	10(4)	1.11	9 $\rightarrow$ 9	23 <sup>+</sup> $\rightarrow$ 21 <sup>+</sup>	E2
8512	1009.7	14(4)	0.95	3 $\rightarrow$ 3	23 <sup>-</sup> $\rightarrow$ 21 <sup>-</sup>	E2
8795	1077.1	37(9)	1.22	1 $\rightarrow$ 1	24 <sup>+</sup> $\rightarrow$ 22 <sup>+</sup>	E2
8905	1040.6	-	1.22	12 $\rightarrow$ 12	(23 <sup>+</sup> ) $\rightarrow$ (21 <sup>+</sup> )	E2
8913	1034.0	-	-	5 $\rightarrow$ 5	23 <sup>-</sup> $\rightarrow$ 21 <sup>-</sup> )	E2
9054	1069.2	12(3)	0.90	2 $\rightarrow$ 2	24 <sup>+</sup> $\rightarrow$ 22 <sup>+</sup>	E2
9179	1078.6	10(5)	0.84	10 $\rightarrow$ 10	24 <sup>+</sup> $\rightarrow$ 22 <sup>+</sup>	E2
9179	1462.3	-	-	10 $\rightarrow$ 1	24 <sup>+</sup> $\rightarrow$ 22 <sup>+</sup>	E2

TABLE I: (continued)

Excitation Energy <sup>a</sup>	Intensity <sup>b</sup>	$R_{DCO}^c$	$Band_i \rightarrow Band_f$	Assignment	Multipolarity	
$E_i(\text{keV})$	$E_\gamma(\text{keV})$	$I_\gamma$	ratio	$J_i^\pi \rightarrow J_f^\pi$		
9381	1117.6	5(1)	0.89	4 $\rightarrow$ 4	24 <sup>-</sup> $\rightarrow$ 22 <sup>-</sup>	E2
9430	1058.5	-	1.25	13 $\rightarrow$ 13	(24 <sup>+</sup> ) $\rightarrow$ (22 <sup>+</sup> )	E2
9528	1117.1	-	-	6 $\rightarrow$ 6	(24 <sup>-</sup> ) $\rightarrow$ (22 <sup>-</sup> )	E2
9566	1078.2	6(2)	1.25	9 $\rightarrow$ 9	25 <sup>+</sup> $\rightarrow$ 23 <sup>+</sup>	E2
9613	1100.1	8(2)	1.37	3 $\rightarrow$ 3	25 <sup>-</sup> $\rightarrow$ 23 <sup>-</sup>	E2
9918	1124.1	-	-	11 $\rightarrow$ 1	(25) $\rightarrow$ 24 <sup>+</sup>	-
9951	1154.5	15(6)	1.11	1 $\rightarrow$ 1	26 <sup>+</sup> $\rightarrow$ 24 <sup>+</sup>	E2
9976	1070.6	-	1.07	12 $\rightarrow$ 12	(25 <sup>+</sup> ) $\rightarrow$ (23 <sup>+</sup> )	E2
9984	1070.7	-	-	5 $\rightarrow$ 5	(25 <sup>-</sup> ) $\rightarrow$ (23 <sup>-</sup> )	E2
10223	1168.6	10(3)	0.97	2 $\rightarrow$ 2	26 <sup>+</sup> $\rightarrow$ 24 <sup>+</sup>	E2
10312	1131.4	5(2)	1.16	10 $\rightarrow$ 10	26 <sup>+</sup> $\rightarrow$ 24 <sup>+</sup>	E2
10312	1516.9	-	-	10 $\rightarrow$ 1	26 <sup>+</sup> $\rightarrow$ 24 <sup>+</sup>	E2
10521	1091.9	-	1.12	13 $\rightarrow$ 13	(26 <sup>+</sup> ) $\rightarrow$ (24 <sup>+</sup> )	E2
10563	1181.2	4(1)	-	4 $\rightarrow$ 4	26 <sup>-</sup> $\rightarrow$ 24 <sup>-</sup>	E2
10707	1178.4	-	-	6 $\rightarrow$ 6	(26 <sup>-</sup> ) $\rightarrow$ (24 <sup>-</sup> )	E2
10751	1184.8	2(2)	-	9 $\rightarrow$ 9	27 <sup>+</sup> $\rightarrow$ 25 <sup>+</sup>	E2
10814	1201.5	6(2)	1.29	3 $\rightarrow$ 3	27 <sup>-</sup> $\rightarrow$ 25 <sup>-</sup>	E2
11029	1151.3	-	-	11 $\rightarrow$ 11	(27) $\rightarrow$ (25)	E2
11079	1102.4	-	1.13	12 $\rightarrow$ 12	(27 <sup>+</sup> ) $\rightarrow$ (25 <sup>+</sup> )	E2
11118	1505.0	-	-	3b $\rightarrow$ 3	(26) $\rightarrow$ 25 <sup>-</sup>	-
11183	1231.2	11(4)	0.86	1 $\rightarrow$ 1	28 <sup>+</sup> $\rightarrow$ 26 <sup>+</sup>	E2
11475	1251.8	6.3(3.8)	0.97	2 $\rightarrow$ 2	28 <sup>+</sup> $\rightarrow$ 26 <sup>+</sup>	E2
11526	1214.2	3(2)	0.91	10 $\rightarrow$ 10	28 <sup>+</sup> $\rightarrow$ 26 <sup>+</sup>	E2
11651	1129.3	-	1.21	13 $\rightarrow$ 13	(28 <sup>+</sup> ) $\rightarrow$ (26 <sup>+</sup> )	E2

TABLE I: (continued)

Excitation $E_i$ (keV)	Energy <sup>a</sup> $E_\gamma$ (keV)	Intensity <sup>b</sup> $I_\gamma$	$R_{DCO}^c$ ratio	$Band_i \rightarrow Band_f$	Assignment $J_i^\pi \rightarrow J_f^\pi$	Multipolarity
11755	1192.0	2(1)	-	4 $\rightarrow$ 4	28 <sup>-</sup> $\rightarrow$ 26 <sup>-</sup>	E2
12035	1282.9	-	-	9 $\rightarrow$ 9	29 <sup>+</sup> $\rightarrow$ 27 <sup>+</sup>	E2
12118	1304.4	5(1)	1.02	3 $\rightarrow$ 3	29 <sup>-</sup> $\rightarrow$ 27 <sup>-</sup>	E2
12246	1164.9	-	-	12 $\rightarrow$ 12	(29 <sup>+</sup> ) $\rightarrow$ (27 <sup>+</sup> )	E2
12290	1221.2	-	-	11 $\rightarrow$ 11	(29) $\rightarrow$ (27)	E2
12492	1309.6	10(3)	1.27	1 $\rightarrow$ 1	30 <sup>+</sup> $\rightarrow$ 28 <sup>+</sup>	E2
12738	1261.2	4(1)	0.93	2 $\rightarrow$ 2	30 <sup>+</sup> $\rightarrow$ 28 <sup>+</sup>	E2
12826	1297.9	-	-	10 $\rightarrow$ 10	30 <sup>+</sup> $\rightarrow$ 28 <sup>+</sup>	E2
12862	1210.9	-	-	13 $\rightarrow$ 13	(30 <sup>+</sup> ) $\rightarrow$ (28 <sup>+</sup> )	E2
12961	1206.5	-	-	4 $\rightarrow$ 4	(30 <sup>-</sup> ) $\rightarrow$ (28 <sup>-</sup> )	E2
13350	1232.0	-	0.8 <sup>e</sup>	14 $\rightarrow$ 3	(30) $\rightarrow$ 29 <sup>-</sup>	-
13413	1377.0	-	-	9 $\rightarrow$ 9	31 <sup>+</sup> $\rightarrow$ 29 <sup>+</sup>	E2
13498	1249.5	-	-	12 $\rightarrow$ 12	(31 <sup>+</sup> ) $\rightarrow$ (29 <sup>+</sup> )	E2
13519	1400.8	4(2)	1.18	3 $\rightarrow$ 3	31 <sup>-</sup> $\rightarrow$ 29 <sup>-</sup>	E2
13592	1301.4	-	-	11 $\rightarrow$ 11	(31) $\rightarrow$ (29)	E2
13881	1388.6	9(3)	1.22	1 $\rightarrow$ 1	32 <sup>+</sup> $\rightarrow$ 30 <sup>+</sup>	E2
14063	1324.5	3.6(2.9)	-	2 $\rightarrow$ 2	32 <sup>+</sup> $\rightarrow$ 30 <sup>+</sup>	E2
14185	1224.3	-	-	4 $\rightarrow$ 4	32 <sup>-</sup> $\rightarrow$ 30 <sup>-</sup>	E2
14196	1370.1	-	-	10 $\rightarrow$ 10	32 <sup>+</sup> $\rightarrow$ 30 <sup>+</sup>	E2
14757	1238.0	-	0.94 <sup>e</sup>	14 $\rightarrow$ 3	(32) $\rightarrow$ 31 <sup>-</sup>	-
14757	1406.9	-	-	14 $\rightarrow$ 14	(32) $\rightarrow$ (30)	E2
14838	1340.6	-	-	12 $\rightarrow$ 12	(33 <sup>+</sup> ) $\rightarrow$ (31 <sup>+</sup> )	E2
14887	1474.7	-	-	9 $\rightarrow$ 9	33 <sup>+</sup> $\rightarrow$ 31 <sup>+</sup>	E2
14981	1389.1	-	-	11 $\rightarrow$ 11	(33) $\rightarrow$ (31)	E2

TABLE I: (continued)

Excitation Energy <sup>a</sup>	Intensity <sup>b</sup>	$R_{DCO}^c$	$Band_i \rightarrow Band_f$	Assignment	Multipolarity	
$E_i(\text{keV})$	$E_\gamma(\text{keV})$	$I_\gamma$	ratio	$J_i^\pi \rightarrow J_f^\pi$		
15007	1486.5	-	-	3 $\rightarrow$ 3	33 <sup>-</sup> $\rightarrow$ 31 <sup>-</sup>	E2
15336	1454.8	6(2)	1.38 <sup>d</sup>	1 $\rightarrow$ 1	34 <sup>+</sup> $\rightarrow$ 32 <sup>+</sup>	E2
15465	1401.6	-	-	2 $\rightarrow$ 2	34 <sup>+</sup> $\rightarrow$ 32 <sup>+</sup>	E2
15476	1290.9	-	-	4 $\rightarrow$ 4	(34 <sup>-</sup> ) $\rightarrow$ 32 <sup>-</sup>	E2
15624	1427.5	-	-	10 $\rightarrow$ 10	34 <sup>+</sup> $\rightarrow$ 32 <sup>+</sup>	E2
16031	1274.0	-	1.42 <sup>e</sup>	14 $\rightarrow$ 14	(34) $\rightarrow$ (32)	E2
16286	1448.0	-	-	12 $\rightarrow$ 12	(35 <sup>+</sup> ) $\rightarrow$ (33 <sup>+</sup> )	E2
16434	1544.2	-	-	9 $\rightarrow$ 9	(35 <sup>+</sup> ) $\rightarrow$ 33 <sup>+</sup>	E2
16463	1481.8	-	-	11 $\rightarrow$ 11	(35) $\rightarrow$ (33)	E2
16780	1444.3	5(2)	1.41 <sup>d</sup>	1 $\rightarrow$ 1	36 <sup>+</sup> $\rightarrow$ 34 <sup>+</sup>	E2
16918	1583.4	-	-	1c $\rightarrow$ 1	(36 <sup>+</sup> ) $\rightarrow$ 34 <sup>+</sup>	E2
16949	1484.3	-	-	2 $\rightarrow$ 2	36 <sup>+</sup> $\rightarrow$ 34 <sup>+</sup>	E2
17117	1493.3	-	-	10 $\rightarrow$ 10	(36 <sup>+</sup> ) $\rightarrow$ 34 <sup>+</sup>	E2
17437	1406.0	-	-	14 $\rightarrow$ 14	(36) $\rightarrow$ (34)	E2
18042	1579.7	-	-	11 $\rightarrow$ 11	(37) $\rightarrow$ (35)	E2
18055	1619.6	-	-	9 $\rightarrow$ 9	(37 <sup>+</sup> ) $\rightarrow$ (35 <sup>+</sup> )	E2
18075	1294.4	-	-	1a $\rightarrow$ 1	(38 <sup>+</sup> ) $\rightarrow$ 36 <sup>+</sup>	E2
18149	1368.4	-	-	1b $\rightarrow$ 1	(38 <sup>+</sup> ) $\rightarrow$ 36 <sup>+</sup>	E2
18529	1610.9	-	-	1c $\rightarrow$ 1c	(38 <sup>+</sup> ) $\rightarrow$ (36 <sup>+</sup> )	E2
18656	1537.7	-	-	10 $\rightarrow$ 10	(38 <sup>+</sup> ) $\rightarrow$ (36 <sup>+</sup> )	E2
18911	1474.0	-	-	14 $\rightarrow$ 14	(38) $\rightarrow$ (36)	E2
19722	1679.7	-	-	11 $\rightarrow$ 11	(39) $\rightarrow$ (37)	E2
20485	1574.0	-	-	14 $\rightarrow$ 14	(40) $\rightarrow$ (38)	E2
21503	1780.9	-	-	11 $\rightarrow$ 11	(41) $\rightarrow$ (39)	E2

TABLE I: (continued)

Excitation Energy <sup>a</sup>	Intensity <sup>b</sup>	$R_{DCO}^c$	$Band_i \rightarrow Band_f$	Assignment	Multipolarity	
$E_i(\text{keV})$	$E_\gamma(\text{keV})$	$I_\gamma$	ratio	$J_i^\pi \rightarrow J_f^\pi$		
22152	1666.8	-	-	14 $\rightarrow$ 14	(42) $\rightarrow$ (40)	E2
23386	1883.2	-	-	11 $\rightarrow$ 11	(43) $\rightarrow$ (41)	E2
25372	1986.3	-	-	11 $\rightarrow$ 11	(45) $\rightarrow$ (43)	E2

<sup>a</sup> Uncertainties of  $\gamma$ -ray energies are between 0.1 and 0.6 keV depending on their intensity.

<sup>b</sup> Intensities are normalized to the 576.5 keV transition with  $I_\gamma = 500$ .

<sup>c</sup> Uncertainties in the DCO ratios are between 0.03 and 0.2.

<sup>d</sup> Ratio obtained from angular distribution matrices.

<sup>e</sup> Uncertainty  $> 50\%$  due to low statistics.




FIG. 3: Sum of triple-gated  $\gamma$ -ray coincidence spectra of Band 9 (upper panel) and Band 10 (lower panel). The spectrum of Band 9 was produced by setting one gate on transitions (229.7 to 766.1 keV) of Band 1 and two gates on transitions (832 to 1619.6 keV) of Band 9. The spectrum of Band 10 was obtained with one gate on the same transitions of Band 1 and two gates on transitions (891.9 to 1537.7 keV) of Band 10. The single asterisks mark peaks of Band 1. The  $\gamma$  rays marked by double asterisks are the linking transitions from Bands 9 and 10, respectively, to Band 1.

Bands 3 to 8 were also known from previous work [12, 18]. We confirm these sequences and place three to four transitions on top of the previously known levels of Bands 3, 5 and 6. The low-spin part of the coupled Bands 5 and 6 was rearranged. Two close-lying  $7^-$  states, at 2647 and 2691 keV, respectively, were previously known [12]. The 2647 keV level was assigned as member of the  $\alpha = 1$  signature branch whereas the 2691 keV state was suggested not to belong to the band. Analyzing triple coincidences, we found a 419.2 keV transition from the  $9^-$  band member to the  $7^-$  level at 2691 keV which was previously not detected, probably because it lies so close in energy to the strong 421.1 keV  $4^+ \rightarrow 2^+$  transition. We rearrange the  $7^-$  states as shown in the level scheme of Fig. 1. In this way the coupled bands continue with  $\Delta I = 1$  and  $\Delta I = 2$  transitions down to the  $5^-$  state. The fairly strong transition of 462.0 keV to the  $7^-$  state at 2647 keV may be explained by mixing of the  $7^-$  levels.

Band 4 is extended by seven transitions. Bands 7 and 8 are weakly populated and could

not be extended to higher spin. DCO ratios measured for some of the in-band transitions as well as decay-out transitions are in accordance with the previous spin assignments [12]. For some high-spin transitions observed in this work we could not determine the multipolarity. However, E2 multipolarity has been assumed for the in-band transitions due to the regularity of the bands.

The two sequences of  $\gamma$  rays labeled Band 9 and 10 have been observed for the first time in our work. Both bands decay into levels of Band 1. Gamma-ray coincidence spectra for the two bands are shown in Fig. 3. The 832.0 keV transition, which we place at the bottom of Band 9, was observed already by Pilotte et al. [12]. The linear polarization asymmetry of this transition is small and negative which confirms the previously suggested mixed M1/E2 multipolarity for this transition which feeds into the  $16^+$  state of Band 1. For an electric dipole one would expect no mixing and, thus, a positive asymmetry. This leads to  $I^\pi = 17^+$  for the lowest level of Band 9.

Band 10 decays into Band 1 via several transitions between 1299 and 1517 keV. Of these only the 1319.7 keV transition is clean and strong enough to allow the determination of the angular correlation ratio. This ratio is compatible with stretched quadrupole, probably E2, multipolarity. Thus, we assign even spins and positive parity to Band 10. Only the in-band transitions near the bottom of Bands 9 and 10 are sufficiently strong for a determination of the angular correlation ratios. They are consistent with stretched quadrupole, probably E2, transitions.

The three new bands, Bands 11, 12 and 13, are only tentatively connected to lower-lying levels and their excitation energies, spins and parities remain uncertain. Gamma-ray coincidence spectra of these bands are displayed in Figs. 4 and 5, respectively. Band 11 is a cascade of eleven transitions, probably of E2 multipolarity, which decays into Band 1. Transitions of Band 1 are observed in coincidence with the new band up to the  $24^+$  level. The transitions of Band 11 show an energy difference which is smoothly increasing from 71 to 105 keV; only the difference between the 1124.1 and 1151.3 keV transitions is much lower. Therefore, the 1124.1 keV  $\gamma$  ray may be a link between the two bands. Other weak  $\gamma$ -ray lines observed in the coincidence spectra, see Fig. 4, with energies of 1095, 1453, 1516, 1833 and 1914 keV might also be candidates for inter-band transitions.

The two other new bands, Bands 12 and 13, are connected by weak inter-band transitions in the low-spin region and, thus, probably form a pair of signature-partner bands. They decay into Bands 1, 2 and 7. Of the linking transitions shown in the level scheme, Fig. 1, only the 1407 keV line is firmly established. Due to the low intensity its DCO ratio has a large uncertainty and the multipolarity could not be unambiguously determined. Hence, the spin assignment to the bands is tentative. The other transitions are weak and only tentatively




FIG. 4: Summed triple-gated coincidence spectrum of Band 11. The spectrum was produced by setting one gate on transitions (229.7 to 1077 keV) of Band 1 and two gates on transitions (1124.1 to 1986.3 keV) of Band 11. The asterisks mark peaks of Band 1. The energies in parentheses correspond to transitions which are observed in coincidence but could not be placed in the level scheme.

assigned. Transitions of Band 1 up to the  $10^+$  and possibly up to the  $14^+$  state are observed in coincidence with Bands 12 and 13, see Fig. 5.

Band 14 has been observed for the first time in this work. It decays into Band 3. Due to the low intensities of the linking transitions the DCO ratios have large uncertainties and their multiplicities could not be reliably determined. Since Band 14 feeds into the  $I^\pi = 29^-$  state of Band 3 we tentatively assign  $I = 30$  to the lowest level of the band.






FIG. 5: Summed double-gated and triple-gated coincidence spectra of Bands 12 and 13. The spectrum of Band 12 (upper panel) was produced by setting one gate on transitions (229.7 to 764.4 keV) of Band 1 and two gates on transitions (384.2 to 1448.0 keV) of Band 12. The spectrum of Band 13 (lower panel) was obtained by setting one gate on the same transitions of Band 1 and two gates on transitions (476 to 1163 keV) of Band 13. The single asterisks mark peaks of Band 1. Double asterisks mark the decay-out transitions. The energies in parentheses correspond to transitions which are observed in coincidence but could not be placed in the level scheme.

#### IV. DISCUSSION

In previous work [12, 18, 32, 33] configuration assignments have been made to the low- and medium-spin regions of Bands 1 to 8 of  $^{124}\text{Ba}$ . Arguments for the configuration assignments are the observed excitation energies, spins, parities, band-crossing frequencies and alignment gains. These quantities can be compared to those observed in neighboring nuclei as well

as to cranked shell model (CSM) [34, 35] and cranked Nilsson-Strutinsky (CNS) [19, 36] calculations. For the coupled bands, signature splittings and  $B(M1)/B(E2)$  ratios provide additional evidence for their configurations.

TABLE II: Shell model origin and quasiparticle labels for low-lying orbitals in  $^{124}\text{Ba}$  [12].

	Shell model states	Nilsson orbitals	$\alpha = +\frac{1}{2}$	$\alpha = -\frac{1}{2}$
Protons	$(d_{5/2}, g_{7/2})$	$[422]_{\frac{3}{2}}^{+}$	$a$	$b$
	$(d_{5/2}, g_{7/2})$	$[420]_{\frac{1}{2}}^{+}$	$c$	$d$
	$g_{9/2}$	$[404]_{\frac{9}{2}}^{+}$	$a'$	$b'$
	$h_{11/2}$	$[550]_{\frac{1}{2}}^{-}$	$f$	$e$
	$h_{11/2}$	$[541]_{\frac{3}{2}}^{-}$	$h$	$g$
Neutrons	$(s_{1/2}, d_{3/2})$	$[411]_{\frac{1}{2}}^{+}$	$A$	$B$
	$(d_{5/2}, g_{7/2})$	$[402]_{\frac{5}{2}}^{+}$	$C$	$D$
	$(d_{5/2}, g_{7/2})$	$[413]_{\frac{5}{2}}^{+}$	$A'$	$B'$
	$h_{11/2}$	$[523]_{\frac{7}{2}}^{-}$	$F$	$E$
	$h_{11/2}$	$[532]_{\frac{5}{2}}^{-}$	$H$	$G$

In the discussion of the configurations of the observed bands we distinguish between the low- and high-spin regions. At low and medium spins pairing correlations are important and comparison of experiment with theory is usually made within the framework of the CSM. Here, we adopt the results of previous work [12] for Bands 1 to 8 and make suggestions for the configuration assignments for the new bands. The quasiparticle levels relevant in this mass region are given in Table II where we adopt the standard notation introduced with the CSM [34]. The Coriolis interaction mixes the original shell model states in the rotating nucleus. Nevertheless, we include the shell model and Nilsson configurations for convenience.

For the higher-spin states, typically for states with  $I > 20$ , calculations have been performed using the configuration-dependent CNS formalism without pairing [19, 36]. The present calculations are similar to those presented for  $^{123}\text{Cs}$  in ref. [21]. Results of these calculations are compared with experimental excitation energies in Figs. 6 and 7 for the positive- and negative-parity bands, respectively.




FIG. 6: Experimental excitation energies relative to a rigid-rotor reference as a function of spin for the positive-parity bands in  $^{124}\text{Ba}$  (upper panel) compared to energies calculated within the framework of the CNS model (lower panel)

The CNS configurations are labeled by the number of particles in the different high- $j$  intruder orbitals outside a closed core. However, it should be noted that in this formalism no separation into core and valence particles is made and all orbitals up to  $N = 8$  are treated on an equal footing. In the case of  $^{124}\text{Ba}$  it is natural to choose  $^{114}\text{Sn}$  with  $Z = 50$  and  $N = 64$  as the core. The configurations may then be identified by the number of proton holes in the  $g_{9/2}$  subshell, the number of protons in  $h_{11/2}$  orbitals and the number of neutrons in  $h_{11/2}$  and  $i_{13/2}$  orbitals. The notation for labeling the configurations is given in the form  $[(p_0)p_1, n_0(n_1)]$ , where  $p_0$  is the number of proton holes in  $g_{9/2}$  orbitals (omitted when




FIG. 7: Experimental excitation energies relative to a rigid-rotor reference as a function of spin for negative-parity bands in  $^{124}\text{Ba}$  (upper panel) compared to energies calculated within the framework of the CNS model (lower panel)

$p_0 = 0$ ),  $p_1$  is the number of  $h_{11/2}$  protons,  $n_0$  is the number of neutrons in the  $h_{11/2}$  subshell and  $n_1$  the number of neutrons in  $i_{13/2}$  orbitals (omitted when  $n_1 = 0$ ). The energy of each configuration is minimized at each spin in the deformation space  $(\varepsilon_2, \varepsilon_4, \gamma)$ , which allows the development of collectivity to be treated as a function of spin.

In the CNS calculations we have used the parameters previously derived for the  $A = 130$  region [37]. As pairing is not included, the results should only be compared with experimental data at high spins where pairing is quenched. We note that there is a difference between




FIG. 8: Dynamic moment of inertia,  $\mathcal{J}^{(2)}$ , as a function of rotational frequency taken as  $\hbar\omega = E_\gamma/2$  for all observed bands.

the absolute values of the experimental and calculated level energies. The reason is that the experimental level energies are given relative to the ground-state energy while the reference of the calculated levels is the liquid drop energy at spin zero. Thus, the difference between the experimental and calculated energies depends essentially on the ground-state shell and pairing energy.

The experimental dynamic moments of inertia, aligned angular momenta and routhians for bands in  $^{124}\text{Ba}$  are shown in Figs. 8, 9 and 10, respectively. A rigid-rotor reference with the parameters  $\mathcal{J}_0 = 17 \text{ h}^2\text{MeV}^{-1}$  and  $\mathcal{J}_1 = 26 \text{ h}^4\text{MeV}^{-3}$  has been used [38]. The experimental crossing frequencies and the alignment gains are listed in Table III. As can be seen in Fig. 9, up to medium frequencies the aligned angular momenta,  $i_x$ , increase roughly in two discrete steps, one at low frequencies, around  $\hbar\omega = 0.37 \text{ MeV}$ , and another one at higher frequencies, around  $\hbar\omega = 0.45 \text{ MeV}$ . The alignment gains may be attributed to the decoupling and alignment of quasiparticles of  $h_{11/2}$  origin, either  $h_{11/2}$  quasiprotons or  $h_{11/2}$




FIG. 9: Aligned angular momentum,  $i_x$ , as a function of rotational frequency. A reference core with Harris parameters  $\mathcal{J}_0 = 17 \hbar^2 \text{MeV}^{-1}$  and  $\mathcal{J}_1 = 26 \hbar^4 \text{MeV}^{-3}$  has been subtracted.

quasineutrons. The smaller differences in alignment between the various bands are then caused by different quasiparticles with lower angular momentum,  $j$ , in the configurations. Furthermore, deformation changes and differences in pairing may influence alignments. At the highest observed frequencies some of the bands show an additional increase in alignment or other irregularities. In the following we will discuss the configurations for the various bands.

### A. Bands 1 and 2

Band 1 shows band crossings at rotational frequencies of 0.37 and 0.49 MeV/ $\hbar$  with alignment gains of 8.2 and 3.3  $\hbar$ , respectively, see Fig. 9. The first alignment has been interpreted as




FIG. 10: Experimental routhians,  $e'$ , as a function of rotational frequency. A reference core with Harris parameters  $\mathcal{J}_0 = 17 \hbar^2 \text{MeV}^{-1}$  and  $\mathcal{J}_1 = 26 \hbar^4 \text{MeV}^{-3}$  has been subtracted.

crossing of the ground band with the two-quasiproton  $h_{11/2}$  band, ef, and the second gain in alignment was suggested to result from a decoupling of an  $h_{11/2}$  neutron pair, EF [12]. Therefore, the configuration of Band 1 changes from the zero-quasiparticle ground-state band (gsb) to a two-quasiproton configuration, ef, and then to the two-quasiproton-two-quasineutron configuration, efEF.

The CNS calculations predict the  $[2,6]$  ( $\pi h_{11/2}^2 \otimes \nu h_{11/2}^6$ ) configurations to be yrast above spin 28 until it is crossed by the steeply downsloping  $[2,4]$  ( $\pi h_{11/2}^2 \otimes \nu h_{11/2}^4$ ) configuration, see Fig. 6. At lower spins, the CNS calculations predict the  $[(2)2,6]$  ( $\pi [(g_{9/2}^-)h_{11/2}^2] \otimes \nu h_{11/2}^6$ ) configuration below the  $[2,6]$  structure. However, the calculated relative positions of the  $[2,6]$  and  $[(2)2,6]$  configurations can depend on the Nilsson parameters and there is some uncertainty in their relative positions. Indeed, in the neighboring  $^{123}\text{Cs}$  nucleus these calculations also predicted the  $[(2)2,6]$  below the  $[2,6]$  configuration in that spin region, while the ob-

TABLE III: Experimental crossing frequencies and aligned angular momenta in  $^{124}\text{Ba}$ .

Band	Before crossing Config.	$i_x$ [ $\hbar$ ]	$\hbar\omega_c$ [MeV]	$\Delta i_x$ [ $\hbar$ ]	Config. above 1 <sup>st</sup> crossing	$\hbar\omega_c$ [MeV]	$\Delta i_x$ [ $\hbar$ ]	Config. above 2 <sup>nd</sup> crossing	$\hbar\omega_c$ [MeV]	$\Delta i_x$ [ $\hbar$ ]
1	0-qp	0	0.37	8.2	ef	0.49	3.3	efEF	-	-
2	0-qp	0	0.41	6.2	EF	0.44	4.4	EFef	0.63	2.5
3	eb	$\approx 5.1$	0.46	6.0	ebEF	-	-	-	-	-
4	ea	5.4	0.44	4.0	eaGH	0.59	$\approx 6.0$	eaGHEF	-	-
5	eb'	5.8	0.44	$> 6.1$	eb'EF	-	-	-	-	-
6	ea'	5.8	0.44	$> 5.8$	ea'EF	-	-	-	-	-
9	efGH	11	-	-	-	-	-	-	-	-
10	efFH	9.5	-	-	-	-	-	-	-	-
12	eb'EA'	8.5	0.52	5.2	eb'EA'GH	-	-	-	-	-
13	eb'FA'	8.5	0.52	4.9	eb'FA'GH	0.64	$> 2.5$	-	-	-

served alignment frequency favors the [2,6] assignment [21]. The difference may also be due to remnants of pairing at high spins. If pairing is not negligible, [(2)2,6] is a six-quasiparticle configuration while [2,6] is a four-quasiparticle configuration. The extra energy necessary to break a pair in order to form a six-quasiparticle excitation may lift the [(2)2,6] configuration in reality to higher energy than predicted by the CNS calculations which neglect pairing. Thus, Band 1 probably corresponds to the [2,6] configuration in the spin region between 22 and 34, in agreement with the previous assignments [12]. The calculated shape parameters for this configuration are  $\varepsilon_2 \approx 0.24$  and  $\gamma \approx 0^\circ$ .

Above the  $I^\pi = 34^+$  state, Band 1 becomes irregular and forks into three branches, see Fig. 1. This is the typical behavior expected for band termination. Terminating bands have first been observed in Dy-Er region [39]. However, there exist analogies between the  $A = 160$  and 125 mass regions [20]. As illustrated in Fig. 11, four protons in  $h_{11/2}$  orbitals outside the  $Z = 64$  semi-closed core in  $^{156}\text{Er}$  may be compared to four neutrons in  $h_{11/2}$  orbitals in  $^{124}\text{Ba}$  which has  $N = 68$  neutrons. Similarly, six neutrons above the  $N = 82$  closed core in  $^{156}\text{Er}$  is analogous to six protons outside the  $Z = 50$  closed core in  $^{124}\text{Ba}$ . Thus, the observed fully aligned configuration  $\pi(h_{11/2})_{16}^4 \otimes \nu[(h_{9/2})_8^2(f_{7/2})_6^2(i_{13/2})_{12}^2]$  of the  $I^\pi = 42^+$  state in  $^{156}\text{Er}$  [40]



is analogous to the configuration  $\pi[(h_{11/2})_{10}^2(g_{7/2}d_{5/2})_{10}^4] \otimes \nu(h_{11/2})_{16}^4$  with  $I^\pi = 36^+$  in  $^{124}\text{Ba}$ .




FIG. 11: Analogy between the  $A = 125$  and  $160$  mass regions: the filling of subshells outside the  $^{114}\text{Sn}$  and  $^{146}\text{Gd}$  cores is illustrated schematically.

The present CNS calculations agree well with the expectations based on this analogy. In Fig. 6 we have differentiated between three types of states of the  $[2,4]$  configuration defined above. The configuration  $[2,4]a$  denotes the states with only four valence neutrons relative to the  $N = 64$  core and they all are located in the  $h_{11/2}$  subshell, while  $[2,4]b$  and  $[2,4]c$  correspond to the states where one neutron and two neutrons, respectively, are excited from the  $d_{5/2}g_{7/2}$  subshell to the  $s_{1/2}d_{3/2}$  orbitals. The  $[2,4]a$  configuration is predicted to become yrast around spin 34 and to terminate at  $I^\pi = 36^+$  in a strongly favored state. Two favored  $38^+$  states and two favored  $40^+$  states are predicted corresponding to the  $[2,4]b$  and  $[2,4]c$  configurations. Experimentally, we observe two  $36^+$  states and three  $38^+$  states. Comparing the relative positions of the calculated states with different configurations with the relative positions of the observed bands shown in Fig. 6, it is most likely that the  $[2,4]a$  ( $\pi[(h_{11/2})_{10}^2(g_{7/2}d_{5/2})_{10}^4] \otimes \nu(h_{11/2})_{16}^4$ ) terminating configuration has to be assigned to the yrast  $36^+$  state. This is a non-collective oblate state, characterized by the deformation parameters  $\varepsilon_2 \approx 0.2$  and  $\gamma = 60^\circ$ . Its total angular momentum is built solely from the maximally aligned angular momenta of the valence particles corresponding to the  $[2,4]a$  configuration. Comparing the relative positions of the higher-energy two nearly degenerate  $36^+$  states, they probably correspond to the  $[2,6]$  and  $[2,4]b$  configurations, respectively. Similarly, the lowest two strongly favored  $38^+$  states probably correspond to the  $[2,4]b$  and  $[2,4]c$  configurations, respectively, while the highest-energy  $38^+$  state has the  $[2,6]$  configuration. The  $38^+$  states belonging to  $[2,4]b$  and  $[2,4]c$  are partially aligned and these configurations are expected to terminate at spins 44 and 40, respectively. We note here that the predicted  $[2,4]$  structure

in  $^{124}\text{Ba}$  is very similar to the predicted [1,4] structure in  $^{123}\text{Cs}$  [21], but with an additional  $h_{11/2}$  proton. Therefore, the  $36^+$  state is analogous to the  $63/2^-$  terminating state in  $^{123}\text{Cs}$ . As shown in ref. [21], the states around this spin in  $^{123}\text{Cs}$  are due to energy minima in the potential energy surface corresponding to zero, one and two particle-hole excitations from neutron ( $d_{5/2}g_{7/2}$ ) to ( $s_{1/2}d_{3/2}$ ) orbitals. Calculations for  $^{124}\text{Ba}$  also show the existence of these minima around spin 36.

Band 2 starts with an alignment of about  $6.2 \hbar$ , see Fig. 9, and shows a band crossing with an alignment gain of  $4.4 \hbar$  at a frequency of  $0.44 \text{ MeV}/\hbar$ . Previously, the two-quasineutron configuration EF, of  $\nu h_{11/2}$  origin, was assigned to this band below  $0.44 \text{ MeV}/\hbar$ . At this frequency the decoupling of an  $h_{11/2}$  quasiproton pair takes place and the configuration changes to the two-quasineutron-two-quasiproton ( $\nu h_{11/2}^2 \otimes \pi h_{11/2}^2$ ) structure, EFef [12]. In this frequency range, Band 1 has the same configuration, however in Band 2 the  $h_{11/2}$  quasineutron pair is almost fully aligned ( $\approx 6.2 \hbar$ ) and the  $h_{11/2}$  quasiproton pair is only partially aligned ( $4.4 \hbar$ ), whereas in Band 1 it is opposite: the  $h_{11/2}$  quasiproton pair is nearly fully aligned ( $8.2 \hbar$ ) and the  $h_{11/2}$  quasineutron pair shows a smaller alignment ( $3.3 \hbar$ ). The close similarity of the configurations of Bands 1 and 2 results in the strong decay from the higher-lying Band 2 into the yrast Band 1.

In the calculations using the CNS formalism the configurations above the first two band crossings in Bands 1 and 2 are the same, i.e. both are represented by a pair of aligned  $h_{11/2}$  protons and  $h_{11/2}$  neutrons, however with a different distribution of the valence protons and neutrons in low- $j$  orbitals. Indeed, a second minimum appears in the calculations with shape parameters  $\varepsilon_2 \approx 0.24$  and  $\gamma \approx -30^\circ$ . The corresponding configuration is labeled [2,6]\* in Fig. 6. This minimum becomes more and more shallow with increasing spin and disappears above spin 34. However, the calculated relative energy difference between the first and second minimum does not agree well with that observed for Bands 1 and 2. At spin 22, both the calculated and experimental energy differences are  $\approx 250 \text{ keV}$ . This difference is predicted to increase with increasing spin up to  $\approx 700 \text{ keV}$ , while experimentally it is constant up to  $I = 28$  and continuously decreases above this spin to about  $30 \text{ keV}$ . This behavior indicates a change in the configuration around spin 28. According to Fig. 6, a possible candidate for the configuration above this spin can be [2,4]b which is predicted to cross the [2,6] configuration at spin 36.

The alignment of Band 2, see Fig. 9, shows an irregularity at  $\hbar\omega = 0.63 \text{ MeV}$  with an alignment gain of  $2.5 \hbar$ , which is not seen in Band 1. In the yrast bands of the neighboring even-even nuclei  $^{126}\text{Ba}$  [13] and  $^{128}\text{Ba}$  [15] a similar behavior with alignment gains of  $\approx 2$  and  $\approx 4 \hbar$ , respectively, has been observed. The alignment of a pair of protons of  $g_{7/2}$  origin was suggested for the crossing around  $\hbar\omega = 0.6 \text{ MeV}$  in  $^{126}\text{Ba}$  [13], whereas an alignment of

a second pair of  $h_{11/2}$  neutrons was proposed for the crossing at this frequency in  $^{128}\text{Ba}$  [15]. These may be alternative explanations for the behavior of Band 2 at high spins in  $^{124}\text{Ba}$ .

## B. Coupled Bands

Information about the quasiparticle configurations of the coupled bands can be extracted from the ratios of reduced magnetic dipole and electric quadrupole transition probabilities,  $B(M1; I \rightarrow I - 1)/B(E2; I \rightarrow I - 2)$ . The experimental ratios have been obtained from the  $\gamma$ -ray-intensity ratios using

$$\frac{B(M1; I \rightarrow I - 1)}{B(E2; I \rightarrow I - 2)} = 0.697 \frac{I_\gamma(M1; I \rightarrow I - 1)}{I_\gamma(E2; I \rightarrow I - 2)} \frac{E_\gamma^5(E2)}{E_\gamma^3(M1)} \frac{1}{1 + \delta^2} \left(\frac{\mu_N}{eb}\right)^2$$

where  $E_\gamma$  and  $I_\gamma$  represent the energy in MeV and the intensity of the  $\gamma$ -ray transitions, respectively. The mixing ratio,  $\delta$ , for the  $\Delta I = 1$  transitions is small and  $\delta^2$  has been set to zero. The experimentally deduced  $B(M1)/B(E2)$  ratios are compared with the theoretical values obtained using the following generalized expression for multi-quasiparticle configurations, formulated in [4, 41] and based on the geometrical model of Dönau and Frauendorf [42, 43]:

$$\begin{aligned} \frac{B(M1; I \rightarrow I - 1)}{B(E2; I \rightarrow I - 2)} &= \frac{12}{5Q_o^2 \cos^2(\gamma + 30^\circ)} \left[ 1 - \frac{K_{tot}^2}{\left(I - \frac{1}{2}\right)^2} \right]^{-2} \\ &\times \left\{ \left( I - \frac{K_{tot}^2}{I^2} \right)^{1/2} \left[ K_1(g_1 - g_R) \left( 1 \pm \frac{\Delta e'}{\hbar\omega} \right) + \sum_\lambda K_\lambda(g_\lambda - g_R) \right] \right. \\ &\quad \left. - \frac{K_{tot}}{I} \left[ (g_1 - g_R)i_1 + \sum_\lambda (g_\lambda - g_R)i_\lambda \right] \right\}^2. \end{aligned}$$

$K_\lambda$ ,  $g_\lambda$  and  $i_\lambda$  stand for the  $K$ -value, gyromagnetic factor and aligned angular momentum, respectively, of the rotation- or Fermi-aligned quasiparticles involved in the configuration.  $K_1$ ,  $g_1$  and  $i_1$  refer to the strongly coupled particle. The approximation  $g_R = Z/A$  was used for the rotational  $g$  factor.  $K_{tot} = K_1 + \sum_\lambda K_\lambda$  denotes the total  $K$ -value of the configuration. The values for the intrinsic  $g$  factors for the different orbitals,  $g_\lambda$ , as well as the quadrupole moment,  $Q_o = 3.9 b$ , have been taken from [13]. The shape parameter  $\gamma$  was set to zero. The aligned angular momenta,  $i_m$ , were determined from the alignment plots, see Fig. 9. The signature splitting,  $\Delta e'$ , was extracted from the routhian plots, see Fig. 10. The parameters used in the calculations are listed in Table IV.

1. Bands 3, 4 and 14

Bands 3 and 4 appear to be a pair of signature-partner bands with a large splitting. It was suggested [12] that Band 3 (the favored signature) corresponds to the quasiproton eb configuration and Band 4 (the unfavored signature) to ea or ec. The  $\pi h_{11/2}g_{7/2}$  and  $\pi(h_{11/2}d_{5/2}) \otimes \nu h_{11/2}^4$  configurations were also assigned to similar bands in the neighboring even-even nuclei  $^{126}\text{Ba}$  [13] and  $^{128}\text{Ba}$  [15], respectively. It was suggested [32, 33, 44] that those bands have a two-quasiproton structure mixed with the octupole vibrational band which accounts for the large signature splitting at low spins. Both bands are crossed by the configuration with an additional  $h_{11/2}$  quasineutron pair, EF.

TABLE IV: Gyromagnetic factors and alignments used for the calculation of  $B(M1)/B(E2)$  ratios.

Bands	Subshell	Nilsson orbital	$g$ -factor	K	$i[\hbar]$
3,4	$\pi h_{11/2}$	$[550]_{\frac{1}{2}}^{-}$	1.17	0.5	5.0
	$\pi g_{7/2}$	$[422]_{\frac{3}{2}}^{+}$	0.72	1.5	1.0
	$\pi d_{5/2}$	$[420]_{\frac{1}{2}}^{+}$	1.38	0.5	1.0
5,6	$\pi h_{11/2}$	$[550]_{\frac{1}{2}}^{-}$	1.17	0.5	5.0
	$\pi g_{9/2}$	$[404]_{\frac{9}{2}}^{+}$	1.27	4.5	1.0
12,13	$\pi h_{11/2}$	$[550]_{\frac{1}{2}}^{-}$	1.17	0.5	5.0
	$\pi d_{5/2}$	$[420]_{\frac{1}{2}}^{+}$	1.38	0.5	1.0
	$\pi g_{9/2}$	$[404]_{\frac{9}{2}}^{+}$	1.27	4.5	0.0
	$\nu h_{11/2}$	$[532]_{\frac{5}{2}}^{-}$	-0.21	2.5	3.5
		$[541]_{\frac{3}{2}}^{-}$	-0.21	1.5	3.5
	$\nu g_{7/2}$	$[402]_{\frac{5}{2}}^{+}$	0.21	-	0.5
		$[404]_{\frac{7}{2}}^{+}$	0.21	3.5	0.5
$\nu d_{5/2}$	$[413]_{\frac{5}{2}}^{+}$	-0.33	2.5	0.5	

The  $B(M1)/B(E2)$  ratios determined from our data for Bands 3 and 4 are compared with the calculated values for the configurations  $\pi h_{11/2}g_{7/2}$  and  $\pi h_{11/2}d_{5/2}$  in Fig. 12. The calculated ratios for the configuration  $\pi h_{11/2}d_{5/2}$  agree better with the experimental data than those

for the  $\pi h_{11/2}g_{7/2}$  configuration. Therefore, we have assigned the configuration  $\pi h_{11/2}d_{5/2}$  to Bands 3 and 4 in the low-spin region. At higher spins these configurations are expected to become more and more mixed.




FIG. 12:  $B(M1)/B(E2)$  ratios derived from measured  $\gamma$ -ray branching ratios for Bands 3 and 4 and calculated values for the  $\pi h_{11/2}d_{5/2}$  and  $\pi h_{11/2}g_{7/2}$  configurations.

Both bands show a band crossing at a rotational frequency of  $\approx 0.45$  MeV/ $\hbar$  with a gain in alignment of 6 and 4  $\hbar$ , respectively. The rather large alignment suggests that aligned  $h_{11/2}$  neutrons or protons are present in the configuration above that frequency. We suggest the configurations ebEF and eaGH for Bands 3 and 4, respectively, in the frequency range around 0.5 MeV/ $\hbar$ . At higher frequencies, Band 3 seems to lose alignment, see Fig. 9. However, this might be caused by a change to a reference core with a smaller deformation. A reduction of the Harris parameters to  $\mathcal{J}_0 = 13 \hbar^2 \text{MeV}^{-1}$  and  $\mathcal{J}_1 = 21 \hbar^4 \text{MeV}^{-3}$  makes the alignment of Band 3 constant.

Band 4 shows a gain in alignment of  $\approx 6 \hbar$  at a frequency of 0.59 MeV/ $\hbar$ , see Fig. 9. Such a large gain can only be caused by  $h_{11/2}$  neutrons or protons. As Band 3 does not show a similar increase at that frequency, the corresponding alignment has to be blocked. Thus, it is most likely that it is caused by a pair of  $h_{11/2}$  neutrons, e.g. EF, which are already aligned in Band 3 at this frequency. At the highest observed frequencies, the configuration

eaGHEF is therefore suggested for Band 4.

At high-spins the [1,6] configuration is suggested by the CNS calculations. As seen in Fig. 7, there is good agreement for the relative positions and slopes between the calculated [1,6] bands and the observed bands 3 and 4 up to spin 26.

Band 14 decays into Band 3 around spin 30. Unfortunately, its spin assignment is not certain. If our tentative assignment is correct, Band 14 has an alignment which is  $\approx 5 \hbar$  higher than that of Band 3, which could be explained by a six-quasiparticle configuration.

## 2. Bands 5 and 6

Bands 5 and 6 form a pair of signature partners with negligible signature splitting. The two-quasiproton configurations,  $eb'$  and  $ea'$ , of  $g_{9/2}h_{11/2}$  origin, seem to be a natural choice for these bands [12]. At a frequency of  $0.44 \text{ MeV}/\hbar$  a band crossing with an alignment gain of  $> 6.1$  and  $> 5.8 \hbar$  is observed for Bands 5 and 6, respectively. The high-spin part of the bands probably contains a pair of decoupled  $h_{11/2}$  neutrons, resulting in the configurations  $eb'EF$  and  $ea'EF$ , respectively.




FIG. 13:  $B(M1)/B(E2)$  ratios derived from measured  $\gamma$ -ray branching ratios for Bands 5 and 6 and calculation for the  $\pi g_{9/2}h_{11/2}$  configuration.

$B(M1)/B(E2)$  ratios have been determined and are compared with calculated ratios for the  $\pi g_{9/2}h_{11/2}$  configuration in Fig. 13. At lower spins a rather large  $B(M1)/B(E2)$  ratio has

been obtained. In this region mixing occurs with other states of the same spin and parity which also accounts for the observed decay-out transitions. Large  $B(M1)/B(E2)$  ratios are calculated for configurations involving neutrons, e.g. for the  $\nu h_{11/2}g_{7/2}$  or  $\nu h_{11/2}s_{1/2}$  configurations. Mixing with such states may explain the observed increase of the  $B(M1)/B(E2)$  ratios near the bottom of Bands 5 and 6. The good agreement at medium spins supports the previous configuration assignment [12] which corresponds to the [(1)1,6] CNS configuration. The theoretically predicted relative positions and slopes of this configuration are in agreement with those of Bands 5 and 6, as seen in Fig. 7.

### 3. Bands 7 and 8

Bands 7 and 8 have been interpreted previously to be built on a  $\gamma$  vibration [18, 45]. The present results do not add significant information to these bands.

### 4. Bands 12 and 13

The new Bands 12 and 13 form a pair of coupled bands without signature splitting. Their alignment is rather large,  $i_x \geq 8 \hbar$ , already at low rotational frequencies, suggesting a multi-quasiparticle configuration. The experimental  $B(M1)/B(E2)$  ratios determined from our data are compared with calculations for three different configurations,  $\pi(h_{11/2}d_{5/2}) \otimes \nu(h_{11/2}g_{7/2})$ ,  $\pi(g_{9/2}h_{11/2}) \otimes \nu(h_{11/2}g_{7/2})$  and  $\pi(g_{9/2}h_{11/2}) \otimes \nu(h_{11/2}d_{5/2})$ , in Fig. 14. As can be seen, the calculated ratios for the configurations  $\pi(g_{9/2}h_{11/2}) \otimes \nu(h_{11/2}g_{7/2})$  and  $\pi(g_{9/2}h_{11/2}) \otimes \nu(h_{11/2}d_{5/2})$  are close to the experimental data. These configurations are expected to be mixed and it is not possible distinguish between them. Thus, we assign the configurations eb'EA' and eb'FA' to Bands 12 and 13, respectively.

At a frequency of  $\omega = 0.5 \text{ MeV}/\hbar$  a further band crossing is observed. The alignment gain of about  $5 \hbar$  may be due to a pair of  $h_{11/2}$  neutrons. As orbitals E and F are already occupied, we suggest that the alignment is due to the GH quasineutron pair.

Two band structures with small signature splitting, the [1,5] and [(1)2,6] configurations, are predicted by the present CNS calculations. They approach the yrast [2,6] configuration in the spin 20 - 35 region, see Fig. 6. The experimental energy difference between the [1,5] and [2,6] configurations is predicted to decrease up to about spin 32 and then to increase at higher spins. For the [(1)2,6] configuration this energy difference is predicted to increase continuously in that spin region. The energy difference between Band 1 and Bands 12 and 13 decreases above spin 24 and becomes very small in the spin 30 - 35 region. This observation




FIG. 14:  $B(M1)/B(E2)$  ratios derived from measured  $\gamma$ -ray branching ratios for Bands 12 and 13 and calculated ratios for three different configurations.

favors the [1,5] assignment over the [(1)2,6] configuration. Another argument against the [(1)2,6] configuration is that, similar to the case of the [(2)2,6] configuration discussed above, it may in reality lie at higher energy than predicted due to the pairing interaction which is not taken into account in the CNS calculations.

### C. Bands 9 and 10

The new Bands 9 and 10 are observed above spins 17 and 18, respectively. They both decay into Band 1, suggesting a similarity in structure. They start with a high alignment which, in the lower part, increases by about  $3\hbar$  and then stays at 11 and  $9.5\hbar$ , respectively, see Fig. 9. At higher spins their alignment pattern looks different. While the alignment of Band 9 decreases, Band 10 continuously gains alignment up to the highest-spin states observed. The rather large alignment suggests that both bands are four-quasiparticle configurations, probably containing a pair of aligned protons coupled to another aligned proton or a neutron pair. We tentatively assign the quasiparticle configurations efGH and efFH to Bands 9 and



10, respectively. The difference in the appearance of the alignment might be caused by differences in the reference core. A reduction of the Harris parameters for Band 9 to  $\mathcal{J}_0 = 13 \hbar^2 \text{MeV}^{-1}$  and  $\mathcal{J}_1 = 21 \hbar^4 \text{MeV}^{-3}$  makes its alignment constant.

Inspection of the results of the CNS calculations presented in Fig. 6 shows that the configuration labeled [1,5] could reproduce the observed properties reasonably well. However, it seems unlikely that Bands 9 and 10 are a pair of signature partners since no inter-band transitions are observed. Furthermore, we have already tentatively assigned this configuration to Bands 12 and 13, as discussed above. Thus, there are no good candidates predicted by the present CNS calculations for the configurations of these bands.

#### D. Band 11

Band 11 is not firmly connected to known levels in  $^{124}\text{Ba}$ . However, the systematics of excitation energies and spins of the bands observed in this nucleus allow an approximate placement in the level scheme and limit the spin range. As the band decays to high-spin states of Band 1, their structures should have some similarity.

Band 11 is observed only above spin 25 or 26 and continues without band crossing up to spin 45 or 46. The most likely candidate from the CNS calculations is the [2,5(1)] configuration, i.e.  $\pi h_{11/2}^2 \otimes \nu h_{11/2}^5 i_{13/2}$ . In its decay to Band 1 it goes over to the [2, 6] configuration with two  $h_{11/2}$  protons and six  $h_{11/2}$  neutrons.

Band 11 shows a gradual decrease in the dynamic moment of inertia in a manner similar to that of several high-spin bands in this mass region. In Fig. 15 the dynamic moments of inertia of Band 11 and similar bands observed in  $^{132}\text{Ce}$  [46–48] and  $^{126}\text{Xe}$  [49] are compared. The  $^{132}\text{Ce}$  band is highly deformed; a transition quadrupole moment of  $Q = 8.8 \text{ b}$  has been measured [46]. In  $^{125,126}\text{Xe}$  several bands, which extend up to very high spins, have been observed recently [49]. However, only for the  $^{126}\text{Xe}$  band shown in Fig. 15 was the quadrupole moment estimated,  $Q \sim 5.2 \text{ b}$  [49]. The dynamic moment of inertia of Band 11 lies between those of  $^{126}\text{Xe}$  and  $^{132}\text{Ce}$  shown in Fig. 15. This systematics suggests that it probably has a larger deformation than the other bands in  $^{124}\text{Ba}$ . The excitation of an  $i_{13/2}$  neutron may cause the larger deformation, however lifetime measurements are needed for a final proof. CSM calculations for  $^{132}\text{Ce}$  show that the frequency dependence of the dynamic moment of inertia is a combined effect of gradual alignment of protons in  $h_{11/2}$  and neutrons in  $h_{9/2}$  and  $h_{11/2}$  orbitals [50], and a similar explanation may apply also to Band 11 in  $^{124}\text{Ba}$ .




FIG. 15: Dynamic moment of inertia,  $\mathcal{J}^{(2)}$ , for Band 11 in  $^{124}\text{Ba}$  as well as for similar bands in  $^{132}\text{Ce}$  [48] and  $^{126}\text{Xe}$  [49], as a function of rotational frequency.

## V. SUMMARY

In summary, high-spin states in  $^{124}\text{Ba}$  have been investigated in two experiments using the Euroball and Gammasphere spectrometers, respectively. Previously known bands have been extended and six new bands have been found. Configurations have been assigned to the band structures, based on comparisons with predictions of the CSM in the lower-spin region and with the CNS model at high spins. The positive-parity yrast band, Band 1, shows the typical behavior of band termination at the highest spins. Configurations for the terminating states which compete with the collective excitations are suggested on the basis of CNS calculations. One of the new bands, Band 11, extends to spins around 46 without band crossings. Its dynamic moment of inertia decreases steadily, similar to those of large-deformed bands in this mass region.

## VI. ACKNOWLEDGMENTS

This work was supported in part by the EU (contract EUROVIV: HPRI-CT-1999-00078), the German BMBF (contract 06 BN 907), the Hungarian Scientific Research Fund OTKA (contract numbers T046901 and T038404), the Bolyai János Foundation of HAS, and by INFN, Italy. The authors are grateful to I. Ragnarsson for helpful discussions of the results of the CSN calculations and the configuration assignments to the band structures in  $^{124}\text{Ba}$ .

- 
- [1] C. M. Petrache, G. LoBianco, P. G. Bizzeti, A. M. Bizzeti-Sona, D. Bazzacco, S. Lunardi, M. Nespolo, G. de Angelis, D. R. Napoli, N. Blasi, et al., *Eur. Phys. J.* **A 16**, 337 (2003).
  - [2] J. F. Smith, V. Medina-Chico, C. J. Chiara, M. P. Carpenter, C. N. Davids, M. Devlin, J. L. Durell, D. B. Fossan, S. J. Freeman, R. V. F. Janssens, et al., *Phys. Rev.* **C 69**, 034339 (2004).
  - [3] F. Seiffert, W. Lieberz, A. Dewald, S. Freund, A. Gelberg, A. Granderrath, D. Lieberz, R. Wirowski, and P. von Brentano, *Nucl. Phys.* **A 554**, 287 (1993).
  - [4] S. Törmänen, S. Juutinen, R. Julin, B. Cederwall, A. Johnson, R. Wyss, P. Ahonen, B. Fant, M. Matsuzaki, J. Nyberg, et al., *Nucl. Phys.* **A 572**, 417 (1994).
  - [5] H. Timmers, J. Simpson, M. A. Riley, T. Bengtsson, M. A. Bentley, F. Hanna, S. M. Mullins, J. F. Sharpey-Schafer, and R. Wyss, *J. Phys. G: Nucl. Part. Phys.* **A 20**, 287 (1994).
  - [6] J. Timár, J. Simpson, E. S. Paul, S. Araddad, C. W. Beausang, M. A. Bentley, M. J. Joyce, and J. F. Sharpey-Schafer, *J. Phys. G: Nucl. Part. Phys.* **21**, 783 (1995).
  - [7] I. Wiedenhöver, J. Yan, U. Neuneyer, R. Wirowski, P. von Brentano, A. Gelberg, N. Yoshida, and T. Otsuka, *Nucl. Phys.* **A 582**, 77 (1995).
  - [8] M. Serris, C. T. Papadopoulos, R. Vlastou, C. A. Kalfas, S. Kossionides, N. Fotiades, S. Harissopoulos, C. W. Beausang, M. J. Joyce, E. S. Paul, et al., *Z. Phys.* **A 358**, 37 (1997).
  - [9] I. Schneider, R. S. Chakrawarthy, I. Wiedenhöver, A. Schmidt, H. Meise, P. Petkov, A. Dewald, P. von Brentano, O. Stuch, K. Jessen, et al., *Phys. Rev.* **C 60**, 014312 (1999).
  - [10] V. Werner, H. Meise, I. Wiedenhöver, A. Gade, and P. von Brentano, *Nucl. Phys.* **A 692**, 451 (2001).
  - [11] R. Wyss, A. Granderrath, R. Bengtsson, P. von Brentano, A. Dewald, A. Gelberg, A. Gizon, J. Gizon, S. Harissopoulos, A. Johnson, et al., *Nucl. Phys.* **A 505**, 337 (1989).

- [12] S. Pilotte, S. Flibotte, S. Monaro, N. Nadon, N. Prevost, P. Taras, H. R. Andrews, D. Horn, V. P. Janzen, D. C. Radford, et al., Nucl. Phys. **A 514**, 545 (1990).
- [13] D. Ward, V. P. Janzen, H. R. Andrews, D. C. Radford, G. C. Ball, D. Horn, J. C. Waddington, J. K. Johansson, F. Banville, J. Gascon, et al., Nucl. Phys. **A 529**, 315 (1991).
- [14] A. Granderath, P. F. Mantica, R. Bengtsson, R. Wyss, P. von Brentano, A. Gelberg, and F. Seiffert, Nucl. Phys. **A 597**, 427 (1996).
- [15] O. Vogel, R. S. Chakrawarthy, A. Dewald, P. Petkov, K. Jessen, J. Gableske, P. von Brentano, D. Bazzacco, A. Gizon, J. Gizon, et al., Eur. Phys. J. **A 4**, 323 (1999).
- [16] J. F. Smith, C. J. Chiara, D. B. Fossan, G. J. Lane, J. M. Sears, I. Thorslund, I. M. Hibbert, R. Wadsworth, I. Y. Lee, and A. O. Macchiavelli, Phys. Lett. **B 483**, 7 (2000).
- [17] C. M. Petrache, G. LoBianco, D. Bazzacco, T. Kröll, S. Lunardi, R. Menegazzo, M. Nespolo, P. Pavan, C. R. Alvarez, G. de Angelis, et al., Eur. Phys. J. **A 12**, 135 (2001).
- [18] J. P. Martin, V. Barci, H. El-Samman, A. Gizon, J. Gizon, W. Klamra, B. M. Nyakó, F. A. Beck, T. Byrski, and J. C. Merdinger, Nucl. Phys. **A 489**, 169 (1988).
- [19] A. V. Afanasjev, D. B. Fossan, G. J. Lane, and I. Ragnarsson, Phys. Rep. **322**, 1 (1999).
- [20] I. Ragnarsson, Z. Xing, T. Bengtsson, and M. A. Riley, Phys. Scripta **Vol. 34**, 651 (1986).
- [21] A. K. Singh, H. Hübel, J. Domscheit, G. B. Hagemann, B. Herskind, D. R. Jensen, J. N. Wilson, R. Clark, M. Cromaz, P. Fallon, et al., Phys. Rev. **C 70**, 034315 (2004).
- [22] A. Al-Khatib, A. K. Singh, H. Hübel, P. Bringel, A. Bürger, A. Neußer-Neffgen, G. Schönwaßer, G. Hagemann, C. R. Hansen, B. Herskind, et al., Acta Phys. Pol. **B 36**, 1029 (2005).
- [23] H. Hübel, Acta Phys. Pol. **B 36**, 1015 (2005).
- [24] I. Y. Lee, Nucl. Phys. **A 520**, 641c (1990).
- [25] F. A. Beck, Prog.Part.Nucl.Phys. **28**, 443 (1992).
- [26] J. Simpson, Z. Phys. **A 358**, 139 (1997).
- [27] J. N. Scheurer, M. Aiche, M. M. Aleonard, G. Barreau, F. Bourguine, D. Boivin, D. Cabaussel, J. F. Chemin, T. P. Doan, J. P. Goudour, et al., Nucl. Instr. Meth. A **385**, 501 (1997).
- [28] J. Gál, G. Hegyesi, J. Molnár, B. M. Nyakó, G. Kalinka, J. N. Scheurer, M. M. Aléonard, J. F. Chemin, J. L. Pedroza, K. Juhász, et al., Nucl. Instr. Meth. A **516**, 502 (2004).
- [29] D. C. Radford, Nucl. Instr. Meth. A **361**, 297 (1995).
- [30] O. Klein and Y. Nishina, Z. Phys. **52**, 853 (1929).

- [31] D. Roßbach, A. Gørgen, H. Hübeler, E. Mergel, G. Schönwaßer, A. N. Wilson, F. Azaiez, A. Astier, D. Bazzacco, M. Bergström, et al., Phys. Lett. **B 513**, 9 (2001).
- [32] P. Mason, G. Benzoni, A. Bracco, F. Camera, B. Million, O. Wieland, S. Leoni, A. K. Singh, A. Al-Khatib, H. Hübeler, et al., Phys. Rev. **C 72**, 064315 (2005).
- [33] R. Piepenbring and J. Leandri, Phys. Lett. **B 267**, 17 (1991).
- [34] R. Bengtsson and S. Frauendorf, Nucl. Phys. **A 314**, 27 (1979).
- [35] R. Bengtsson, S. Frauendorf, and F. R. May, At. Nucl. Data Tables **35**, 15 (1986).
- [36] T. Bengtsson and I. Ragnarsson, Nucl. Phys. **A 436**, 14 (1985).
- [37] A. V. Afanasjev and I. Ragnarsson, Nucl. Phys. **A 608**, 176 (1996).
- [38] D. M. Todd, R. Aryaeinejad, D. J. G. Love, A. H. Nelson, P. J. Nolan, P. J. Smith, , and P. J. Twin, J. Phys. G: Nucl. Part. Phys. **10**, 1407 (1984).
- [39] A. V. Afanasjev and I. Ragnarsson, Nucl. Phys. **A 591**, 387 (1995).
- [40] F. S. Stephens, M. A. Delplanque, R. M. Diamond, A. O. Macchiavelli, and J. E. Draper, Phys. Rev. Lett. **54**, 2584 (1985).
- [41] D. C. Radford, H. R. Andrews, G. C. Ball, D. Horn, D. Ward, F. Banville, S. Flibotte, S. Monaro, S. Pilotte, P. Taras, et al., Nucl. Phys. **A 545**, 665 (1992).
- [42] F. Dönaue and S. Frauendorf, Proc. Conf. on High Angular Momentum Properties of Nuclei (Oak Ridge, 1982) ed. N.R. Johnson (New York: Harwood Academic) p. 143 (1983).
- [43] F. Dönaue, Nucl. Phys. **A 471**, 469 (1987).
- [44] R. Wyss, A. Johnson, D. J. G. Love, M. J. Godfrey, and S. M. Mullins, Z. Phys. **A 332**, 241 (1989).
- [45] T. Komatsubara, T. Hosoda, H. Sakamoto, T. Aoki, and K. Furuno, Nucl. Phys. **A 496**, 605 (1989).
- [46] A. J. Kirwan, G. C. Ball, P. J. Bishop, M. J. Godfrey, P. J. Nolan, D. J. Thornley, D. J. G. Love, and A. H. Nelson, Phys. Rev. Lett. **58**, 467 (1987).
- [47] P. J. Nolan, A. Kirwan, D. J. G. Love, A. H. Nelson, D. J. Unwin, and P. J. Twin., J. Phys. G: Nucl. Part. Phys. **11**, L17 (1985).
- [48] E. S. Paul, P. T. W. Choy, C. Andreoiu, A. J. Boston, A. O. Evans, C. Fox, S. Gros, P. J. Nolan, G. Rainovski, J. A. Sampson, et al., Phys. Rev. **C 71**, 054309 (2005).
- [49] C. R. Hansen, G. B. Hagemann, B. Herskind, D. R. Jensen, G. Sletten, J. N. Wilson, S. Ødegård, P. Bringel, C. Engelhardt, H. Hübeler, et al., AIP Conf. Proc. **764**, 46 (2005).

- [50] R. Wyss, J. Nyberg, A. Johnson, R. Bengtsson, and W. Nazarewicz, Phys. Lett. **B 215**, 211 (1988).