



**HAL**  
open science

## Lifetime difference and CP-violating phase in the $B_{s^0}$ system

V.M. Abazov, B. Abbott, M. Abolins, B.S. Acharya, M. Adams, T. Adams,  
M. Agelou, S.H. Ahn, M. Ahsan, G.D. Alexeev, et al.

► **To cite this version:**

V.M. Abazov, B. Abbott, M. Abolins, B.S. Acharya, M. Adams, et al.. Lifetime difference and CP-violating phase in the  $B_{s^0}$  system. Physical Review Letters, 2007, 98, pp.121801. 10.1103/PhysRevLett.98.121801 . in2p3-00123626

**HAL Id: in2p3-00123626**

**<https://hal.in2p3.fr/in2p3-00123626>**

Submitted on 21 Sep 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Lifetime Difference and CP-Violating Phase in the  $B_s^0$  System

V.M. Abazov,<sup>35</sup> B. Abbott,<sup>75</sup> M. Abolins,<sup>65</sup> B.S. Acharya,<sup>28</sup> M. Adams,<sup>51</sup> T. Adams,<sup>49</sup> E. Aguilo,<sup>5</sup> S.H. Ahn,<sup>30</sup> M. Ahsan,<sup>59</sup> G.D. Alexeev,<sup>35</sup> G. Alkhazov,<sup>39</sup> A. Alton,<sup>64,\*</sup> G. Alverson,<sup>63</sup> G.A. Alves,<sup>2</sup> M. Anastasoie,<sup>34</sup> L.S. Ancu,<sup>34</sup> T. Andeen,<sup>53</sup> S. Anderson,<sup>45</sup> B. Andrieu,<sup>16</sup> M.S. Anzelc,<sup>53</sup> Y. Arnoud,<sup>13</sup> M. Arov,<sup>52</sup> A. Askew,<sup>49</sup> B. Åsman,<sup>40</sup> A.C.S. Assis Jesus,<sup>3</sup> O. Atramentov,<sup>49</sup> C. Autermann,<sup>20</sup> C. Avila,<sup>7</sup> C. Ay,<sup>23</sup> F. Badaud,<sup>12</sup> A. Baden,<sup>61</sup> L. Bagby,<sup>52</sup> B. Baldin,<sup>50</sup> D.V. Bandurin,<sup>59</sup> P. Banerjee,<sup>28</sup> S. Banerjee,<sup>28</sup> E. Barberis,<sup>63</sup> A.-F. Barfuss,<sup>14</sup> P. Bargassa,<sup>80</sup> P. Baringer,<sup>58</sup> C. Barnes,<sup>43</sup> J. Barreto,<sup>2</sup> J.F. Bartlett,<sup>50</sup> U. Bassler,<sup>16</sup> D. Bauer,<sup>43</sup> S. Beale,<sup>5</sup> A. Bean,<sup>58</sup> M. Begalli,<sup>3</sup> M. Begel,<sup>71</sup> C. Belanger-Champagne,<sup>40</sup> L. Bellantoni,<sup>50</sup> A. Bellavance,<sup>67</sup> J.A. Benitez,<sup>65</sup> S.B. Beri,<sup>26</sup> G. Bernardi,<sup>16</sup> R. Bernhard,<sup>22</sup> L. Berntzon,<sup>14</sup> I. Bertram,<sup>42</sup> M. Besançon,<sup>17</sup> R. Beuselinck,<sup>43</sup> V.A. Bezzubov,<sup>38</sup> P.C. Bhat,<sup>50</sup> V. Bhatnagar,<sup>26</sup> M. Binder,<sup>24</sup> C. Biscarat,<sup>19</sup> I. Blackler,<sup>43</sup> G. Blazey,<sup>52</sup> F. Blekman,<sup>43</sup> S. Blessing,<sup>49</sup> D. Bloch,<sup>18</sup> K. Bloom,<sup>67</sup> A. Boehnlein,<sup>50</sup> D. Boline,<sup>62</sup> T.A. Bolton,<sup>59</sup> G. Borissov,<sup>42</sup> K. Bos,<sup>33</sup> T. Bose,<sup>77</sup> A. Brandt,<sup>78</sup> R. Brock,<sup>65</sup> G. Brooijmans,<sup>70</sup> A. Bross,<sup>50</sup> D. Brown,<sup>78</sup> N.J. Buchanan,<sup>49</sup> D. Buchholz,<sup>53</sup> M. Buehler,<sup>81</sup> V. Buescher,<sup>22</sup> S. Burdin,<sup>50</sup> S. Burke,<sup>45</sup> T.H. Burnett,<sup>82</sup> E. Busato,<sup>16</sup> C.P. Buszello,<sup>43</sup> J.M. Butler,<sup>62</sup> P. Calfayan,<sup>24</sup> S. Calvet,<sup>14</sup> J. Cammin,<sup>71</sup> S. Caron,<sup>33</sup> W. Carvalho,<sup>3</sup> B.C.K. Casey,<sup>77</sup> N.M. Cason,<sup>55</sup> H. Castilla-Valdez,<sup>32</sup> S. Chakrabarti,<sup>17</sup> D. Chakraborty,<sup>52</sup> K. Chan,<sup>5</sup> K.M. Chan,<sup>71</sup> A. Chandra,<sup>48</sup> F. Charles,<sup>18</sup> E. Cheu,<sup>45</sup> F. Chevallier,<sup>13</sup> D.K. Cho,<sup>62</sup> S. Choi,<sup>31</sup> B. Choudhary,<sup>27</sup> L. Christofek,<sup>77</sup> T. Christoudias,<sup>43</sup> D. Claes,<sup>67</sup> B. Clément,<sup>18</sup> C. Clément,<sup>40</sup> Y. Coadou,<sup>5</sup> M. Cooke,<sup>80</sup> W.E. Cooper,<sup>50</sup> M. Corcoran,<sup>80</sup> F. Couderc,<sup>17</sup> M.-C. Cousinou,<sup>14</sup> B. Cox,<sup>44</sup> S. Crépe-Renaudin,<sup>13</sup> D. Cutts,<sup>77</sup> M. Cwiok,<sup>29</sup> H. da Motta,<sup>2</sup> A. Das,<sup>62</sup> B. Davies,<sup>42</sup> G. Davies,<sup>43</sup> K. De,<sup>78</sup> P. de Jong,<sup>33</sup> S.J. de Jong,<sup>34</sup> E. De La Cruz-Burelo,<sup>64</sup> C. De Oliveira Martins,<sup>3</sup> J.D. Degenhardt,<sup>64</sup> F. Déliot,<sup>17</sup> M. Demarteau,<sup>50</sup> R. Demina,<sup>71</sup> D. Denisov,<sup>50</sup> S.P. Denisov,<sup>38</sup> S. Desai,<sup>50</sup> H.T. Diehl,<sup>50</sup> M. Diesburg,<sup>50</sup> M. Doidge,<sup>42</sup> A. Dominguez,<sup>67</sup> H. Dong,<sup>72</sup> L.V. Dudko,<sup>37</sup> L. Dufflot,<sup>15</sup> S.R. Dugad,<sup>28</sup> D. Duggan,<sup>49</sup> A. Duperrin,<sup>14</sup> J. Dyer,<sup>65</sup> A. Dyshkant,<sup>52</sup> M. Eads,<sup>67</sup> D. Edmunds,<sup>65</sup> J. Ellison,<sup>48</sup> V.D. Elvira,<sup>50</sup> Y. Enari,<sup>77</sup> S. Eno,<sup>61</sup> P. Ermolov,<sup>37</sup> H. Evans,<sup>54</sup> A. Evdokimov,<sup>36</sup> V.N. Evdokimov,<sup>38</sup> A.V. Ferapontov,<sup>59</sup> T. Ferbel,<sup>71</sup> F. Fiedler,<sup>24</sup> F. Filthaut,<sup>34</sup> W. Fisher,<sup>50</sup> H.E. Fisk,<sup>50</sup> M. Ford,<sup>44</sup> M. Fortner,<sup>52</sup> H. Fox,<sup>22</sup> S. Fu,<sup>50</sup> S. Fuess,<sup>50</sup> T. Gadfort,<sup>82</sup> C.F. Galea,<sup>34</sup> E. Gallas,<sup>50</sup> E. Galyaev,<sup>55</sup> C. Garcia,<sup>71</sup> A. Garcia-Bellido,<sup>82</sup> V. Gavrilov,<sup>36</sup> P. Gay,<sup>12</sup> W. Geist,<sup>18</sup> D. Gelé,<sup>18</sup> C.E. Gerber,<sup>51</sup> Y. Gershtein,<sup>49</sup> D. Gillberg,<sup>5</sup> G. Ginther,<sup>71</sup> N. Gollub,<sup>40</sup> B. Gómez,<sup>7</sup> A. Goussiou,<sup>55</sup> P.D. Grannis,<sup>72</sup> H. Greenlee,<sup>50</sup> Z.D. Greenwood,<sup>60</sup> E.M. Gregores,<sup>4</sup> G. Grenier,<sup>19</sup> Ph. Gris,<sup>12</sup> J.-F. Grivaz,<sup>15</sup> A. Grohsjean,<sup>24</sup> S. Grünendahl,<sup>50</sup> M.W. Grünewald,<sup>29</sup> F. Guo,<sup>72</sup> J. Guo,<sup>72</sup> G. Gutierrez,<sup>50</sup> P. Gutierrez,<sup>75</sup> A. Haas,<sup>70</sup> N.J. Hadley,<sup>61</sup> P. Haefner,<sup>24</sup> S. Hagopian,<sup>49</sup> J. Haley,<sup>68</sup> I. Hall,<sup>75</sup> R.E. Hall,<sup>47</sup> L. Han,<sup>6</sup> K. Hanagaki,<sup>50</sup> P. Hansson,<sup>40</sup> K. Harder,<sup>44</sup> A. Harel,<sup>71</sup> R. Harrington,<sup>63</sup> J.M. Hauptman,<sup>57</sup> R. Hauser,<sup>65</sup> J. Hays,<sup>43</sup> T. Hebbeker,<sup>20</sup> D. Hedin,<sup>52</sup> J.G. Hegeman,<sup>33</sup> J.M. Heinmiller,<sup>51</sup> A.P. Heinson,<sup>48</sup> U. Heintz,<sup>62</sup> C. Hensel,<sup>58</sup> K. Herner,<sup>72</sup> G. Hesketh,<sup>63</sup> M.D. Hildreth,<sup>55</sup> R. Hirosky,<sup>81</sup> J.D. Hobbs,<sup>72</sup> B. Hoeneisen,<sup>11</sup> H. Hoeth,<sup>25</sup> M. Hohlfeld,<sup>15</sup> S.J. Hong,<sup>30</sup> R. Hooper,<sup>77</sup> P. Houben,<sup>33</sup> Y. Hu,<sup>72</sup> Z. Hubacek,<sup>9</sup> V. Hynek,<sup>8</sup> I. Iashvili,<sup>69</sup> R. Illingworth,<sup>50</sup> A.S. Ito,<sup>50</sup> S. Jabeen,<sup>62</sup> M. Jaffré,<sup>15</sup> S. Jain,<sup>75</sup> K. Jakobs,<sup>22</sup> C. Jarvis,<sup>61</sup> A. Jenkins,<sup>43</sup> R. Jesik,<sup>43</sup> K. Johns,<sup>45</sup> C. Johnson,<sup>70</sup> M. Johnson,<sup>50</sup> A. Jonckheere,<sup>50</sup> P. Jonsson,<sup>43</sup> A. Juste,<sup>50</sup> D. Käfer,<sup>20</sup> S. Kahn,<sup>73</sup> E. Kajfasz,<sup>14</sup> A.M. Kalinin,<sup>35</sup> J.M. Kalk,<sup>60</sup> J.R. Kalk,<sup>65</sup> S. Kappler,<sup>20</sup> D. Karmanov,<sup>37</sup> J. Kasper,<sup>62</sup> P. Kasper,<sup>50</sup> I. Katsanos,<sup>70</sup> D. Kau,<sup>49</sup> R. Kaur,<sup>26</sup> R. Kehoe,<sup>79</sup> S. Kermiche,<sup>14</sup> N. Khalatyan,<sup>62</sup> A. Khanov,<sup>76</sup> A. Kharchilava,<sup>69</sup> Y.M. Kharzheev,<sup>35</sup> D. Khatidze,<sup>70</sup> H. Kim,<sup>31</sup> T.J. Kim,<sup>30</sup> M.H. Kirby,<sup>34</sup> B. Klima,<sup>50</sup> J.M. Kohli,<sup>26</sup> J.-P. Konrath,<sup>22</sup> M. Kopal,<sup>75</sup> V.M. Korablev,<sup>38</sup> J. Kotcher,<sup>73</sup> B. Kothari,<sup>70</sup> A. Koubarovsky,<sup>37</sup> A.V. Kozelov,<sup>38</sup> D. Krop,<sup>54</sup> A. Kryemadhi,<sup>81</sup> T. Kuhl,<sup>23</sup> A. Kumar,<sup>69</sup> S. Kunori,<sup>61</sup> A. Kupco,<sup>10</sup> T. Kurča,<sup>19</sup> J. Kvita,<sup>8</sup> D. Lam,<sup>55</sup> S. Lammers,<sup>70</sup> G. Landsberg,<sup>77</sup> J. Lazoflores,<sup>49</sup> P. Lebrun,<sup>19</sup> W.M. Lee,<sup>50</sup> A. Leflat,<sup>37</sup> F. Lehner,<sup>41</sup> V. Lesne,<sup>12</sup> J. Leveque,<sup>45</sup> P. Lewis,<sup>43</sup> J. Li,<sup>78</sup> L. Li,<sup>48</sup> Q.Z. Li,<sup>50</sup> S.M. Lietti,<sup>4</sup> J.G.R. Lima,<sup>52</sup> D. Lincoln,<sup>50</sup> J. Linnemann,<sup>65</sup> V.V. Lipaev,<sup>38</sup> R. Lipton,<sup>50</sup> Z. Liu,<sup>5</sup> L. Lobo,<sup>43</sup> A. Lobodenko,<sup>39</sup> M. Lokajicek,<sup>10</sup> A. Lounis,<sup>18</sup> P. Love,<sup>42</sup> H.J. Lubatti,<sup>82</sup> M. Lynker,<sup>55</sup> A.L. Lyon,<sup>50</sup> A.K.A. Maciel,<sup>2</sup> R.J. Madaras,<sup>46</sup> P. Mättig,<sup>25</sup> C. Magass,<sup>20</sup> A. Magerkurth,<sup>64</sup> N. Makovec,<sup>15</sup> P.K. Mal,<sup>55</sup> H.B. Malbouisson,<sup>3</sup> S. Malik,<sup>67</sup> V.L. Malyshev,<sup>35</sup> H.S. Mao,<sup>50</sup> Y. Maravin,<sup>59</sup> B. Martin,<sup>13</sup> R. McCarthy,<sup>72</sup> A. Melnitchouk,<sup>66</sup> A. Mendes,<sup>14</sup> L. Mendoza,<sup>7</sup> P.G. Mercadante,<sup>4</sup> M. Merkin,<sup>37</sup> K.W. Merritt,<sup>50</sup> A. Meyer,<sup>20</sup> J. Meyer,<sup>21</sup> M. Michaut,<sup>17</sup> H. Miettinen,<sup>80</sup> T. Millet,<sup>19</sup> J. Mitrevski,<sup>70</sup> J. Molina,<sup>3</sup> R.K. Mommsen,<sup>44</sup> N.K. Mondal,<sup>28</sup> J. Monk,<sup>44</sup> R.W. Moore,<sup>5</sup> T. Moulík,<sup>58</sup> G.S. Muanza,<sup>19</sup> M. Mulders,<sup>50</sup> M. Mulhearn,<sup>70</sup> O. Mundal,<sup>22</sup> L. Mundim,<sup>3</sup> E. Nagy,<sup>14</sup> M. Naimuddin,<sup>50</sup> M. Narain,<sup>77</sup> N.A. Naumann,<sup>34</sup> H.A. Neal,<sup>64</sup> J.P. Negret,<sup>7</sup> P. Neustroev,<sup>39</sup> H. Nilsen,<sup>22</sup>

C. Noeding,<sup>22</sup> A. Nomerotski,<sup>50</sup> S.F. Novaes,<sup>4</sup> T. Nunnemann,<sup>24</sup> V. O'Dell,<sup>50</sup> D.C. O'Neil,<sup>5</sup> G. Obrant,<sup>39</sup>  
 C. Ochando,<sup>15</sup> V. Oguri,<sup>3</sup> N. Oliveira,<sup>3</sup> D. Onoprienko,<sup>59</sup> N. Oshima,<sup>50</sup> J. Osta,<sup>55</sup> R. Otec,<sup>9</sup> G.J. Otero y Garzón,<sup>51</sup>  
 M. Owen,<sup>44</sup> P. Padley,<sup>80</sup> M. Pangilinan,<sup>62</sup> N. Parashar,<sup>56</sup> S.-J. Park,<sup>71</sup> S.K. Park,<sup>30</sup> J. Parsons,<sup>70</sup> R. Partridge,<sup>77</sup>  
 N. Parua,<sup>72</sup> A. Patwa,<sup>73</sup> G. Pawloski,<sup>80</sup> P.M. Perea,<sup>48</sup> K. Peters,<sup>44</sup> Y. Peters,<sup>25</sup> P. Pétroff,<sup>15</sup> M. Petteni,<sup>43</sup>  
 R. Piegaia,<sup>1</sup> J. Piper,<sup>65</sup> M.-A. Pleier,<sup>21</sup> P.L.M. Podesta-Lerma,<sup>32,§</sup> V.M. Podstavkov,<sup>50</sup> Y. Pogorelov,<sup>55</sup> M.-E. Pol,<sup>2</sup>  
 A. Pompoš,<sup>75</sup> B.G. Pope,<sup>65</sup> A.V. Popov,<sup>38</sup> C. Potter,<sup>5</sup> W.L. Prado da Silva,<sup>3</sup> H.B. Prosper,<sup>49</sup> S. Protopopescu,<sup>73</sup>  
 J. Qian,<sup>64</sup> A. Quadt,<sup>21</sup> B. Quinn,<sup>66</sup> M.S. Rangel,<sup>2</sup> K.J. Rani,<sup>28</sup> K. Ranjan,<sup>27</sup> P.N. Ratoff,<sup>42</sup> P. Renkel,<sup>79</sup>  
 S. Reucroft,<sup>63</sup> M. Rijssenbeek,<sup>72</sup> I. Ripp-Baudot,<sup>18</sup> F. Rizatdinova,<sup>76</sup> S. Robinson,<sup>43</sup> R.F. Rodrigues,<sup>3</sup>  
 C. Royon,<sup>17</sup> P. Rubinov,<sup>50</sup> R. Ruchti,<sup>55</sup> G. Sajot,<sup>13</sup> A. Sánchez-Hernández,<sup>32</sup> M.P. Sanders,<sup>16</sup> A. Santoro,<sup>3</sup>  
 G. Savage,<sup>50</sup> L. Sawyer,<sup>60</sup> T. Scanlon,<sup>43</sup> D. Schaile,<sup>24</sup> R.D. Schamberger,<sup>72</sup> Y. Scheglov,<sup>39</sup> H. Schellman,<sup>53</sup>  
 P. Schieferdecker,<sup>24</sup> C. Schmitt,<sup>25</sup> C. Schwanenberger,<sup>44</sup> A. Schwartzman,<sup>68</sup> R. Schwienhorst,<sup>65</sup> J. Sekaric,<sup>49</sup>  
 S. Sengupta,<sup>49</sup> H. Severini,<sup>75</sup> E. Shabalina,<sup>51</sup> M. Shamim,<sup>59</sup> V. Shary,<sup>17</sup> A.A. Shchukin,<sup>38</sup> R.K. Shivpuri,<sup>27</sup>  
 D. Shpakov,<sup>50</sup> V. Siccardi,<sup>18</sup> R.A. Sidwell,<sup>59</sup> V. Simak,<sup>9</sup> V. Sirotenko,<sup>50</sup> P. Skubic,<sup>75</sup> P. Slattery,<sup>71</sup> D. Smirnov,<sup>55</sup>  
 R.P. Smith,<sup>50</sup> G.R. Snow,<sup>67</sup> J. Snow,<sup>74</sup> S. Snyder,<sup>73</sup> S. Söldner-Rembold,<sup>44</sup> L. Sonnenschein,<sup>16</sup> A. Sopczak,<sup>42</sup>  
 M. Sosebee,<sup>78</sup> K. Soustruznik,<sup>8</sup> M. Souza,<sup>2</sup> B. Spurlock,<sup>78</sup> J. Stark,<sup>13</sup> J. Steele,<sup>60</sup> V. Stolin,<sup>36</sup> A. Stone,<sup>51</sup>  
 D.A. Stoyanova,<sup>38</sup> J. Strandberg,<sup>64</sup> S. Strandberg,<sup>40</sup> M.A. Strang,<sup>69</sup> M. Strauss,<sup>75</sup> R. Ströhmer,<sup>24</sup> D. Strom,<sup>53</sup>  
 M. Strovink,<sup>46</sup> L. Stutte,<sup>50</sup> S. Sumowidagdo,<sup>49</sup> P. Svoisky,<sup>55</sup> A. Sznajder,<sup>3</sup> M. Talby,<sup>14</sup> P. Tamburello,<sup>45</sup>  
 W. Taylor,<sup>5</sup> P. Telford,<sup>44</sup> J. Temple,<sup>45</sup> B. Tiller,<sup>24</sup> F. Tissandier,<sup>12</sup> M. Titov,<sup>22</sup> V.V. Tokmenin,<sup>35</sup> M. Tomoto,<sup>50</sup>  
 T. Toole,<sup>61</sup> I. Torchiani,<sup>22</sup> T. Trefzger,<sup>23</sup> S. Trincaz-Duvoid,<sup>16</sup> D. Tsybychev,<sup>72</sup> B. Tuchming,<sup>17</sup> C. Tully,<sup>68</sup>  
 P.M. Tuts,<sup>70</sup> R. Unalan,<sup>65</sup> L. Uvarov,<sup>39</sup> S. Uvarov,<sup>39</sup> S. Uzunyan,<sup>52</sup> B. Vachon,<sup>5</sup> P.J. van den Berg,<sup>33</sup> B. van Eijk,<sup>35</sup>  
 R. Van Kooten,<sup>54</sup> W.M. van Leeuwen,<sup>33</sup> N. Varelas,<sup>51</sup> E.W. Varnes,<sup>45</sup> A. Vartapetian,<sup>78</sup> I.A. Vasilyev,<sup>38</sup>  
 M. Vaupel,<sup>25</sup> P. Verdier,<sup>19</sup> L.S. Vertogradov,<sup>35</sup> M. Verzocchi,<sup>50</sup> F. Villeneuve-Seguiet,<sup>43</sup> P. Vint,<sup>43</sup> J.-R. Vlimant,<sup>16</sup>  
 E. Von Toerne,<sup>59</sup> M. Voutilainen,<sup>67,‡</sup> M. Vreeswijk,<sup>33</sup> H.D. Wahl,<sup>49</sup> L. Wang,<sup>61</sup> M.H.L.S Wang,<sup>50</sup> J. Warchol,<sup>55</sup>  
 G. Watts,<sup>82</sup> M. Wayne,<sup>55</sup> G. Weber,<sup>23</sup> M. Weber,<sup>50</sup> H. Weerts,<sup>65</sup> A. Wenger,<sup>22,#</sup> N. Wermes,<sup>21</sup> M. Wetstein,<sup>61</sup>  
 A. White,<sup>78</sup> D. Wicke,<sup>25</sup> G.W. Wilson,<sup>58</sup> S.J. Wimpenny,<sup>48</sup> M. Wobisch,<sup>50</sup> D.R. Wood,<sup>63</sup> T.R. Wyatt,<sup>44</sup>  
 Y. Xie,<sup>77</sup> S. Yacoob,<sup>53</sup> R. Yamada,<sup>50</sup> M. Yan,<sup>61</sup> T. Yasuda,<sup>50</sup> Y.A. Yatsunenko,<sup>35</sup> K. Yip,<sup>73</sup> H.D. Yoo,<sup>77</sup>  
 S.W. Youn,<sup>53</sup> C. Yu,<sup>13</sup> J. Yu,<sup>78</sup> A. Yurkewicz,<sup>72</sup> A. Zatserklyaniy,<sup>52</sup> C. Zeitnitz,<sup>25</sup> D. Zhang,<sup>50</sup> T. Zhao,<sup>82</sup>  
 B. Zhou,<sup>64</sup> J. Zhu,<sup>72</sup> M. Zielinski,<sup>71</sup> D. Zieminska,<sup>54</sup> A. Zieminski,<sup>54</sup> V. Zutshi,<sup>52</sup> and E.G. Zverev<sup>37</sup>  
 (DØ Collaboration)

<sup>1</sup> Universidad de Buenos Aires, Buenos Aires, Argentina

<sup>2</sup> LAFEX, Centro Brasileiro de Pesquisas Físicas, Rio de Janeiro, Brazil

<sup>3</sup> Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil

<sup>4</sup> Instituto de Física Teórica, Universidade Estadual Paulista, São Paulo, Brazil

<sup>5</sup> University of Alberta, Edmonton, Alberta, Canada, Simon Fraser University, Burnaby, British Columbia, Canada, York University, Toronto, Ontario, Canada, and McGill University, Montreal, Quebec, Canada

<sup>6</sup> University of Science and Technology of China, Hefei, People's Republic of China

<sup>7</sup> Universidad de los Andes, Bogotá, Colombia

<sup>8</sup> Center for Particle Physics, Charles University, Prague, Czech Republic

<sup>9</sup> Czech Technical University, Prague, Czech Republic

<sup>10</sup> Center for Particle Physics, Institute of Physics, Academy of Sciences of the Czech Republic, Prague, Czech Republic

<sup>11</sup> Universidad San Francisco de Quito, Quito, Ecuador

<sup>12</sup> Laboratoire de Physique Corpusculaire, IN2P3-CNRS, Université Blaise Pascal, Clermont-Ferrand, France

<sup>13</sup> Laboratoire de Physique Subatomique et de Cosmologie, IN2P3-CNRS, Université de Grenoble 1, Grenoble, France

<sup>14</sup> CPPM, IN2P3-CNRS, Université de la Méditerranée, Marseille, France

<sup>15</sup> Laboratoire de l'Accélérateur Linéaire, IN2P3-CNRS et Université Paris-Sud, Orsay, France

<sup>16</sup> LPNHE, IN2P3-CNRS, Universités Paris VI and VII, Paris, France

<sup>17</sup> DAPNIA/Service de Physique des Particules, CEA, Saclay, France

<sup>18</sup> IPHC, IN2P3-CNRS, Université Louis Pasteur, Strasbourg, France, and Université de Haute Alsace, Mulhouse, France

<sup>19</sup> IPNL, Université Lyon 1, CNRS/IN2P3, Villeurbanne, France and Université de Lyon, Lyon, France

<sup>20</sup> III. Physikalisches Institut A, RWTH Aachen, Aachen, Germany

<sup>21</sup> Physikalisches Institut, Universität Bonn, Bonn, Germany

<sup>22</sup> Physikalisches Institut, Universität Freiburg, Freiburg, Germany

<sup>23</sup> Institut für Physik, Universität Mainz, Mainz, Germany

<sup>24</sup> Ludwig-Maximilians-Universität München, München, Germany

<sup>25</sup> Fachbereich Physik, University of Wuppertal, Wuppertal, Germany

<sup>26</sup> Panjab University, Chandigarh, India

<sup>27</sup> Delhi University, Delhi, India

- <sup>28</sup> *Tata Institute of Fundamental Research, Mumbai, India*  
<sup>29</sup> *University College Dublin, Dublin, Ireland*  
<sup>30</sup> *Korea Detector Laboratory, Korea University, Seoul, Korea*  
<sup>31</sup> *SungKyunKwan University, Suwon, Korea*  
<sup>32</sup> *CINVESTAV, Mexico City, Mexico*  
<sup>33</sup> *FOM-Institute NIKHEF and University of Amsterdam/NIKHEF, Amsterdam, The Netherlands*  
<sup>34</sup> *Radboud University Nijmegen/NIKHEF, Nijmegen, The Netherlands*  
<sup>35</sup> *Joint Institute for Nuclear Research, Dubna, Russia*  
<sup>36</sup> *Institute for Theoretical and Experimental Physics, Moscow, Russia*  
<sup>37</sup> *Moscow State University, Moscow, Russia*  
<sup>38</sup> *Institute for High Energy Physics, Protvino, Russia*  
<sup>39</sup> *Petersburg Nuclear Physics Institute, St. Petersburg, Russia*  
<sup>40</sup> *Lund University, Lund, Sweden, Royal Institute of Technology and Stockholm University, Stockholm, Sweden, and Uppsala University, Uppsala, Sweden*  
<sup>41</sup> *Physik Institut der Universität Zürich, Zürich, Switzerland*  
<sup>42</sup> *Lancaster University, Lancaster, United Kingdom*  
<sup>43</sup> *Imperial College, London, United Kingdom*  
<sup>44</sup> *University of Manchester, Manchester, United Kingdom*  
<sup>45</sup> *University of Arizona, Tucson, Arizona 85721, USA*  
<sup>46</sup> *Lawrence Berkeley National Laboratory and University of California, Berkeley, California 94720, USA*  
<sup>47</sup> *California State University, Fresno, California 93740, USA*  
<sup>48</sup> *University of California, Riverside, California 92521, USA*  
<sup>49</sup> *Florida State University, Tallahassee, Florida 32306, USA*  
<sup>50</sup> *Fermi National Accelerator Laboratory, Batavia, Illinois 60510, USA*  
<sup>51</sup> *University of Illinois at Chicago, Chicago, Illinois 60607, USA*  
<sup>52</sup> *Northern Illinois University, DeKalb, Illinois 60115, USA*  
<sup>53</sup> *Northwestern University, Evanston, Illinois 60208, USA*  
<sup>54</sup> *Indiana University, Bloomington, Indiana 47405, USA*  
<sup>55</sup> *University of Notre Dame, Notre Dame, Indiana 46556, USA*  
<sup>56</sup> *Purdue University Calumet, Hammond, Indiana 46323, USA*  
<sup>57</sup> *Iowa State University, Ames, Iowa 50011, USA*  
<sup>58</sup> *University of Kansas, Lawrence, Kansas 66045, USA*  
<sup>59</sup> *Kansas State University, Manhattan, Kansas 66506, USA*  
<sup>60</sup> *Louisiana Tech University, Ruston, Louisiana 71272, USA*  
<sup>61</sup> *University of Maryland, College Park, Maryland 20742, USA*  
<sup>62</sup> *Boston University, Boston, Massachusetts 02215, USA*  
<sup>63</sup> *Northeastern University, Boston, Massachusetts 02115, USA*  
<sup>64</sup> *University of Michigan, Ann Arbor, Michigan 48109, USA*  
<sup>65</sup> *Michigan State University, East Lansing, Michigan 48824, USA*  
<sup>66</sup> *University of Mississippi, University, Mississippi 38677, USA*  
<sup>67</sup> *University of Nebraska, Lincoln, Nebraska 68588, USA*  
<sup>68</sup> *Princeton University, Princeton, New Jersey 08544, USA*  
<sup>69</sup> *State University of New York, Buffalo, New York 14260, USA*  
<sup>70</sup> *Columbia University, New York, New York 10027, USA*  
<sup>71</sup> *University of Rochester, Rochester, New York 14627, USA*  
<sup>72</sup> *State University of New York, Stony Brook, New York 11794, USA*  
<sup>73</sup> *Brookhaven National Laboratory, Upton, New York 11973, USA*  
<sup>74</sup> *Langston University, Langston, Oklahoma 73050, USA*  
<sup>75</sup> *University of Oklahoma, Norman, Oklahoma 73019, USA*  
<sup>76</sup> *Oklahoma State University, Stillwater, Oklahoma 74078, USA*  
<sup>77</sup> *Brown University, Providence, Rhode Island 02912, USA*  
<sup>78</sup> *University of Texas, Arlington, Texas 76019, USA*  
<sup>79</sup> *Southern Methodist University, Dallas, Texas 75275, USA*  
<sup>80</sup> *Rice University, Houston, Texas 77005, USA*  
<sup>81</sup> *University of Virginia, Charlottesville, Virginia 22901, USA*  
<sup>82</sup> *University of Washington, Seattle, Washington 98195, USA*

(Dated: January 9, 2007)

From an analysis of the decay  $B_s^0 \rightarrow J/\psi\phi$  we obtain the width difference between the light and heavy mass eigenstates,  $\Delta\Gamma \equiv (\Gamma_L - \Gamma_H) = 0.17 \pm 0.09$  (stat)  $\pm 0.02$  (syst)  $\text{ps}^{-1}$  and the CP-violating phase  $\phi_s = -0.79 \pm 0.56$  (stat)  $^{+0.14}_{-0.01}$  (syst). Under the hypothesis of no CP violation ( $\phi_s \equiv 0$ ), we obtain  $1/\bar{\Gamma} = \bar{\tau}(B_s^0) = 1.52 \pm 0.08^{+0.01}_{-0.03}$  ps and  $\Delta\Gamma = 0.12^{+0.08}_{-0.10} \pm 0.02$   $\text{ps}^{-1}$ . The data sample corresponds to an integrated luminosity of  $1.1 \text{ fb}^{-1}$  accumulated with the D0 detector at

the Tevatron.

PACS numbers: 13.25.Hw, 11.30.Er

In the standard model (SM), the light ( $L$ ) and heavy ( $H$ ) eigenstates of the mixed  $B_s^0$  system are expected to have a sizeable mass and decay width difference,  $\Delta M \equiv M_H - M_L$  and  $\Delta\Gamma \equiv \Gamma_L - \Gamma_H$ . The CP-violating phase, defined as the relative phase of the off-diagonal elements of the mass and decay matrices in the  $B_s^0 - \bar{B}_s^0$  basis, is predicted to be small. Thus, to a good approximation the two mass eigenstates are expected to be CP eigenstates. New phenomena may alter the CP-violating mixing phase  $\phi_s$ , leading to a reduction of the observed  $\Delta\Gamma$  compared to the SM prediction [1]  $\Delta\Gamma_{SM}$ :  $\Delta\Gamma = \Delta\Gamma_{SM} \times \cos\phi_s$ . While the mass difference has recently been measured to high precision [2, 3], the CP-violating phase remains unknown.

The decay  $B_s^0 \rightarrow J/\psi\phi$ , proceeding through the quark process  $b \rightarrow c\bar{c}s$ , gives rise to both CP-even and CP-odd final states. It is possible to separate the two CP components of the decay  $B_s^0 \rightarrow J/\psi\phi$ , and thus to measure the lifetime difference, through a study of the time-dependent angular distribution of the decay products of the  $J/\psi$  and  $\phi$  mesons. Moreover, with a sizeable lifetime difference, there is sensitivity to the mixing phase through the interference terms between the CP-even and CP-odd waves.

In Ref. [4] we presented an analysis of the decay chain  $B_s^0 \rightarrow J/\psi\phi$ ,  $J/\psi \rightarrow \mu^+\mu^-$ ,  $\phi \rightarrow K^+K^-$  based on the first  $\approx 450 \text{ pb}^{-1}$  of  $p\bar{p}$  data at a center-of-mass energy of 1.96 TeV collected with the D0 detector [5]. In that analysis, we extracted three parameters characterizing the  $B_s^0$  system and its decay  $B_s^0 \rightarrow J/\psi\phi$ : the average lifetime,  $\bar{\tau} = 1/\bar{\Gamma}$ , where  $\bar{\Gamma} \equiv (\Gamma_H + \Gamma_L)/2$ ;  $\Delta\Gamma/\bar{\Gamma}$ ; and the relative rate of the decay to the CP-odd states at time zero. Here we present new results, based on a two-fold increase in statistics. In addition to  $\bar{\tau}$  and  $\Delta\Gamma$ , we extract for the first time the CP-violating phase  $\phi_s$ . We also measure the magnitudes of the decay amplitudes, and their relative phases.

The data, collected between June 2002 and January 2006, correspond to an integrated luminosity of  $1.1 \text{ fb}^{-1}$ . The selected events include two reconstructed muons of opposite charge, with a transverse momentum greater than 1.5 GeV and pseudorapidity  $|\eta| < 2$ . Each muon is required to be detected as a track segment in at least one of the three layers of the muon system, and to be matched to a central track. One muon is required to have segments both inside and outside the toroid magnet. We require the events to satisfy a muon trigger that does not include a cut on the impact parameter.

To select the  $B_s^0$  candidate sample, we set the minimum values of momenta in the transverse plane for  $B_s^0$ ,  $\phi$ , and  $K$  meson candidates at 6.0 GeV, 1.5 GeV, and 0.7 GeV, respectively.  $J/\psi$  candidates are accepted if




FIG. 1: The invariant mass distribution of the  $(J/\psi, \phi)$  system for  $B_s^0$  candidates. The curves are projections of the maximum likelihood fit (see text).

the invariant mass of the muon pair is in the range 2.9 – 3.3 GeV. For events in the central rapidity region (an event is considered to be central if the higher  $p_T$  muon has  $|\eta_{\mu 1}| < 1$ ), we require the transverse momentum of the  $J/\psi$  meson to exceed 4 GeV. Successful candidates are constrained to the world average mass of the  $J/\psi$  meson [6]. Decay products of the  $\phi$  candidates are required to satisfy a fit to a common vertex and to have an invariant mass in the range 1.01 – 1.03 GeV. We require the  $(J/\psi, \phi)$  pair to be consistent with coming from a common vertex, and to have an invariant mass in the range 5.0 – 5.8 GeV. In the case of multiple  $\phi$  meson candidates, we select the one with the highest transverse momentum. Monte Carlo (MC) studies show that the  $p_T$  spectrum of the  $\phi$  mesons coming from  $B_s^0$  decay is harder than the spectrum of a pair of random tracks from hadronization. We define the signed decay length of a  $B_s^0$  meson  $L_{xy}^B$  as the vector pointing from the primary vertex to the decay vertex projected on the  $B_s^0$  transverse momentum. To reconstruct the primary vertex, we select tracks with  $p_T > 0.3 \text{ GeV}$  that are not used as decay products of the  $B_s^0$  candidate, and apply a constraint to the average beam spot position. The proper decay length,  $ct$ , is defined by the relation  $ct = L_{xy}^B \cdot M_{B_s^0} / p_T$  where  $M_{B_s^0}$  is the measured mass of the  $B_s^0$  candidate. The distribution of the proper decay length uncertainty  $\sigma(ct)$  of  $B_s^0$  mesons peaks around  $25 \mu\text{m}$ . We accept events with  $\sigma(ct) < 60 \mu\text{m}$ . The invariant mass distribution of the accepted 23343 candidates is shown in Fig. 1. The curves are projections of the maximum likelihood fit, described below. The fit assigns  $1039 \pm 45$  (stat) events to the  $B_s^0$  decay.

We perform a simultaneous unbinned maximum likelihood fit to the proper decay length, three decay angles, and mass. The likelihood function  $\mathcal{L}$  is given by:

$$\mathcal{L} = \prod_{i=1}^N [f_{sig} \mathcal{F}_{sig}^i + (1 - f_{sig}) \mathcal{F}_{bck}^i], \quad (1)$$

where  $N$  is the total number of events, and  $f_{sig}$  is the fraction of signal in the sample. The function  $\mathcal{F}_{sig}^i$  describes the distribution of the signal in mass, proper decay length, and the decay angles, and  $\mathcal{F}_{bck}^i$  is the product of the background mass, proper decay length, and angular probability density functions. Background is di-

vided into two categories. ‘‘Prompt’’ background is due to directly produced  $J/\psi$  mesons accompanied by random tracks arising from hadronization. This background is distinguished from ‘‘non-prompt’’ background, where the  $J/\psi$  meson is a product of a  $B$  hadron decay while the tracks forming the  $\phi$  candidate emanate from a multi-body decay of the same  $B$  hadron or from hadronization.

The time evolution of the angular distribution of the products of the decay of *flavor untagged*  $B_s^0$  mesons, i.e., summed over  $B_s^0$  and  $\bar{B}_s^0$ , expressed in terms of the linear polarization amplitudes  $A_x$  and their relative phases  $\delta_i$  is [1]:

$$\begin{aligned} \frac{d^3\Gamma(t)}{d\cos\theta d\varphi d\cos\psi} &\propto 2|A_0(0)|^2 \mathcal{T}_+ \cos^2\psi (1 - \sin^2\theta \cos^2\varphi) + \sin^2\psi \{ |A_{\parallel}(0)|^2 \mathcal{T}_+ (1 - \sin^2\theta \sin^2\varphi) + |A_{\perp}(0)|^2 \mathcal{T}_- \sin^2\theta \} \\ &+ \frac{1}{\sqrt{2}} \sin 2\psi |A_0(0)| |A_{\parallel}(0)| \cos(\delta_2 - \delta_1) \mathcal{T}_+ \sin^2\theta \sin 2\varphi \\ &+ \left\{ \frac{1}{\sqrt{2}} |A_0(0)| |A_{\perp}(0)| \cos \delta_2 \sin 2\psi \sin 2\theta \cos \varphi \right. \\ &\left. - |A_{\parallel}(0)| |A_{\perp}(0)| \cos \delta_1 \sin^2\psi \sin 2\theta \sin \varphi \right\} \frac{1}{2} (e^{-\Gamma_H t} - e^{-\Gamma_L t}) \sin \phi_s. \end{aligned} \quad (2)$$

where  $\mathcal{T}_{+/-} = \frac{1}{2} ((1 \pm \cos \phi_s) e^{-\Gamma_L t} + (1 \mp \cos \phi_s) e^{-\Gamma_H t})$ .

In the coordinate system of the  $J/\psi$  rest frame (where the  $\phi$  meson moves in the  $x$  direction, the  $z$  axis is perpendicular to the decay plane of  $\phi \rightarrow K^+K^-$ , and  $p_y(K^+) \geq 0$ ), the transversity polar and azimuthal angles  $(\theta, \varphi)$  describe the direction of the  $\mu^+$ , and  $\psi$  is the angle between  $\vec{p}(K^+)$  and  $-\vec{p}(J/\psi)$  in the  $\phi$  rest frame.

We model the acceptance in the three angles by fits using polynomial functions, with parameters determined using Monte Carlo simulations. We have used the *SVV\_HELAMP* model in the *EVTGEN* generator [7], interfaced to the *PYTHIA* program [8]. Simulated events were reweighted to match the kinematic distributions observed in the data.

The lifetime distribution shape of the background is described as a sum of a prompt component, simulated as a Gaussian function centered at zero, and a non-prompt component, simulated as a superposition of one exponential for the negative  $ct$  region and two exponentials for the positive  $ct$  region, with free slopes and normalization. The mass distributions of the backgrounds are parametrized by first-order polynomials. The distributions in the transversity polar and azimuthal an-

gles are parametrized as  $1 + X_{2x} \cos^2\theta + X_{4x} \cos^4\theta$  and  $1 + Y_{1x} \cos(2\varphi) + Y_{2x} \cos^2(2\varphi)$ , respectively. For the background dependence on the angle  $\psi$ , we use the function  $1 + Z_{2x} \cos^2(\psi)$ . We also allow for a background term analogous to the interference term of the CP-even waves, with one free coefficient. For each of the above background functions we use two separate sets of parameters for the prompt and non-prompt components.

Our results for the hypothesis of CP conservation and for the case of free  $\phi_s$ , are presented in Table I. For the CP-violating phase, which has a four-fold ambiguity discussed below, the fit value closest to the SM prediction of  $-0.03$  [1] is  $\phi_s = -0.79 \pm 0.56$ . Figures 2 – 5 show the fit projections on the angular distributions and the proper decay length. Figure 6 shows the  $\Delta \ln(\mathcal{L}) = 0.5$  error ellipse contour (corresponding to the confidence level of 39%) in the plane  $(\Delta\Gamma, \phi_s)$ . As seen from Eq. 2, the sign of  $\sin \phi_s$  is reversed with the simultaneous reversal of the signs of  $\cos \delta_1$  and  $\cos \delta_2$ . For the case  $\cos \delta_1 < 0$  and  $\cos \delta_2 > 0$ , our measurement correlates two possible solutions for  $\phi_s$  with the two signs of  $\Delta\Gamma$ :  $\phi_s = -0.79 \pm 0.56$ ,  $\Delta\Gamma > 0$ , and  $\phi_s = 2.35 \pm 0.56$ ,  $\Delta\Gamma < 0$ . For the case  $\cos \delta_1 > 0$  and  $\cos \delta_2 < 0$  the two solutions

are  $\phi_s = 0.79 \pm 0.56$ ,  $\Delta\Gamma > 0$ , and  $\phi_s = -2.35 \pm 0.56$ ,  $\Delta\Gamma < 0$ .

TABLE I: Maximum likelihood fit results. Sign ambiguities are discussed in the text.

Observable	CP conserved	free $\phi_s$
$\Delta\Gamma$ (ps <sup>-1</sup> )	$0.12^{+0.08}_{-0.10}$	$0.17^{+0.09}_{-0.09}$
$\frac{1}{\Gamma} = \bar{\tau}$ (ps)	$1.52^{+0.08}_{-0.08}$	$1.49 \pm 0.08$
$\phi_s$	$\equiv 0$	$-0.79 \pm 0.56$
$ A_0(0) ^2 -  A_{\parallel}(0) ^2$	$0.38 \pm 0.05$	$0.37 \pm 0.06$
$A_{\perp}(0)$	$0.45 \pm 0.05$	$0.46 \pm 0.06$
$\delta_1 - \delta_2$	$2.6 \pm 0.4$	$2.6 \pm 0.4$
$\delta_1$	–	$3.3 \pm 1.0$
$\delta_2$	–	$0.7 \pm 1.1$




FIG. 2: The transversity polar angle distribution for the signal-enhanced subsample:  $ct/\sigma(ct) > 5$  and signal mass range. The curves show: the signal contribution, dotted (red); the background, light solid (green); and total, solid (blue) [color online].

We perform a test using pseudo-experiments with similar statistical sensitivity, generated with the same parameters as obtained in this analysis under the condition of no CP violation. When fits allowing for CP violation are performed,  $\approx 50\%$  of the experiments have a fitted  $\cos(\phi_s)$  less than the measured value. About 80% of experiments have the statistical uncertainty of  $\phi_s$  greater than that for data.

We verify the procedure by performing fits on MC samples passed through the full chain of detector simulation, event reconstruction, and maximum likelihood fitting. We assign systematic uncertainties due to the statistical precision of this procedure test. We also repeat the fits to the data with the parameters describing the acceptance varied by  $\pm 1\sigma$ .




FIG. 3: The transversity azimuthal angle distribution for the signal-enhanced subsample:  $ct/\sigma(ct) > 5$  and signal mass range. The curves show: the signal contribution, dotted (red); the background, light solid (green); and total, solid (blue) [color online].




FIG. 4: The  $\psi$  angle distribution for the signal-enhanced subsample:  $ct/\sigma(ct) > 5$  and signal mass range. The curves show: the signal contribution, dotted (red); the background, light solid (green); and total, solid (blue) [color online].

Uncertainties from the data processing reflect the stability of the results with respect to different versions of the track and vertex reconstruction algorithms. The “interference” term in the background model accounts for the collective effect of various physics processes. However, its presence may be partially due to the detector acceptance effects. Therefore, we interpret the difference between fits with and without this term as a systematic uncertainty associated with the background model. Effects of the imperfect detector alignment are estimated using a modified geometry of the the silicon microstrip tracker, with silicon sensors moved within the known uncertainty. The effects of systematic uncertainties are listed in Table II.

From a fit to the CP-conserving time-dependent angu-




FIG. 5: The proper decay length,  $ct$ , of the  $B_s^0$  candidates in the signal mass region. The curves show: the signal contribution, dashed (red); the CP-even (dotted) and CP-odd (dashed-dotted) contributions of the signal, the background, light solid (green); and total, solid (blue) [color online].

TABLE II: Sources of systematic uncertainty in the results of the analysis of the decay  $B_s^0 \rightarrow J/\psi\phi$ .

Source	$c\tau(B_s^0)$ $\mu\text{m}$	$\Delta\Gamma$ $\text{ps}^{-1}$	$R_\perp$	$\phi_s$
Procedure test	$\pm 2.0$	$\pm 0.02$	$\pm 0.01$	—
Acceptance	$\pm 0.5$	$\pm 0.001$	$\pm 0.003$	$\pm 0.01$
Reco. algorithm	$-8.0, +1.3$	$+0.001$	$\pm 0.01$	$-0.01$
Background model	$+1.0$	$+0.01$	$-0.01$	$+0.14$
Alignment	$\pm 2.0$	—	—	—
Total	$-8.8, +3.3$	$\pm 0.02$	$\pm 0.02$	$-0.01, +0.14$




FIG. 6: The  $\Delta \ln(\mathcal{L}) = 0.5$  contour (error ellipse) in the plane  $(\Delta\Gamma, \phi_s)$  for the fit to the  $B_s^0 \rightarrow J/\psi\phi$  data. Also shown is the band representing the relation  $\Delta\Gamma = \Delta\Gamma_{SM} \times |\cos(\phi_s)|$ , with  $\Delta\Gamma_{SM} = 0.10 \pm 0.03 \text{ ps}^{-1}$  [10]. The 4-fold ambiguity is discussed in the text.

lar distribution of the untagged decay  $B_s^0 \rightarrow J/\psi\phi$ , we obtain the average lifetime of the  $B_s^0$  system,  $\bar{\tau}(B_s^0) = 1.52 \pm 0.08 \text{ (stat)} \pm 0.03 \text{ (syst)} \text{ ps}$  and the width difference between the two mass eigenstates,  $\Delta\Gamma = 0.12 \pm 0.08 \text{ (stat)} \pm 0.02 \text{ (syst)} \text{ ps}^{-1}$ .

Allowing for CP violation in  $B_s^0$  mixing, we provide the first direct constraint on the CP-violating phase,  $\phi_s = -0.79 \pm 0.56 \text{ (stat)} \pm 0.01 \text{ (syst)}$ .

We thank U. Nierste for useful discussions. We thank the staffs at Fermilab and collaborating institutions, and acknowledge support from the DOE and NSF (USA); CEA and CNRS/IN2P3 (France); FASI, Rosatom and RFBR (Russia); CAPES, CNPq, FAPERJ, FAPESP and FUNDUNESP (Brazil); DAE and DST (India); Colciencias (Colombia); CONACyT (Mexico); KRF and KOSEF (Korea); CONICET and UBACyT (Argentina); FOM (The Netherlands); PPARC (United Kingdom); MSMT (Czech Republic); CRC Program, CFI, NSERC and WestGrid Project (Canada); BMBF and DFG (Germany); SFI (Ireland); The Swedish Research Council (Sweden); Research Corporation; Alexander von Humboldt Foundation; and the Marie Curie Program.

- [1] I. Dunietz, R. Fleischer, and U. Nierste, hep-ph/0012219.
- [2] D0 Collaboration, V.M. Abazov *et al.*, Phys. Rev. Lett. **97**, 021802 (2006).
- [3] A. Abulencia *et al.* [CDF Collaboration], submitted to Phys. Rev. Lett., arXiv:hep-ex/0609040.
- [4] D0 Collaboration, V. M. Abazov *et al.*, Phys. Rev. Lett. **95**, 171801 (2005).
- [5] V.M. Abazov *et al.* (D0 Collaboration), “The upgraded D0 detector,” Nucl. Instrum. Methods Phys. Res. A **565**, 463 (2006).
- [6] W. M. Yao *et al.* (Particle Data Group), J. Phys. G **33**, 1(2006) (URL: <http://pdg.lbl.gov>).
- [7] A. Ryd, D. Lange, <http://www.slac.stanford.edu/~lange/EvtGen/>
- [8] H. U. Bengtsson and T. Sjostrand, Comp. Phys. Comm. **46**, 43 (1987).
- [9] K. Anikeev *et al.*, “B physics at the Tevatron: Run II and beyond,” FERMILAB-Pub-01/197, hep-ph/0201071, p. 360.
- [10] M. Beneke, G. Buchalla, C. Greub, A. Lenz and U. Nierste, “Next-to-leading order QCD corrections to the lifetime difference of B/s mesons,” Phys. Lett. B **459** (1999) 631 [arXiv:hep-ph/9808385]; input parameters updated in March 2006.