



**HAL**  
open science

## New Measurements of the Transverse Beam Asymmetry for Elastic Electron Scattering from Selected Nuclei

S. Abrahamyan, A. Acha, A. Afanasev, Z. Ahmed, H. Albataineh, K. Aniol,  
D. S. Armstrong, W. Armstrong, J. Arrington, T. Averett, et al.

► **To cite this version:**

S. Abrahamyan, A. Acha, A. Afanasev, Z. Ahmed, H. Albataineh, et al.. New Measurements of the Transverse Beam Asymmetry for Elastic Electron Scattering from Selected Nuclei. *Physical Review Letters*, 2012, 109, pp.192501. 10.1103/PhysRevLett.109.192501 . in2p3-00745218

**HAL Id: in2p3-00745218**

**<https://hal.in2p3.fr/in2p3-00745218>**

Submitted on 14 Nov 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## New Measurements of the Transverse Beam Asymmetry for Elastic Electron Scattering from Selected Nuclei

S. Abrahamyan,<sup>45</sup> A. Acha,<sup>10</sup> A. Afanasev,<sup>11</sup> Z. Ahmed,<sup>33</sup> H. Albataineh,<sup>6</sup> K. Aniol,<sup>3</sup> D. S. Armstrong,<sup>7</sup> W. Armstrong,<sup>35</sup> J. Arrington,<sup>1</sup> T. Averett,<sup>7</sup> B. Babineau,<sup>23</sup> S. L. Bailey,<sup>7</sup> J. Barber,<sup>40</sup> A. Barbieri,<sup>43</sup> A. Beck,<sup>25</sup> V. Bellini,<sup>15</sup> R. Beminiwattha,<sup>27</sup> H. Benaoum,<sup>33</sup> J. Benesch,<sup>36</sup> F. Benmokhtar,<sup>8</sup> P. Bertin,<sup>6</sup> T. Bielarski,<sup>41</sup> W. Boeglin,<sup>10</sup> P. Bosted,<sup>36</sup> F. Butaru,<sup>5</sup> E. Burtin,<sup>5</sup> J. Cahoon,<sup>40</sup> A. Camsonne,<sup>36</sup> M. Canan,<sup>28</sup> P. Carter,<sup>8</sup> C. C. Chang,<sup>39</sup> G. D. Cates,<sup>43</sup> Y.-C. Chao,<sup>36</sup> C. Chen,<sup>12</sup> J.-P. Chen,<sup>36</sup> Seonho Choi,<sup>31</sup> E. Chudakov,<sup>36</sup> E. Cisbani,<sup>17</sup> B. Craver,<sup>17</sup> F. Cusanno,<sup>17,\*</sup> M. M. Dalton,<sup>43</sup> R. De Leo,<sup>14</sup> K. de Jager,<sup>36,43</sup> W. Deconinck,<sup>25,7</sup> P. Decowski,<sup>32</sup> D. Deepa,<sup>28</sup> X. Deng,<sup>43</sup> A. Deur,<sup>36</sup> D. Dutta,<sup>26</sup> A. Etile,<sup>6</sup> C. Ferdi,<sup>6</sup> R. J. Feuerbach,<sup>36</sup> J. M. Finn,<sup>7,†</sup> D. Flay,<sup>35</sup> G. B. Franklin,<sup>4</sup> M. Friend,<sup>4</sup> S. Frullani,<sup>17</sup> E. Fuchey,<sup>6,35</sup> S. A. Fuchs,<sup>7</sup> K. Fuoti,<sup>40</sup> F. Garibaldi,<sup>17</sup> E. Gasser,<sup>6</sup> R. Gilman,<sup>30,36</sup> A. Giusa,<sup>15</sup> A. Glamazdin,<sup>20</sup> L. E. Glesener,<sup>7</sup> J. Gomez,<sup>36</sup> M. Gorchtein,<sup>18,46</sup> J. Grames,<sup>36</sup> K. Grimm,<sup>7</sup> C. Gu,<sup>43</sup> O. Hansen,<sup>36</sup> J. Hansknecht,<sup>36</sup> O. Hen,<sup>34</sup> D. W. Higinbotham,<sup>36</sup> R. S. Holmes,<sup>33</sup> T. Holmstrom,<sup>23</sup> C. J. Horowitz,<sup>18</sup> J. Hoskins,<sup>7</sup> J. Huang,<sup>25,24</sup> T. B. Humensky,<sup>37</sup> C. E. Hyde,<sup>28,6</sup> H. Ibrahim,<sup>28,2</sup> F. Itard,<sup>6</sup> C.-M. Jen,<sup>33</sup> E. Jensen,<sup>7</sup> X. Jiang,<sup>30,24</sup> G. Jin,<sup>43</sup> S. Johnston,<sup>40</sup> J. Katich,<sup>7</sup> L. J. Kaufman,<sup>40,‡</sup> A. Kelleher,<sup>25</sup> K. Kliakhandler,<sup>34</sup> P. M. King,<sup>27</sup> A. Kolarkar,<sup>38</sup> S. Kowalski,<sup>25</sup> E. Kuchina,<sup>30</sup> K. S. Kumar,<sup>40</sup> L. Lagamba,<sup>14</sup> D. Lambert,<sup>32</sup> P. LaViolette,<sup>40</sup> J. Leacock,<sup>44</sup> J. Leckey IV,<sup>7</sup> J. H. Lee,<sup>7,27,§</sup> J. J. LeRose,<sup>36</sup> D. Lhuillier,<sup>5</sup> R. Lindgren,<sup>43</sup> N. Liyanage,<sup>43</sup> N. Lubinsky,<sup>29</sup> J. Mammei,<sup>40</sup> F. Mammoliti,<sup>15</sup> D. J. Margaziotis,<sup>3</sup> P. Markowitz,<sup>10</sup> M. Mazouz,<sup>21</sup> K. McCormick,<sup>30</sup> A. McCreary,<sup>36</sup> D. McNulty,<sup>40</sup> D. G. Meekins,<sup>36</sup> L. Mercado,<sup>40</sup> Z.-E. Meziani,<sup>35</sup> R. W. Michaels,<sup>36</sup> M. Mihovilovic,<sup>19</sup> B. Moffit,<sup>7</sup> P. Monaghan,<sup>12</sup> N. Muangma,<sup>25</sup> C. Muñoz-Camacho,<sup>6</sup> S. Nanda,<sup>36</sup> V. Nelyubin,<sup>43</sup> D. Neyret,<sup>5</sup> Nuruzzaman,<sup>26</sup> Y. Oh,<sup>31</sup> K. Otis,<sup>40</sup> A. Palmer,<sup>23</sup> D. Parno,<sup>4</sup> K. D. Paschke,<sup>43</sup> S. K. Phillips,<sup>41,||</sup> M. Poelker,<sup>36</sup> R. Pomatsalyuk,<sup>20</sup> M. Posik,<sup>35</sup> M. Potokar,<sup>19</sup> K. Prok,<sup>5</sup> A. J. R. Puckett,<sup>25,24</sup> X. Qian,<sup>9</sup> Y. Qiang,<sup>25,¶</sup> B. Quinn,<sup>4</sup> A. Rakhman,<sup>33</sup> P. E. Reimer,<sup>1</sup> B. Reitz,<sup>36</sup> S. Riordan,<sup>40,43</sup> J. Roche,<sup>27,36</sup> P. Rogan,<sup>40</sup> G. Ron,<sup>22</sup> G. Russo,<sup>15</sup> K. Saenboonruang,<sup>43</sup> A. Saha,<sup>36</sup> B. Sawatzky,<sup>36</sup> A. Shahinyan,<sup>45,36</sup> R. Silwal,<sup>43</sup> J. Singh,<sup>43</sup> S. Sirca,<sup>19</sup> K. Slifer,<sup>41</sup> R. Snyder,<sup>43</sup> P. Solvignon,<sup>36</sup> P. A. Souder,<sup>33,\*\*</sup> M. L. Sperduto,<sup>15</sup> R. Subedi,<sup>43</sup> M. L. Stutzman,<sup>36</sup> R. Suleiman,<sup>36</sup> V. Sulkosky,<sup>25,7</sup> C. M. Sutura,<sup>15</sup> W. A. Tobias,<sup>43</sup> W. Troth,<sup>23</sup> G. M. Urciuoli,<sup>16</sup> P. Ulmer,<sup>28</sup> A. Vacheret,<sup>5</sup> E. Voutier,<sup>21</sup> B. Waidyawansa,<sup>27</sup> D. Wang,<sup>43</sup> K. Wang,<sup>43</sup> J. Wexler,<sup>40</sup> A. Whitbeck,<sup>38</sup> R. Wilson,<sup>13</sup> B. Wojtsekhowski,<sup>36</sup> X. Yan,<sup>42</sup> H. Yao,<sup>35</sup> Y. Ye,<sup>42</sup> Z. Ye,<sup>12,43</sup> V. Yim,<sup>40</sup> L. Zana,<sup>33</sup> X. Zhan,<sup>1</sup> J. Zhang,<sup>36</sup> Y. Zhang,<sup>30</sup> X. Zheng,<sup>43</sup> V. Ziskin,<sup>25</sup> and P. Zhu<sup>42</sup>

(HAPPEX and PREX Collaborations)

<sup>1</sup>Argonne National Laboratory, Argonne, Illinois 60439, USA

<sup>2</sup>Cairo University, Giza 12613, Egypt

<sup>3</sup>California State University, Los Angeles, Los Angeles, California 90032, USA

<sup>4</sup>Carnegie Mellon University, Pittsburgh, Pennsylvania 15213, USA

<sup>5</sup>CEA Saclay, DAPNIA/SPhN, F-91191 Gif-sur-Yvette, France

<sup>6</sup>Clermont Université, Université Blaise Pascal, CNRS/IN2P3, Laboratoire de Physique  
Corpusculaire, FR-63000 Clermont-Ferrand, France

<sup>7</sup>College of William and Mary, Williamsburg, Virginia 23187, USA

<sup>8</sup>Christopher Newport University, Newport News, Virginia 23606, USA

<sup>9</sup>Duke University, TUNL, Durham, North Carolina, 27706, USA

<sup>10</sup>Florida International University, Miami, Florida 33199, USA

<sup>11</sup>The George Washington University, Washington, D.C. 20052, USA

<sup>12</sup>Hampton University, Hampton, Virginia 23668, USA

<sup>13</sup>Harvard University, Cambridge, Massachusetts 02138, USA

<sup>14</sup>INFN, Sezione di Bari and University of Bari, I-70126 Bari, Italy

<sup>15</sup>INFN, Sezione di Catania and University of Catania, I-95123 Catania, Italy

<sup>16</sup>INFN, Sezione di Roma, I-00161 Rome, Italy

<sup>17</sup>INFN, Sezione di Roma, gruppo Sanità, I-00161 Rome, Italy

<sup>18</sup>Indiana University, Bloomington, Indiana 47405, USA

<sup>19</sup>Institut Jožef Stefan, 3000 SI-1001 Ljubljana, Slovenia

<sup>20</sup>Kharkov Institute of Physics and Technology, Kharkov 61108, Ukraine

<sup>21</sup>LPSC, Université Joseph Fourier, CNRS/IN2P3, INPG, Grenoble, France

<sup>22</sup>Lawrence Berkeley National Laboratory, Berkeley, California 94720, USA

<sup>23</sup>Longwood University, Farmville, Virginia 23909, USA

- <sup>24</sup>Los Alamos National Laboratory, Los Alamos, New Mexico 87545, USA  
<sup>25</sup>Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA  
<sup>26</sup>Mississippi State University, Mississippi State, Mississippi 39762, USA  
<sup>27</sup>Ohio University, Athens, Ohio 45701, USA  
<sup>28</sup>Old Dominion University, Norfolk, Virginia 23529, USA  
<sup>29</sup>Rensselaer Polytechnic Institute, Troy, New York 12180, USA  
<sup>30</sup>Rutgers University, The State University of New Jersey, New Brunswick, New Jersey 08901, USA  
<sup>31</sup>Seoul National University, Seoul 151-742, South Korea  
<sup>32</sup>Smith College, Northampton, Massachusetts 01063, USA  
<sup>33</sup>Syracuse University, Syracuse, New York 13244, USA  
<sup>34</sup>Tel Aviv University, P.O. Box 39040, Tel-Aviv 69978, Israel  
<sup>35</sup>Temple University, Philadelphia, Pennsylvania 19122, USA  
<sup>36</sup>Thomas Jefferson National Accelerator Facility, Newport News, Virginia 23606, USA  
<sup>37</sup>University of Chicago, Chicago, Illinois, 60637, USA  
<sup>38</sup>University of Kentucky, Lexington, Kentucky 40506, USA  
<sup>39</sup>University of Maryland, College Park, Maryland, 20742, USA  
<sup>40</sup>University of Massachusetts Amherst, Amherst, Massachusetts 01003, USA  
<sup>41</sup>University of New Hampshire, Durham, New Hampshire 03824, USA  
<sup>42</sup>University of Science and Technology of China, Hefei, Anhui 230026, People's Republic of China  
<sup>43</sup>University of Virginia, Charlottesville, Virginia 22903, USA  
<sup>44</sup>Virginia Polytechnic Institute and State University, Blacksburg, Virginia 24061, USA  
<sup>45</sup>Yerevan Physics Institute, Yerevan, Armenia  
<sup>46</sup>Institute für Kernphysik, Universität Mainz, 55128 Mainz, Germany
- (Received 30 August 2012; published 5 November 2012)

We have measured the beam-normal single-spin asymmetry  $A_n$  in the elastic scattering of 1–3 GeV transversely polarized electrons from  $^1\text{H}$  and for the first time from  $^4\text{He}$ ,  $^{12}\text{C}$ , and  $^{208}\text{Pb}$ . For  $^1\text{H}$ ,  $^4\text{He}$ , and  $^{12}\text{C}$ , the measurements are in agreement with calculations that relate  $A_n$  to the imaginary part of the two-photon exchange amplitude including inelastic intermediate states. Surprisingly, the  $^{208}\text{Pb}$  result is significantly smaller than the corresponding prediction using the same formalism. These results suggest that a systematic set of new  $A_n$  measurements might emerge as a new and sensitive probe of the structure of heavy nuclei.

DOI: [10.1103/PhysRevLett.109.192501](https://doi.org/10.1103/PhysRevLett.109.192501)

PACS numbers: 25.30.Bf, 27.10.+h, 27.80.+w

Traditionally, fixed-target electron scattering has been analyzed in terms of the one-boson (photon or  $Z$ ) exchange approximation. For scattering off heavy nuclei, distorted waves, based on solutions to the Dirac equation in the strong electric field of the nucleus, are also required to describe the data. Recently, the inclusion of the exchange of one or more additional photons has been necessary for the interpretation of precision data. The electric form factors  $G_E^p$  extracted in elastic electron-proton scattering using two different techniques, Rosenbluth separation and polarization observables, were inconsistent [1–3]. The latter should be less sensitive to higher-order electromagnetic effects, and calculations including two-photon exchange provide a plausible explanation for the difference [4–8]. Another example is corrections to the parity-violating asymmetry  $A_{\text{PV}}$  in the same process, which provides a measurement of the weak charge of the proton and serves as a sensitive test of the electroweak theory. For interpreting  $A_{\text{PV}}$ ,  $\gamma$ - $Z$  box diagrams are important [9], as well as two-photon exchange [10]. Theoretical calculations of two-photon exchange processes are difficult, because an integral over all off-shell proton intermediate state contributions must be made.

The effect of the extra boson is relatively small on the measured cross section or asymmetry for the above examples. On the other hand, the beam-normal spin asymmetry  $A_n$  for elastic electron scattering at GeV energies is dominated by two (or more)  $\gamma$  exchange. Several measurements of  $A_n$  at GeV energies for the proton have been reported [11–14]. Several theoretical papers report computed values of  $A_n$  that are in qualitative agreement with the data when they include the effects of inelastic intermediate hadronic states [15–18].

The beam-normal, or transverse asymmetry,  $A_n$  is a direct probe of higher-order photon exchange as time-reversal symmetry dictates that  $A_n$  is zero at first Born approximation. Afanasev *et al.* [4] and Gorchtein and Horowitz [19] have calculated  $A_n$ , in a two-photon exchange approximation, but including a full range of intermediate excited states. Gorchtein and Horowitz predict that  $A_n$  scales roughly as the ratio of mass number  $A$  to  $Z$  and is not strongly  $Z$ -dependent. In contrast, Cooper and Horowitz [20] calculate Coulomb distortion effects and work to all orders in photon exchanges by numerically solving the Dirac equation. However, they consider only the elastic intermediate state. They find that elastic

intermediate state contributions, while in general small, increase strongly with  $Z$ .

To predict  $A_n$  for nuclear targets, Afanasev used a unitarity-based model [18] with the total photoproduction cross section and the Compton slope as input; his prediction for  ${}^4\text{He}$  is consistent with the value of  $A_n$  reported in this Letter. However, there is not yet a calculation of  $A_n$  that includes both Coulomb distortion effects and a full range of excited intermediate states. Measuring  $A_n$  as a function of  $Z$  might reveal the role of dispersion effects relative to Coulomb distortions and motivate more detailed calculations. To this end, in this Letter, we report data on the beam-normal spin asymmetry  $A_n$  on the targets  ${}^1\text{H}$ ,  ${}^4\text{He}$ ,  ${}^{12}\text{C}$ , and  ${}^{208}\text{Pb}$ .

To observe the beam-normal single-spin asymmetry, the electron beam-spin vector  $\vec{P}_e$  must have a component normal to the scattering plane defined by the unit vector  $\hat{k}$  perpendicular to the plane, where  $\hat{k} = \vec{k}/|k|$ ;  $\vec{k} = \vec{k}_e \times \vec{k}_{\text{out}}$ , where  $\vec{k}_e$  and  $\vec{k}_{\text{out}}$  are, respectively, the incident and scattered electron momenta. The measured beam-normal single-spin asymmetry is then defined as  $A_n^m = (\sigma_{\uparrow} - \sigma_{\downarrow})/(\sigma_{\uparrow} + \sigma_{\downarrow})$ , where  $\sigma_{\uparrow(\downarrow)}$  is the cross section for beam electron spin parallel (antiparallel) to  $\hat{k}$ . The measured asymmetry  $A_n^m$  is related to  $A_n$  by

$$A_n^m = A_n \vec{P}_e \cdot \hat{k}, \quad (1)$$

where  $\phi$  is the angle between  $\hat{k}$  and  $\vec{P}_e$ :  $\cos\phi = \vec{P}_e \cdot \hat{k}/|P_e|$ .

The measurements were carried out in Hall A at the Thomas Jefferson National Accelerator Facility. The data were obtained as a part of a study of systematic uncertainties for three experiments designed to measure  $A_{\text{PV}}$  in elastic electron scattering, since  $A_n$  can contribute to the extracted  $A_{\text{PV}}$  if the beam polarization has a transverse component and the apparatus lacks perfect symmetry.

The data were obtained in 2004 for the  ${}^1\text{H}$  and  ${}^4\text{He}$  targets where the primary goal was to measure  $A_{\text{PV}}$  in order to determine the strange form factors in the nucleon [21,22]. The  ${}^{12}\text{C}$  and  ${}^{208}\text{Pb}$  data were obtained in 2010 where the goal was to determine the radius of the distribution of neutrons [23,24]. The kinematics for each target is given in Table I: the central acceptance angle of the spectrometers  $\theta$ , the beam energy  $E_b$ , the acceptance-averaged 4-momentum transfer  $Q^2$ , and the average accepted  $\cos\phi$  [Eq. (1)]. The uncertainties in  $Q^2$  were 1% for the  ${}^1\text{H}$  and  ${}^4\text{He}$  data and 1.3% for the  ${}^{12}\text{C}$  and  ${}^{208}\text{Pb}$  data [21–23].

All of the targets except  ${}^{12}\text{C}$  were cooled with helium gas at about 20 K. The  $\text{LH}_2$  and high pressure He targets featured rapid vertical flow of the fluid. In addition, the beam was rastered over a 4 mm  $\times$  4 mm square for all targets. The 0.55 mm thick isotopically pure  ${}^{208}\text{Pb}$  target was sandwiched between two 150  $\mu\text{m}$  diamond foils, and the edges were cooled with the cold helium. Electrons elastically scattered from the targets were focused onto detectors in the focal plane of the Hall A high resolution

TABLE I. Kinematic values for the various targets.

Target	H	${}^4\text{He}$	${}^{12}\text{C}$	${}^{208}\text{Pb}$
$\theta$	6°	6°	5°	5°
$Q^2(\text{GeV}^2)$	0.0989	0.0773	0.009 84	0.008 81
$E_b(\text{GeV})$	3.026	2.750	1.063	1.063
$\langle\cos\phi\rangle$	0.968	0.967	0.963	0.967

spectrometers [25]. The transverse asymmetry is modulated by the sine of the azimuthal electron scattering angle, and so the electron polarization was set vertical. This ensured that the acceptance of the two spectrometers, which are symmetrically placed to accept horizontally scattered events, contained the maximum and minimum of the asymmetry. The momentum resolution of the spectrometers ensured that essentially only elastic events were accepted.

To measure the asymmetry, Cherenkov light was produced in a radiator and collected by a photomultiplier tube (PMT), whose output sent to an analog-to-digital converter and integrated over a fixed time period of constant helicity. These detectors had to withstand the radiation damage caused by the high signal flux and also provide a uniform response to the electrons so that integrating the signals did not increase fluctuations. For the  ${}^{208}\text{Pb}$  and  ${}^{12}\text{C}$  data, each spectrometer had two 3.5 cm by 14 cm quartz detectors oriented at 45° to the direction of the electrons in the spectrometer, one in front that was 5 mm thick and one behind that was 1 cm thick. For the  ${}^1\text{H}$  and  ${}^4\text{He}$  data, a five-layer sandwich of quartz and brass provided sufficient energy resolution.

The electron beam originated from a GaAs photocathode illuminated by circularly polarized light [26]. By reversing the sign of the laser circular polarization, the direction of the spin at the target could be reversed rapidly [27]. A half-wave ( $\lambda/2$ ) plate was periodically inserted into the laser optical path which passively reversed the sign of the electron beam polarization. Roughly equal statistics were thus accumulated with opposite signs for the measured asymmetry, which suppressed many systematic effects. The direction of the polarization could be controlled by a Wien filter and solenoidal lenses near the injector [28]. The accelerated beam was directed into Hall A, where its intensity, energy, and trajectory on target were inferred from the response of several monitoring devices.

Each period of constant spin direction is referred to as a “window.” The beam monitors, target, detector components, and electronics were designed so that the fluctuations in the fractional difference in the PMT response between a pair of successive windows were dominated by scattered electron counting statistics. To keep spurious beam-induced asymmetries under control at well below the parts per million level, careful attention was given to the design and configuration of the laser optics leading to the photocathode [27].

The spin-reversal rate was 30 Hz for the  $^1\text{H}$  and  $^4\text{He}$  data and 240 Hz for the  $^{12}\text{C}$  and  $^{208}\text{Pb}$  data. The integrated response of each detector PMT and beam monitor was digitized and recorded for each window. In the 30 Hz case, the raw spin-direction asymmetry  $A_{\text{raw}}$  in each spectrometer arm was computed from the detector response normalized to the beam intensity for each window pair. At the faster reversal, quadruplets of windows with either of the patterns  $+- -+$  or  $-+ + -$  were used to suppress the significant 60 Hz line noise. In either case, the sequence of these patterns was chosen with a pseudorandom number generator.

Loose requirements were imposed on beam quality, removing periods of beam intensity, position, or energy instability, removing about 25% of the total data sample. No spin-direction-dependent cuts were applied. Since we measure the difference between two horizontal detectors, the dominant source of noise due to the beam arose from position fluctuations in the horizontal direction, which change the acceptance of the spectrometers in opposite directions. Noise in the beam energy or current largely cancels. In contrast, a measurement of the sum of the detectors in the  $A_{\text{PV}}$  case is relatively more sensitive to beam energy or current fluctuations and less to the beam position.

As explained in detail in Refs. [21–23], the window-to-window differences in the asymmetry from beam jitter were reduced by using the correlations to beam position differences from precision beam position monitors,  $\Delta x_i$ , by defining a correction  $A_{\text{beam}} = \sum c_i \Delta x_i$ . The  $c_i$  were measured several times each hour from calibration data where the beam was modulated by using steering coils and an accelerating cavity. The largest  $c_i$  was for  $^{208}\text{Pb}$  and was on

the order of 50 ppb/nm. The spread in the resulting  $A_n^m = A_{\text{raw}} - A_{\text{beam}}$  was observed to be dominated by counting statistics. For example, for  $^{208}\text{Pb}$ , which had the highest rate and hence the smallest statistical uncertainty for a window, this spread corresponded to a rate of about 1 GHz at a beam current of  $70 \mu\text{A}$ . About 1 d was spent at each  $\lambda/2$  setting on each target.

The values of  $A_n^m$  were consistent from run to run as shown in Fig. 1. The asymmetries in each spectrometer arm were of opposite sign as expected [ $\hat{k}$  in Eq. (1) reverses sign]. After correcting for the  $\lambda/2$  reversals, the magnitudes of  $A_n^m$  are consistent within statistical uncertainties. The reduced  $\chi^2$  for a constant fit to the  $A_n^m$  runs is close to 1 for each target type. The average  $A_{\text{beam}}$  corrections were negligible. The physics asymmetry  $A_n$  is calculated from  $A_n^m$  by correcting for the beam polarization  $P_e$ , the average value of  $\cos\phi$  as given in Table I, and the background subtractions from the Al windows in the  $\text{LH}_2$  and  $^4\text{He}$  targets and the diamond surrounding the lead foil.

Nonlinearity in the PMT response was limited to 1% in bench tests that mimicked running conditions. The total relative nonlinearity between the PMT response and those of the beam intensity monitors was limited to 1.5% by studies. An acceptance correction accounted for the nonlinear dependence of the asymmetry with  $Q^2$ . A significant systematic uncertainty in  $\langle Q^2 \rangle$  is in the determination of the absolute scale of  $\theta_{\text{lab}}$ . A nuclear recoil technique with a dedicated calibration run using a water cell target [22] was used to set a scale uncertainty on  $\langle Q^2 \rangle$  of  $<0.2\%$ .

Beam polarization measurements [ $P_e$  in Eq. (1)] were made during the runs for the four nuclei. The beam polarization was inferred from longitudinal polarization




FIG. 1. Plots of the asymmetries for carbon and lead. The top row shows for the  $^{12}\text{C}$  target the left high resolution spectrometer (HRS) and right HRS asymmetries in the left and right panels, respectively. The bottom row shows the same sequence for the  $^{208}\text{Pb}$  target. The data have been sign-corrected for the  $\lambda/2$  plate insertions.



TABLE II.  $A_{PV}$  uncertainty contributions in units of  $10^{-6}$  or parts per million.

Target	$^1\text{H}$	$^4\text{He}$	$^{12}\text{C}$	$^{208}\text{Pb}$
False asymmetry	0.14	0.11	0.02	0.12
Beam polarization	0.21	0.33	0.08	0.003
Linearity	0.07	0.15	0.06	0.004
Target windows	0.06	0.12	0.00	0.062
Total systematic	0.27	0.41	0.10	0.14
Statistical	1.52	1.39	0.36	0.21

measurements taken before and after the transverse polarization data taking. A solenoid was used to control the orientation of the polarization between the longitudinal and transverse (vertical) directions. The polarization was verified to be purely vertical to within  $\pm 2^\circ$  with a Mott polarimeter located in an injector 5 MeV extraction line. The vertical component of the polarization set at the injector is conserved after passing through the Continuous Electron Beam Accelerator Facility accelerator, a result of accelerating and transporting the polarized beam in planes flat with respect to one another. The extent to which the beam-spin tune degrades the vertical polarization orientation in the Continuous Electron Beam Accelerator Facility has been studied and determined to be  $\pm 1^\circ$  [29]. A small longitudinal component of the electron spin introduces a negligible parity-violating contribution to the measured asymmetry.

For  $^{12}\text{C}$  and  $^{208}\text{Pb}$ , the longitudinal polarization measurements included data taken with a Compton polarimeter, yielding  $P_e = 0.8820 \pm 0.012 \pm 0.012$ . An independent Møller polarimeter gave  $P_e = 0.9049 \pm 0.001 \pm 0.011$  for  $^{12}\text{C}$  and  $^{208}\text{Pb}$ . We used the average of these two measurements. For the  $^1\text{H}$  and  $^4\text{He}$  data, only the Møller polarimeter was used. For  $^1\text{H}$  data  $P_e = 75.1 \pm 1.7\%$ , and for the  $^4\text{He}$  data  $P_e = 84.2 \pm 1.7\%$ .

A summary of the systematic and statistical uncertainties is shown in Table II. The central values of  $A_n$  for each nucleus and the total combined statistical and systematic uncertainties added in quadrature are displayed in the first two rows of Table III. For  $^1\text{H}$ , our result is consistent with a previously reported measurement [12] for the same  $Q^2$  but at a lower beam energy (0.85 GeV).

TABLE III. The measured  $A_n$  and derived  $\hat{A}_n$  values [Eq. (2)] for the four nuclei along with the corresponding total uncertainties  $A/Z$  and  $Q$ .

Target	H	$^4\text{He}$	$^{12}\text{C}$	$^{208}\text{Pb}$
$A_n$ (ppm)	-6.80	-13.97	-6.49	0.28
$\sigma(A_n)$ (ppm)	$\pm 1.54$	$\pm 1.45$	$\pm 0.38$	$\pm 0.25$
$\sqrt{Q^2}$ (GeV)	0.31	0.28	0.099	0.094
$A/Z$	1.0	2.0	2.0	2.53
$\hat{A}_n$ (ppm/GeV)	-21.9	-24.9	-32.8	+1.2
$\sigma(\hat{A}_n)$ (ppm/GeV)	$\pm 5.0$	$\pm 2.6$	$\pm 1.9$	$\pm 1.1$

We now discuss the observed trends of the first ever measurements of  $A_n$  for target nuclei with  $A > 2$ . In our kinematic range, the calculations in Ref. [19] scale approximately with  $Z$ ,  $A$ , and  $\sqrt{Q^2}$  as

$$A_n = \hat{A}_n \frac{QA}{Z}, \quad (2)$$

where  $\hat{A}_n$  is approximately constant, with a small additional dependence on the incident beam energy  $E_{\text{beam}}$ :  $\sim -25$  ppm/GeV for  $E_{\text{beam}} \sim 3$  GeV (for  $^1\text{H}$  and  $^4\text{He}$ ) and  $\sim -30$  ppm/GeV for  $E_{\text{beam}} \sim 1$  GeV (for  $^{12}\text{C}$  and  $^{208}\text{Pb}$ ). In the last two rows of Table III, we see that the extracted  $\hat{A}_n$  from the three lower  $Z$  nuclei are consistent with this empirical trend, while the  $^{208}\text{Pb}$  result is consistent with zero. The  $^{208}\text{Pb}$  result is in strong disagreement with the theoretical prediction as shown in Fig. 2, which plots the measurement results and their predictions [19].

Motivated by this large observed disagreement, we initiate a discussion of the potential dynamics by first noting that the scattering angle for all four measurements was roughly the same (Table I). If dispersion corrections play a bigger role than predicted, one might expect larger disagreements for  $A_n$  measurements taken at lower beam energy, as is the case for  $^{12}\text{C}$  and  $^{208}\text{Pb}$ . However, the measured  $A_n$  for  $^{12}\text{C}$  is quite consistent with theoretical expectations. In Fig. 3, we plot the fractional difference between the measured values and the predictions of Ref. [19] as a function of  $Z$ . The trend suggests that Coulomb distortions are playing a very significant role at large  $Z$ , underscoring the potential interest in additional  $A_n$  measurements with intermediate  $Z$  nuclei.

In conclusion, we have measured the beam-normal single-spin asymmetry  $A_n$  for  $^1\text{H}$ ,  $^4\text{He}$ ,  $^{12}\text{C}$ , and  $^{208}\text{Pb}$  and find good agreement for  $^1\text{H}$ ,  $^4\text{He}$ , and  $^{12}\text{C}$  with the calculations in Ref. [19], which include a dispersion integral over intermediate excited states. However, they




FIG. 2 (color). Extracted physics asymmetries  $A_n$  vs  $Q$ . Each curve, specific to a particular nucleus as indicated, is a theoretical calculation from Ref. [19].




FIG. 3 (color). Percent fractional deviation of  $A_n$  measurements normalized to the respective theory prediction vs target nucleus  $Z$ .

are only to order  $\alpha^2$  (two-photon exchange) and neglect Coulomb distortions. On the other hand,  $A_n$  for  $^{208}\text{Pb}$  is measured to be very small and disagrees completely with theoretical calculations. Coulomb distortions were shown in Ref. [20] to grow rapidly with  $Z$ . On the other hand, the weight of dispersion corrections varies with the incident beam energy. Thus, new theoretical calculations that treat dispersion corrections and Coulomb distortions simultaneously as well as a systematic set of  $A_n$  measurements for a range of  $Z$  at various beam energies might lead to new insights into the structure of heavy nuclei.

We thank the entire staff of JLab for their efforts to develop and maintain the polarized beam and the experimental apparatus. This work was supported by DOE Contract No. DE-AC05-84ER40150 Modification No. M175, under which the Southeastern Universities Research Association (SURA) operates JLab, Contract No. DE-AC02-06CH11357 for Argonne National Lab, and by the Department of Energy, the National Science Foundation, the INFN (Italy), and the Commissariat à l'Énergie Atomique (France).

\*Present address: Technische Universitaet Muenchen, Excellence Cluster Universe, Garching b. Muenchen, Germany.

†Deceased.

‡Present address: Indiana University, Bloomington, IN 47405, USA.

§Present address: Institute for Basic Science, Daejeon, South Korea.

||Present address: University of New Hampshire, Durham, NH 03824, USA.

¶Present address: Thomas Jefferson National Accelerator Facility, Newport News, VA 23606, USA.

\*\*souder@physics.syr.edu

- [1] C. F. Perdrisat, V. Punjabi, and M. Vanderhaeghen, *Prog. Part. Nucl. Phys.* **59**, 694 (2007).
- [2] J. Arrington, C. D. Roberts, and J. M. Zanotti, *J. Phys. G* **34**, S23 (2007).
- [3] J. Arrington, *Phys. Rev. C* **68**, 034325 (2003).
- [4] A. V. Afanasev, S. J. Brodsky, C. E. Carlson, Y.-C. Chen, and M. Vanderhaeghen, *Phys. Rev. D* **72**, 013008 (2005).
- [5] C. E. Carlson and M. Vanderhaeghen, *Annu. Rev. Nucl. Part. Sci.* **57**, 171 (2007).
- [6] J. Arrington, W. Melnitchouk, and J. A. Tjon, *Phys. Rev. C* **76**, 035205 (2007).
- [7] P. G. Blunden, W. Melnitchouk, and J. A. Tjon, *Phys. Rev. C* **72**, 034612 (2005).
- [8] J. Arrington, P. G. Blunden, and W. Melnitchouk, *Prog. Part. Nucl. Phys.* **66**, 782 (2011).
- [9] M. Gorchtein, C. J. Horowitz, and M. J. Ramsey-Musolf, *Phys. Rev. C* **84**, 015502 (2011).
- [10] A. V. Afanasev and C. E. Carlson, *Phys. Rev. Lett.* **94**, 212301 (2005).
- [11] S. P. Wells *et al.*, *Phys. Rev. C* **63**, 064001 (2001).
- [12] F. E. Maas *et al.*, *Phys. Rev. Lett.* **94**, 082001 (2005).
- [13] D. S. Armstrong *et al.*, *Phys. Rev. Lett.* **99**, 092301 (2007).
- [14] D. Androic *et al.*, *Phys. Rev. Lett.* **107**, 022501 (2011).
- [15] A. V. Afanasev and N. P. Merenkov, *Phys. Lett. B* **599**, 48 (2004).
- [16] B. Pasquini and M. Vanderhaeghen, *Eur. Phys. J. A* **24S2**, 29 (2005).
- [17] B. Pasquini and M. Vanderhaeghen, *Phys. Rev. C* **70**, 045206 (2004).
- [18] A. V. Afanasev, [arXiv:0711.3065](https://arxiv.org/abs/0711.3065).
- [19] M. Gorchtein and C. J. Horowitz, *Phys. Rev. C* **77**, 044606 (2008).
- [20] E. D. Cooper and C. J. Horowitz, *Phys. Rev. C* **72**, 034602 (2005).
- [21] A. Acha *et al.*, *Phys. Rev. Lett.* **98**, 032301 (2007).
- [22] K. A. Aniol *et al.*, *Phys. Rev. Lett.* **96**, 022003 (2006).
- [23] S. Abrahamyan *et al.* (PREX Collaboration), *Phys. Rev. Lett.* **108**, 112502 (2012).
- [24] C. J. Horowitz, S. J. Pollock, P. A. Souder and R. Michaels, *Phys. Rev. C* **63**, 025501 (2001).
- [25] J. Alcorn *et al.*, *Nucl. Instrum. Methods Phys. Res., Sect. A* **522**, 294 (2004).
- [26] C. K. Sinclair, P. Adderley, B. Dunham, J. Hansknecht, P. Hartmann, M. Poelker, J. Price, P. Rutt, W. Schneider, and M. Steigerwald, *Phys. Rev. ST Accel. Beams* **10**, 023501 (2007); P. A. Adderley, J. Clark, J. Grames, J. Hansknecht, K. Surlis-Law, D. Machie, M. Poelker, M. L. Stutzman, and R. Suleiman, *Phys. Rev. ST Accel. Beams* **13**, 010101 (2010).
- [27] K. D. Paschke, *Eur. Phys. J. A* **32**, 549 (2007).
- [28] J. Grames *et al.*, in *Proceedings of the 2011 Particle Accelerator Conference, New York, 2011* (IEEE, New York, 2011).
- [29] J. M. Grames, Ph.D. thesis, Jefferson Laboratory [Institution Report No. JLAB-R-00-001, 2000].