



HAL
open science

Measurement of the neutrino-oxygen neutral-current interaction cross section by observing nuclear de-excitation γ -rays

K. Abe, J. Adam, H. Aihara, T. Akiri, C. Andreopoulos, S. Aoki, A. Ariga, T. Ariga, S. Assylbekov, D. Autiero, et al.

► To cite this version:

K. Abe, J. Adam, H. Aihara, T. Akiri, C. Andreopoulos, et al.. Measurement of the neutrino-oxygen neutral-current interaction cross section by observing nuclear de-excitation γ -rays. *Physical Review D*, 2014, 90 (7), pp.072012. 10.1103/PhysRevD.90.072012 . in2p3-00962301

HAL Id: in2p3-00962301

<https://hal.in2p3.fr/in2p3-00962301>

Submitted on 14 Nov 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Measurement of the neutrino-oxygen neutral-current interaction cross section by observing nuclear deexcitation γ rays

K. Abe,⁴⁶ J. Adam,³² H. Aihara,^{45,23} T. Akiri,⁹ C. Andreopoulos,⁴⁴ S. Aoki,²⁴ A. Ariga,² T. Ariga,² S. Assylbekov,⁸ D. Autiero,²⁹ M. Barbi,³⁹ G. J. Barker,⁵⁴ G. Barr,³⁵ M. Bass,⁸ M. Batkiewicz,¹³ F. Bay,¹¹ S. W. Bentham,²⁶ V. Berardi,¹⁵ B. E. Berger,^{8,23} S. Berkman,⁴ I. Bertram,²⁶ S. Bhadra,⁵⁸ F. d. M. Blaszczyk,²⁸ A. Blondel,¹² C. Bojecho,⁵¹ S. Bordini,¹⁵ S. B. Boyd,⁵⁴ D. Brailsford,¹⁷ A. Bravar,¹² C. Bronner,²³ N. Buchanan,⁸ R. G. Calland,²⁷ J. Caravaca Rodríguez,¹⁵ S. L. Cartwright,⁴² R. Castillo,¹⁵ M. G. Catanesi,¹⁸ A. Cervera,¹⁶ D. Cherdack,⁸ G. Christodoulou,²⁷ A. Clifton,⁸ J. Coleman,²⁷ S. J. Coleman,⁷ G. Collazuol,²⁰ K. Connolly,⁵⁵ L. Cremonesi,³⁸ A. Dabrowska,¹³ I. Danko,³⁷ R. Das,⁸ S. Davis,⁵⁵ P. de Perio,⁴⁹ G. De Rosa,¹⁹ T. Dealtry,^{44,35} S. R. Dennis,^{54,44} C. Densham,⁴⁴ D. Dewhurst,³⁵ F. Di Lodovico,³⁸ S. Di Luise,¹¹ O. Drapier,¹⁰ T. Duboyski,³⁸ K. Duffy,³⁵ F. Dufour,¹² J. Dumarchez,³⁶ S. Dytman,³⁷ M. Dziewiecki,⁵³ S. Emery-Schrenk,⁶ A. Ereditato,² L. Escudero,¹⁶ A. J. Finch,²⁶ G. A. Fiorentini Aguirre,⁵⁸ M. Friend,^{14,†} Y. Fujii,^{14,†} Y. Fukuda,³⁰ A. P. Furmanski,⁵⁴ V. Galymov,²⁹ A. Gaudin,⁵¹ S. Giffin,³⁹ C. Giganti,³⁶ K. Gilje,³² D. Goeldi,² T. Golan,⁵⁷ J. J. Gomez-Cadenas,¹⁶ M. Gonin,¹⁰ N. Grant,²⁶ D. Gudin,²² D. R. Hadley,⁵⁴ L. Haegel,¹² A. Haesler,¹² M. D. Haigh,⁵⁴ P. Hamilton,¹⁷ D. Hansen,³⁷ T. Hara,²⁴ M. Hartz,^{23,50} T. Hasegawa,^{14,†} N. C. Hastings,³⁹ Y. Hayato,^{46,23} C. Hearty,^{4,‡} R. L. Helmer,⁵⁰ M. Hierholzer,² J. Hignight,³² A. Hillairet,⁵¹ A. Himmel,⁹ T. Hiraki,²⁵ S. Hirota,²⁵ J. Holeczek,⁴³ S. Horikawa,¹¹ K. Huang,²⁵ A. K. Ichikawa,²⁵ K. Ieki,²⁵ M. Ieva,¹⁵ M. Ikeda,⁴⁶ J. Imber,³² J. Insler,²⁸ T. J. Irvine,⁴⁷ T. Ishida,^{14,†} T. Ishii,^{14,†} S. J. Ives,¹⁷ E. Iwai,¹⁴ K. Iwamoto,⁴⁰ K. Iyogi,⁴⁶ A. Izmaylov,^{16,22} A. Jacob,³⁵ B. Jamieson,⁵⁶ R. A. Johnson,⁷ S. Johnson,⁷ J. H. Jo,³² P. Jonsson,¹⁷ C. K. Jung,^{32,§} M. Kabirnezhad,³¹ A. C. Kaboth,¹⁷ T. Kajita,^{47,§} H. Kakuno,⁴⁸ J. Kameda,⁴⁶ Y. Kanazawa,⁴⁵ D. Karlen,^{51,50} I. Karpikov,²² T. Katori,³⁸ E. Kearns,^{3,23,§} M. Khabibullin,²² A. Khotjantsev,²² D. Kielczewska,⁵² T. Kikawa,²⁵ A. Kilinski,³¹ J. Kim,⁴ S. King,³⁸ J. Kisiel,⁴³ P. Kitching,¹ T. Kobayashi,^{14,†} L. Koch,⁴¹ A. Kolaceke,³⁹ A. Konaka,⁵⁰ L. L. Kormos,²⁶ A. Korzenev,¹² K. Koseki,^{14,†} Y. Koshio,^{33,§} I. Kreslo,² W. Kropp,⁵ H. Kubo,²⁵ Y. Kudenko,^{22,||} S. Kumaratunga,⁵⁰ R. Kurjata,⁵³ T. Kutter,²⁸ J. Lagoda,³¹ K. Laihem,⁴¹ I. Lamont,²⁶ E. Larkin,⁵⁴ M. Laveder,²⁰ M. Lawe,⁴² M. Lazos,²⁷ K. P. Lee,⁴⁷ C. Licciardi,³⁹ T. Lindner,⁵⁰ C. Lister,⁵⁴ R. P. Litchfield,⁵⁴ A. Longhin,²⁰ L. Ludovici,²¹ M. Macaire,⁶ L. Magaletti,¹⁸ K. Mahn,⁵⁰ M. Malek,¹⁷ S. Manly,⁴⁰ A. D. Marino,⁷ J. Marteau,²⁹ J. F. Martin,⁴⁹ S. Martynenko,²² T. Maruyama,^{14,†} J. Marzec,⁵³ E. L. Mathie,³⁹ V. Matveev,²² K. Mavrokoridis,²⁷ E. Mazzucato,⁶ M. McCarthy,⁴ N. McCauley,²⁷ K. S. McFarland,⁴⁰ C. McGrew,³² A. Mefodiev,²² C. Metelko,²⁷ M. Mezzetto,²⁰ P. Mijakowski,³¹ C. A. Miller,⁵⁰ A. Minamino,²⁵ O. Mineev,²² S. Mine,⁵ A. Missert,⁷ M. Miura,^{46,§} L. Monfregola,¹⁶ S. Moriyama,^{46,§} Th. A. Mueller,¹⁰ A. Murakami,²⁵ M. Murdoch,²⁷ S. Murphy,¹¹ J. Myslik,^{25,23} T. Nagasaki,^{14,†} T. Nakadaira,^{14,†} M. Nakahata,^{46,23} T. Nakai,³⁴ K. Nakamura,^{23,14,†} S. Nakayama,^{46,§} T. Nakaya,^{25,23} K. Nakayoshi,^{14,†} C. Nantais,⁴ D. Naples,³⁷ C. Nielsen,⁴ M. Nirkko,² K. Nishikawa,^{14,†} Y. Nishimura,⁴⁷ J. Nowak,²⁶ H. M. O'Keeffe,²⁶ R. Ohta,^{14,†} K. Okumura,^{47,23} T. Okusawa,³⁴ W. Oryszczak,⁵² S. M. Oser,⁴ T. Ovsyannikova,²² R. A. Owen,³⁸ Y. Oyama,^{14,†} V. Palladino,¹⁹ J. L. Palomino,³² V. Paolone,³⁷ D. Payne,²⁷ G. F. Pearce,⁴⁴ O. Perevozchikov,²⁸ J. D. Perkin,⁴² Y. Petrov,⁴ L. Pickard,⁴² E. S. Pinzon Guerra,⁵⁸ C. Pistillo,² P. Plonski,⁵³ E. Poplawska,³⁸ B. Popov,^{36,¶} M. Posiadala-Zezula,⁵² J.-M. Poutissou,⁵⁰ R. Poutissou,⁵⁰ P. Przewlocki,³¹ B. Quilain,¹⁰ E. Radicioni,¹⁸ P. N. Ratoff,²⁶ M. Ravonel,¹² M. A. M. Rayner,¹² A. Redij,² M. Reeves,²⁶ E. Reinherz-Aronis,⁸ C. Riccio,¹⁹ F. Retiere,⁵⁰ A. Robert,³⁶ P. A. Rodrigues,⁴⁰ P. Rojas,⁸ E. Rondio,³¹ S. Roth,⁴¹ A. Rubbia,¹¹ D. Ruterbories,⁴⁰ R. Sacco,³⁸ K. Sakashita,^{14,†} F. Sánchez,¹⁵ F. Sato,¹⁴ E. Scantamburlo,¹² K. Scholberg,^{9,§} S. Schoppmann,⁴¹ J. Schwehr,⁸ M. Scott,⁵⁰ Y. Seiya,³⁴ T. Sekiguchi,^{14,†} H. Sekiya,^{46,23,§} D. Sgalaberna,¹¹ F. Shaker,⁵⁶ M. Shiozawa,^{46,23} S. Short,³⁸ Y. Shustrov,²² P. Sinclair,¹⁷ B. Smith,¹⁷ R. J. Smith,³⁵ M. Smy,⁵ J. T. Sobczyk,⁵⁷ H. Sobel,^{5,23} M. Sorel,¹⁶ L. Southwell,²⁶ P. Stamoulis,¹⁶ J. Steinmann,⁴¹ B. Still,³⁸ Y. Suda,⁴⁵ A. Suzuki,²⁴ K. Suzuki,²⁵ S. Y. Suzuki,^{14,†} Y. Suzuki,^{23,23} T. Szegłowski,⁴³ R. Tacik,^{39,50} M. Tada,^{14,†} S. Takahashi,²⁵ A. Takeda,⁴⁶ Y. Takeuchi,^{24,23} H. K. Tanaka,^{46,§} H. A. Tanaka,^{4,‡} M. M. Tanaka,^{14,†} I. J. Taylor,³² D. Terhorst,⁴¹ R. Terri,³⁸ L. F. Thompson,⁴² A. Thorley,²⁷ S. Tobayama,⁴ W. Toki,⁸ T. Tomura,⁴⁶ Y. Totsuka,^{*} C. Touramanis,²⁷ T. Tsukamoto,^{14,†} M. Tzanov,²⁸ Y. Uchida,¹⁷ K. Ueno,⁴⁶ A. Vacheret,³⁵ M. Vagins,^{23,5} G. Vasseur,⁶ T. Wachala,¹³ A. V. Waldron,³⁵ C. W. Walter,^{9,§} D. Wark,^{44,35} M. O. Wascko,¹⁷ A. Weber,^{44,35} R. Wendell,^{46,§} R. J. Wilkes,⁵⁵ M. J. Wilking,³² C. Wilkinson,⁴² Z. Williamson,³⁵ J. R. Wilson,³⁸ R. J. Wilson,⁸ T. Wongjirad,⁹ Y. Yamada,^{14,†} K. Yamamoto,³⁴ C. Yanagisawa,^{32,**} T. Yano,²⁴ S. Yen,⁵⁰ N. Yershov,²² M. Yokoyama,^{45,§} T. Yuan,⁷ M. Yu,⁵⁸ A. Zalewska,¹³ J. Zalipska,³¹ L. Zambelli,^{14,†} K. Zaremba,⁵³ M. Ziembicki,⁵³ E. D. Zimmerman,⁷ M. Zito,⁶ and J. Żmuda⁵⁷

(T2K Collaboration)

¹Department of Physics, Centre for Particle Physics, University of Alberta, Edmonton, Alberta, Canada

²Laboratory for High Energy Physics (LHEP), Albert Einstein Center for Fundamental Physics, University of Bern, Bern, Switzerland

³Department of Physics, Boston University, Boston, Massachusetts, USA

- ⁴*Department of Physics and Astronomy, University of British Columbia, Vancouver, British Columbia, Canada*
- ⁵*Department of Physics and Astronomy, University of California, Irvine, Irvine, California, USA*
- ⁶*IRFU, CEA Saclay, Gif-sur-Yvette, France*
- ⁷*Department of Physics, University of Colorado at Boulder, Boulder, Colorado, USA*
- ⁸*Department of Physics, Colorado State University, Fort Collins, Colorado, USA*
- ⁹*Department of Physics, Duke University, Durham, North Carolina, USA*
- ¹⁰*Laboratoire Leprince-Ringuet, Ecole Polytechnique, IN2P3-CNRS, Palaiseau, France*
- ¹¹*Institute for Particle Physics, ETH Zurich, Zurich, Switzerland*
- ¹²*Section de Physique, University of Geneva, DPNC, Geneva, Switzerland*
- ¹³*H. Niewodniczanski Institute of Nuclear Physics PAN, Cracow, Poland*
- ¹⁴*High Energy Accelerator Research Organization (KEK), Tsukuba, Ibaraki, Japan*
- ¹⁵*Institut de Fisica d'Altes Energies (IFAE), Bellaterra (Barcelona), Spain*
- ¹⁶*IFIC (CSIC and University of Valencia), Valencia, Spain*
- ¹⁷*Department of Physics, Imperial College London, London, United Kingdom*
- ¹⁸*Dipartimento Interuniversitario di Fisica, INFN Sezione di Bari and Università e Politecnico di Bari, Bari, Italy*
- ¹⁹*Dipartimento di Fisica, INFN Sezione di Napoli and Università di Napoli, Napoli, Italy*
- ²⁰*Dipartimento di Fisica, INFN Sezione di Padova and Università di Padova, Padova, Italy*
- ²¹*INFN Sezione di Roma and Università di Roma "La Sapienza," Roma, Italy*
- ²²*Institute for Nuclear Research of the Russian Academy of Sciences, Moscow, Russia*
- ²³*Kavli Institute for the Physics and Mathematics of the Universe (WPI), Todai Institutes for Advanced Study, University of Tokyo, Kashiwa, Chiba, Japan*
- ²⁴*Kobe University, Kobe, Japan*
- ²⁵*Department of Physics, Kyoto University, Kyoto, Japan*
- ²⁶*Physics Department, Lancaster University, Lancaster, United Kingdom*
- ²⁷*Department of Physics, University of Liverpool, Liverpool, United Kingdom*
- ²⁸*Department of Physics and Astronomy, Louisiana State University, Baton Rouge, Louisiana, USA*
- ²⁹*Université de Lyon, Université Claude Bernard Lyon 1, IPN Lyon (IN2P3), Villeurbanne, France*
- ³⁰*Department of Physics, Miyagi University of Education, Sendai, Japan*
- ³¹*National Centre for Nuclear Research, Warsaw, Poland*
- ³²*Department of Physics and Astronomy, State University of New York at Stony Brook, Stony Brook, New York, USA*
- ³³*Department of Physics, Okayama University, Okayama, Japan*
- ³⁴*Department of Physics, Osaka City University, Osaka, Japan*
- ³⁵*Department of Physics, Oxford University, Oxford, United Kingdom*
- ³⁶*Laboratoire de Physique Nucléaire et de Hautes Energies (LPNHE), UPMC, Université Paris Diderot, CNRS/IN2P3, Paris, France*
- ³⁷*Department of Physics and Astronomy, University of Pittsburgh, Pittsburgh, Pennsylvania, USA*
- ³⁸*School of Physics and Astronomy, Queen Mary University of London, London, United Kingdom*
- ³⁹*Department of Physics, University of Regina, Regina, Saskatchewan, Canada*
- ⁴⁰*Department of Physics and Astronomy, University of Rochester, Rochester, New York, USA*
- ⁴¹*III. Physikalisches Institut, RWTH Aachen University, Aachen, Germany*
- ⁴²*University of Sheffield, Department of Physics and Astronomy, Sheffield, United Kingdom*
- ⁴³*Institute of Physics, University of Silesia, Katowice, Poland*
- ⁴⁴*Rutherford Appleton Laboratory, Harwell Oxford, and Daresbury Laboratory, STFC, Warrington, United Kingdom*
- ⁴⁵*Department of Physics, University of Tokyo, Tokyo, Japan*
- ⁴⁶*Kamioka Observatory, Institute for Cosmic Ray Research, University of Tokyo, Kamioka, Japan*
- ⁴⁷*Research Center for Cosmic Neutrinos, Institute for Cosmic Ray Research, University of Tokyo, Kashiwa, Japan*
- ⁴⁸*Department of Physics, Tokyo Metropolitan University, Tokyo, Japan*
- ⁴⁹*Department of Physics, University of Toronto, Toronto, Ontario, Canada*
- ⁵⁰*TRIUMF, Vancouver, British Columbia, Canada*
- ⁵¹*Department of Physics and Astronomy, University of Victoria, Victoria, British Columbia, Canada*
- ⁵²*Faculty of Physics, University of Warsaw, Warsaw, Poland*
- ⁵³*Institute of Radioelectronics, Warsaw University of Technology, Warsaw, Poland*
- ⁵⁴*Department of Physics, University of Warwick, Coventry, United Kingdom*
- ⁵⁵*Department of Physics, University of Washington, Seattle, Washington, USA*
- ⁵⁶*Department of Physics, University of Winnipeg, Winnipeg, Manitoba, Canada*

⁵⁷*Faculty of Physics and Astronomy, Wroclaw University, Wroclaw, Poland*⁵⁸*Department of Physics and Astronomy, York University, Toronto, Ontario, Canada*

(Received 13 March 2014; published 31 October 2014)

We report the first measurement of the neutrino-oxygen neutral-current quasielastic (NCQE) cross section. It is obtained by observing nuclear deexcitation γ rays which follow neutrino-oxygen interactions at the Super-Kamiokande water Cherenkov detector. We use T2K data corresponding to 3.01×10^{20} protons on target. By selecting only events during the T2K beam window and with well-reconstructed vertices in the fiducial volume, the large background rate from natural radioactivity is dramatically reduced. We observe 43 events in the 4–30 MeV reconstructed energy window, compared with an expectation of 51.0, which includes an estimated 16.2 background events. The background is primarily nonquasielastic neutral-current interactions and has only 1.2 events from natural radioactivity. The flux-averaged NCQE cross section we measure is $1.55 \times 10^{-38} \text{ cm}^2$ with a 68% confidence interval of $(1.22, 2.20) \times 10^{-38} \text{ cm}^2$ at a median neutrino energy of 630 MeV, compared with the theoretical prediction of $2.01 \times 10^{-38} \text{ cm}^2$.

DOI: [10.1103/PhysRevD.90.072012](https://doi.org/10.1103/PhysRevD.90.072012)

PACS numbers: 25.30.Pt, 21.10.Pc, 23.20.Lv, 29.40.Ka

I. INTRODUCTION

Nuclear deexcitation γ rays are a useful tool for detecting neutrino-nucleus neutral-current (NC) interactions where the final-state neutrino and associated nucleon are not observed in a Cherenkov detector. These interactions have previously been observed in neutrino-carbon interactions [1,2]. The most well-known γ -ray production process on oxygen is inelastic scattering, $\nu + {}^{16}\text{O} \rightarrow \nu + {}^{16}\text{O}^*$, where the residual oxygen nucleus can deexcite by emitting a nucleon or γ rays with energies between 1 and 10 MeV. This process can be used to detect supernova neutrinos [3], which have an average energy of 20–30 MeV. Most theoretical work on γ -ray production in NC interactions has been performed in this low neutrino energy range with the assumption that it is applicable up to neutrino energies of several hundred MeV [4–7].

A recent calculation of γ -ray production in neutrino NC interactions shows that quasielastic (QE) nucleon knockout $\nu + {}^{16}\text{O} \rightarrow \nu + p + {}^{15}\text{N}^*(\nu + n + {}^{15}\text{O}^*)$ overwhelms the inelastic process at $E_\nu \gtrsim 200 \text{ MeV}$ [8]. The NCQE cross section is calculated to be more than an order of magnitude larger than the NC inelastic cross section from Ref. [7] at $E_\nu \approx 500 \text{ MeV}$. We can observe this interaction in two ways: by observing the “primary” γ rays produced when the residual nucleus deexcites or by observing the “secondary” γ rays produced when knocked-out nucleons interact with other nuclei in the water. The primary γ rays occur only when the knocked-out nucleon comes from

the $1p_{3/2}$ or $1s_{1/2}$ states, while the secondary γ rays can be produced by nucleons released during deexcitation or by interactions of the original knocked-out nucleons and so can occur for all nucleon states. Both types of γ rays, produced in interactions of atmospheric neutrinos, are a major background for the study of astrophysical neutrinos in the 10 MeV range [9,10], and a direct measurement of the rate of this process with a known neutrino source will be useful for ongoing and proposed projects [11–14].

This paper reports the first measurement of the neutrino-oxygen NCQE cross section via the detection of deexcitation γ rays from both primary and secondary interactions. The neutrinos are produced by using the narrow-band neutrino beam at the Japan Proton Accelerator Research Complex (J-PARC) and measured with the Super-Kamiokande (SK) water Cherenkov detector.

II. THE T2K EXPERIMENT

The Tokai-to-Kamioka (T2K) experiment [15] is a long-baseline neutrino oscillation experiment consisting of a neutrino beam and several near detectors and using Super-Kamiokande as a far detector. It is designed to search for $\nu_\mu \rightarrow \nu_e$ appearance, which is sensitive to the neutrino mixing angle θ_{13} , and to precisely measure the mixing angle θ_{23} and the mass difference $|\Delta m_{32}^2|$ by ν_μ disappearance.

The accelerator at J-PARC provides a 30 GeV proton beam which collides with a graphite target to produce charged mesons. Positively charged pions and kaons are collected and focused by magnetic horns and ultimately decay in flight to produce primarily muon neutrinos inside a 96 m long cavity filled with helium gas. The proton beam is directed 2.5° away from SK. The off-axis neutrino beam has a narrow peak with median energy 630 MeV at SK because of the two-body decay kinematics of the π^+ which dominate the focused beam. This peak energy was chosen because it corresponds to the first maximum in the neutrino oscillation probability at the location of the far detector. The narrow energy peak also allows for the measurement of

*Deceased.

†Also at J-PARC, Tokai, Japan.

‡Also at Institute of Particle Physics, Canada.

§Also at Kavli IPMU (WPI), The University of Tokyo, Japan.

||Also at Moscow Institute of Physics and Technology and National Research Nuclear University “MEPhI,” Moscow, Russia.

¶Also at JINR, Dubna, Russia.

**Also at BMCC/CUNY, Science Department, New York, NY, USA.

the NC cross section at a particular energy. Typically, it is not possible to make energy-dependent measurements of this cross section, because the invisible outgoing neutrino makes accurate energy reconstruction impossible.

The T2K experiment has several near detectors located 280 m from the neutrino production target. The on-axis near detector, INGRID, which consists of 16 modules made up of alternating layers of iron and plastic scintillator arranged in a cross, monitors the neutrino beam direction. The off-axis near detectors, ND280, measure the neutrino beam spectrum and composition for the oscillation analyses. The neutrino measurements at the INGRID and ND280 detectors are consistent with expectations [16], but this information is not used to constrain systematic uncertainties in this analysis so that an absolute cross-section measurement can be made.

Super-Kamiokande [11] is a cylindrical water Cherenkov detector consisting of 50 ktons of ultrapure water, located 295 km from the neutrino target at J-PARC. It was built in the middle of Mount Ikenoyama, near the town of Kamioka, 1000 m below the peak. The tank is optically separated into two regions which share the same water. The inner detector (ID) is a cylinder containing the 22.5 kton fiducial volume and is instrumented with 11 129 inward-facing photomultiplier tubes (PMTs). The outer detector extends 2 m outward from all sides of the ID and is instrumented with 1885 outward-facing PMTs. It serves as a veto counter against cosmic-ray muons as well as a shield for γ rays and neutrons emitted from radioactive nuclei in the surrounding rock and stainless steel support structure.

III. EVENT SIMULATION

T2K events at SK are simulated in three stages. First, the neutrino beam line is simulated to predict the flux and energy spectrum of neutrinos arriving at SK. Next, the interactions of those neutrinos with the nuclei in the SK detector are simulated, including final-state interactions within the nucleus. Finally, the SK detector response to all of the particles leaving the nucleus is simulated.

FLUKA [17] is used to simulate hadron production in the target based on the measured proton beam profile. Hadron production data from NA61/SHINE at CERN [18,19] are used to tune the simulation and evaluate the systematic error. The regions of phase space not covered by the NA61/SHINE data directly are tuned by extrapolating the data using the empirical parameterization developed by Bonesini *et al.* [20]. Once particles leave the production target, they are transported through the magnetic horns, target hall, decay volume, and beam dump by using a GEANT3 [21] simulation with GCALOR [22] for hadronic interactions. The initial composition of the beam is 93% ν_μ , 6% $\bar{\nu}_\mu$, and 1% ν_e (the 0.1% $\bar{\nu}_e$ component is not considered in this analysis). The ν_μ ($\bar{\nu}_\mu$) flux is 94% (92%) from the decay in flight of pions, while approximately half of the ν_e flux is from muon decays and the other half is from the

kaon decays. A more detailed description of the neutrino flux prediction and its uncertainty can be found in Ref. [23].

Neutrino interactions based on the above flux are simulated by using the NEUT event generator [24,25]. The NCQE cross section on oxygen is simulated by using a spectral function model [26,27] with the BBBA05 form factor parameterization [28], which is then reweighted as a function of neutrino energy to match the recent theoretical calculations from Ref. [8]. In order to simulate the deexcitation γ -ray emission, it is necessary to identify which state the remaining nucleus is in after the neutrino interaction. In the simple shell model, the nucleons in ^{16}O occupy three states: $1p_{1/2}$, $1p_{3/2}$, and $1s_{1/2}$ with knockout energies of 12.1, 18.4, and 42 MeV, respectively (3.54 MeV more for neutrons), though precise electron scattering measurements have shown significant deviations from this mean-field scenario [29–31]. Approximately 20% of the total spectral strength is pushed by NN correlations out of the Fermi sea into continuum states [32]. The electron scattering results are incorporated into a realistic spectral function model in Ref. [26], which was integrated in momentum and energy in Ref. [8] to calculate the spectroscopic factors for the shell states shown in Table I.

Since the $1p_{1/2}$ hole is already the ground state, it produces no primary γ rays. The $1p_{3/2}$ proton and neutron hole states have three possible energy levels, the most common of which (branching ratio 87%) has the lowest energy and decays by releasing a 6.32 or 6.18 MeV photon for a proton or neutron hole, respectively. That is the only $1p_{3/2}$ neutron-hole decay that produces a photon, but for the proton holes one of the higher energy levels also releases a 9.93 MeV photon, bringing the total branching ratio from γ -ray production up to 92% [29,33]. The $1s_{1/2}$ hole state can decay via a variety of channels, usually including additional nucleon emission because of the large binding energy of the knocked-out nucleon. The branching ratios of the $1s_{1/2}$ proton hole state are estimated by using the result of the $^{16}\text{O}(p, 2p)^{15}\text{N}$ experiment (RCNP-E148) [34], with a 3–6 MeV γ ray produced 22% of the time and a > 6 MeV γ ray produced 15% of the time. Only protons were studied in the experiment, but the proton and neutron

TABLE I. The spectroscopic factors and branching ratios for primary γ -ray production for the three residual excited nuclear states simulated in this analysis. The spectroscopic factors are calculated in Ref. [8], while the branching ratios were measured with electron and proton scattering [29,33,34].

	$1p_{1/2}$	$1p_{3/2}$	$1s_{1/2}$
Spectroscopic factors	0.632	0.703	0.422
γ -ray branching ratios:			
> 6 MeV from p hole	0%	91.8%	14.7%
> 6 MeV from n hole	0%	86.9%	14.7%
3–6 MeV from either	0%	0%	27.8%

energy levels are expected to be very similar, since ^{16}O is an isoscalar nucleus. There have been no measurements of photon production from the continuum states, so they are assumed to produce no primary γ rays, though this assumption is taken into account in the systematic uncertainties in Sec. IV C.

Non-QE NC interactions make up the largest neutrino-induced background component and predominantly consist of NC single-pion production where the pion is absorbed during final-state interactions in the nucleus. This resonant production is simulated by using the Rein-Sehgal model [35], the position dependence within the nucleus is calculated with the model from Ref. [36], and the scale of the microscopic pion interaction probabilities in the nuclear medium is determined from fits to pion scattering data [37–39]. The simulation of primary deexcitation γ rays from this process is based on measurements of π^- absorption at rest on H_2O at CERN [40]. These pion-absorption interactions can also release nucleons which go on to produce secondary γ rays as described below. More details about NEUT, including the models used to simulate the smaller charged-current (CC) backgrounds, can be found in Refs. [15,25].

SK's GEANT3-based simulation [21] is used to transport all the particles leaving the nucleus through the detector, to produce and transport the Cherenkov light, and to simulate the response of the photodetectors and electronics. Charged pions with momenta above $500 \text{ MeV}/c$ are simulated with GCALOR [22], while lower momentum pions are simulated with a custom routine based on the NEUT cascade model for final state hadrons. GCALOR also simulates the interactions of nucleons with nuclei in the water, including the production of secondary γ rays. In this simulation, secondary γ rays are typically produced in multiples: 95% of events with secondary γ rays have at least two, some originating from multiple interactions of neutrons in the water. The total secondary γ -ray energy per event is distributed widely with a peak around 7 MeV and a long tail towards higher energies.

There is an additional signal-like contribution from the inelastic process, $\nu + ^{16}\text{O} \rightarrow \nu + ^{16}\text{O}^*$. However, since there is no accurate estimation of γ -ray production induced by the NC inelastic process in the T2K energy range, we do not subtract its contribution in the final result. If we assume that the rate of γ -ray production after a inelastic interaction is similar to that of a nucleon knockout reaction and extrapolate the NC inelastic cross section predicted in Ref. [7] to the energy region of this analysis, we expect its contribution to be no larger than a few percent of our final sample.

IV. ANALYSIS

The results presented in this paper are based on T2K run 1–3 data from 3.01×10^{20} protons on target [41]. The expected number of beam-related events after the

TABLE II. Observed and expected numbers of events in T2K runs 1–3. The CC samples are based on the flux at SK including three-flavor oscillations (parameters described in the text). The NC samples are based on the unoscillated flux. The ν_e NCQE events are treated as signal, but the $\bar{\nu}_\mu$ NCQE are considered background since there is a different predicted cross section for antineutrinos.

Beam-related expectation	ν_μ	ν_e	$\bar{\nu}_\mu$
NCQE	34.33	0.46	0.69
NC non-QE	11.59	0.26	0.45
CC	2.01	0.0014	0.025
Signal		34.80	
Background (beam)		15.02	
Beam-unrelated		1.20	
Observed events		43	

selections described in the next section are summarized in Table II, which categorizes them by neutrino flavor and interaction mode. For the computation of the CC components, we assume three-flavor oscillations with $|\Delta m_{32}^2| = 2.44 \times 10^{-3} \text{ eV}^2$, $\sin^2\theta_{23} = 0.50$, and $\sin^2 2\theta_{13} = 0.097$. The majority of the beam-related background comes from nonquasielastic NC events, in particular, single-pion production followed by pion absorption within the nucleus. The CC background comes from interactions where the outgoing charged lepton has low momentum and is misidentified as an electron or where the charged lepton itself is below the Cherenkov threshold but deexcitation γ rays are emitted.

The expected number of beam-unrelated events after all selections are applied is estimated to be 1.2 by sampling events at least $5 \mu\text{s}$ before the T2K beam trigger so that no beam-related activity is included. The measured event rate is normalized to the total live time of the analyzed beam spills. Since the beam-unrelated background is directly measured with data outside the beam window, the systematic uncertainty associated with it is small.

A. Event selection

The event trigger for this analysis requires events to be within the 1 ms T2K beam window and have at least 25 PMT hits within 200 ns. This trigger has an estimated efficiency or 99.5% for 4 MeV events and is the lowest threshold trigger used in any T2K or Super-K analysis. It is possible only because of the sharp reduction in accidental backgrounds due to the beam timing requirement.

The reconstruction of the event vertex, direction, and energy is the same as that used in the SK solar neutrino analysis [42]. The event vertex is found by a maximum likelihood fit to the timing residuals of the Cherenkov light, accounting for the dark noise rate [43]. The vertex resolution is approximately 125 cm at 4 MeV and improves to below 50 cm above 12 MeV. The event direction is reconstructed by comparing the observed hit pattern with

the MC expectation for a single electron ring using a likelihood function. The reconstructed energy is defined as the total energy of a single electron that would have produced all Cherenkov photons in the event. It is calculated by using the effective number of hit PMTs N_{eff} , which corrects for the rate of multiple hits on a single PMT, scattered and reflected light, the water quality, the dark noise rate, the photocathode coverage as a function of time and position, and the PMT quantum efficiency. The relationship between N_{eff} and energy is determined by using a MC simulation of monoenergetic electrons. We use this definition because it is used by the SK low-energy reconstruction tools, though we know many events have multiple particles and a variety of particle species. The vertex resolution, water quality, and energy scale are calibrated by using a variety of sources, described in detail in Ref. [44].

The first selections applied are a cut on the reconstructed energy, only allowing events between 4 and 30 MeV, a standard fiducial volume cut of 2 m from the detector wall, and a tighter event timing cut. The neutrino beam spill has a bunch structure, reflecting the underlying proton bunch structure, with 6 or 8 bunches separated by 581 ns gaps, delivered every 3 s. The ± 100 ns timing cut contains the neutrino beam bunch width which has an observed rms of 24 ns at SK. The time is synchronized between the near and far sites by using a common-view GPS system, and the bunch timing is calibrated by using the higher energy T2K neutrino events at SK.

Further selection cuts are applied based on the event vertex and reconstruction quality to remove beam-unrelated background, similar to those used in SK solar [42] and supernova relic neutrino analyses [9]. These cut criteria are simultaneously optimized in an energy-dependent way to maximize the figure of merit defined as $N_{\text{beam}} / \sqrt{N_{\text{beam}} + N_{\text{unrel}}}$, where N_{beam} and N_{unrel} denote the number of expected beam-related and beam-unrelated events, respectively. The cut optimization is done separately for each of the three T2K run periods, since the beam intensities and beam bunch structures differ.

Most of the beam-unrelated background comes from radioactive impurities in the PMT glass, cases, and support structure and so is concentrated near the ID wall. Cuts on the distance from the nearest wall, \mathcal{D}_1 , and the distance from the wall along the backward direction of the reconstructed track, \mathcal{D}_2 , together effectively eliminate background events produced at or near the ID wall. A minimum cut of 2 m is applied for both, with the cut on \mathcal{D}_1 increasing linearly below 4.75 MeV to about 3.2 m and the cut on \mathcal{D}_2 increasing linearly below 5.75 MeV to about 10 m.

Beam-unrelated background events that pass the fiducial cuts typically have reconstruction errors which move the vertex to the center of the tank. These errors can be identified based on the distribution of hits in time and space. The hit time distribution should be a sharp peak after time-of-flight correction from the correct vertex, which we

quantify as the timing goodness g_t . The hit pattern should also be azimuthally symmetric around the reconstructed particle direction, which we test by using g_p , the Kolmogorov-Smirnov distance between the observed hit distribution and a perfectly symmetric one. The reconstruction quality cut criterium Q_{rec} is defined as the hyperbolic combination of these two parameters: $Q_{\text{rec}} \equiv g_t^2 - g_p^2$ and is shown in Fig. 1. The cut on Q_{rec} is also energy dependent and varies from about 0.25 at its tightest at the low end of the energy spectrum to effectively no cut above 11 MeV. More detailed descriptions of g_t and g_p are found in Ref. [45].

Before selection, the beam-unrelated background rate from natural radioactivity is 284 counts per second, or 1.2 million events expected during the 1 ms beam windows used for other T2K analyses [46]. Applying the tight timing cut reduces this background to 1816 events. The fiducial and reconstruction quality cuts further reduce the beam-unrelated background to 1.77 events, a contamination of 2.4%. These beam-unrelated selection cuts reduce the estimated NCQE signal efficiency to 74%. Among the selected signal events, we estimate 97% have true vertices within the fiducial volume.

Finally, to suppress the beam-related CC interaction events, two additional cuts are applied: a preactivity cut and a Cherenkov opening angle cut. The preactivity cut rejects electrons produced in muon decays with more than 99.9% efficiency by rejecting events which occur less than 20 μs after a high-energy event, defined as a group of 22 or more hits in a 30 ns window. The likelihood of this selection rejecting a signal event because of accidental dark noise hits is less than 0.1%.




FIG. 1 (color online). Distribution of the reconstruction quality parameter Q_{rec} after the beam-unrelated selection cuts (timing, fiducial volume, Q_{rec}) have been applied. The inset shows the distribution before the energy-dependent cut on Q_{rec} but including the timing and fiducial volume cuts. The T2K run 1–3 data are represented by points with statistical error bars, and the expectation is represented by stacked histograms showing the NCQE signal and the NC non-QE, CC, and beam-unrelated background components.

For this low-energy sample, the Cherenkov angle of an event is defined as the peak of the distribution of Cherenkov angles calculated for every combination of three PMTs with hits, assuming the reconstructed vertex as the origin, following the technique from Ref. [9]. For a single particle this peak will be close to the opening angle of the particle, while the more isotropic light distributions from multiple particles will have peaks close to 90° . The Cherenkov angle depends on the velocity of the particle, approaching 42° as the velocity approaches c . The electrons produced by the deexcitation γ rays selected in this analysis are highly relativistic and so peak at 42° . The heavier muons from ν_μ CC events have smaller opening angles, peaking around 28° ; the higher momentum muons with larger opening angles having already been removed by the energy cut at 30 MeV. These low-momentum muons are removed by a cut at 34° . The Cherenkov angle distribution for events passing all other selection criteria can be seen in Fig. 2. The data-expectation disagreement in the multi- γ peak is likely due to the approximations made in the model of γ -ray emission induced by secondary neutron interactions used by GEANT3 and GCALOR, particularly the multiplicity of the secondary neutrons.

After all selections, 51.0 events are expected, of which 34.8 are expected to be NCQE signal for a purity of 69%. The overall selection efficiency is estimated to be 70% relative to the number of true NCQE events in the true fiducial volume which produce either primary or secondary γ rays (approximately 25% of NCQE events produce no photons and are consequently unobservable). The beam-unrelated contamination remains 2.4% after the final beam-related selections, with the 1.77 events after only




FIG. 2 (color online). The Cherenkov angle distribution in data and MC expectation after the beam-unrelated selections and the preactivity cut. The expectation has a three-peak structure corresponding to low-momentum muons around 28° , single γ rays around 42° , and multiple γ rays around 90° . A selection cut is applied at 34° to remove the muon events, but no attempt is made to separate single- and multiple- γ events.

the beam-unrelated selections reduced to 1.2 background events in the final sample.

B. Observed events

Figure 3 shows the observed event timing distribution in a region from -1 to $5 \mu\text{s}$ with respect to the beam trigger time, before the tight ± 100 ns timing cut on each bunch has been applied. Six events are found outside the tight bunch time windows, which is consistent with the 3.6 beam-unrelated events expected for this amount of integrated live time. These events are separate from the 1.2 beam-unrelated events expected to fall within the 200 ns bunch windows.

After all cuts, 43 events remain in the 4–30 MeV reconstructed energy range, compared with 51.0 expected. The vertex distribution of the sample is shown in Fig. 4, in which no nonuniformity or biases with respect to the neutrino beam direction are found. The energy distribution of the data after all the selection cuts is shown in Fig. 5. A peak due to 6 MeV prompt deexcitation γ rays is clearly seen in the data, and the observed distribution matches well with the expectation. The high-energy tail originates primarily from the contribution of additional secondary γ rays overlapping the primary γ rays.

C. Systematic uncertainties

The sources of systematic uncertainty on the expected number of signal and background events and their size are summarized in Table III. The methods for calculating these uncertainties are described below.

The flux errors, calculated in correlated energy bins, are determined based on beam monitoring, constraints from external measurements (particularly NA61/SHINE [18,19]), and Monte Carlo studies of focusing parameters




FIG. 3 (color online). ΔT_0 distribution of the data sample after all selection cuts except for beam timing, compared to the bunch center positions determined from high energy T2K neutrino events, indicated by eight dashed vertical lines. The on-timing and off-timing events are shown in solid and hashed, respectively.




FIG. 4 (color online). Vertex distribution of the final data sample in Y versus X after all selections have been applied. The solid and dotted lines indicate the boundaries of the inner detector and fiducial volume, respectively. The neutrino beam direction is indicated by the arrow.

(e.g., horn current, beam alignment, etc.) [23]. The dominant uncertainty on the flux comes from the production of hadrons in the target, though the beam alignment and off-axis angle become significant at energies between 700 MeV and 1.2 GeV, the high-energy side of the focusing peak. The neutrino interaction uncertainties which affect the normalization of the background are evaluated by comparing NEUT predictions to external neutrino-nucleus data sets in an energy region similar to T2K [16].

The 15% systematic uncertainty on primary γ -ray production in the signal (and the QE component of the CC background) comes from several sources. The largest contribution comes from the uncertainty on the $1p_{3/2}$




FIG. 5 (color online). Comparison of the reconstructed energy spectrum between the selected data sample and the expectation after all selection cuts have been applied. The CC component is actually about twice as large as the beam-unrelated background, but it is less apparent since it is spread across all energies.

TABLE III. Summary of systematic uncertainties on the expected number of signal and background events. While the CC component has the largest uncertainty, it has a relatively small effect on the final result, since there are relatively few CC events in the final sample.

Fraction of sample	Signal	Background		
	NCQE	NC non-QE	CC	Unrel.
Fraction of sample	68%	26%	4%	2%
Flux	11%	10%	12%	...
Cross sections	...	18%	24%	...
Primary γ production	15%	3%	9%	...
Secondary γ production	13%	13%	7.6%	...
Detector response	2.2%	2.2%	2.2%	...
Oscillation parameters	10%	...
Total systematic error	23%	25%	31%	0.8%

spectroscopic factor. We take the 34% difference between the value used in our simulation (from [8]) and the value calculated by Ejiri, 0.940 [47], which gives an 11% change in the number of selected NCQE events. The second-largest contribution is from final-state nuclear interactions: NEUT assumes that the deexcitation γ -ray production is the same whether the final state contains a single nucleon or multiple nucleons. We estimate the systematic uncertainty introduced by this assumption by observing the change in the number of signal events with the extreme alternate assumption that no deexcitation γ rays are released from events with multi-nucleon final states, which gives a 9% change in the number of selected NCQE. Both of these errors have a diluted effect on the final sample, since NCQE events can be observed via either primary or secondary γ rays.

The $1s_{1/2}$ spectral function has a 30% uncertainty, also calculated by model comparisons between Refs. [8] and [47], but only gives a 1% contribution to the final error. The uncertainty in the $1s_{1/2}$ branching ratios is also accounted for based on the measurement from Kobayashi *et al.* [34], giving an additional 1% error. The uncertainty due to not simulating photon production in the NN -induced continuum states is estimated by assuming these events emit photons in the same way as the $1s_{1/2}$ state, a reasonable approximation since both have binding energies well above the particle emission threshold, giving a 3% error. These systematics have only a small effect on the final selected sample, since only a small fraction of $1s_{1/2}$ deexcitations produce photons above 6 MeV.

For the non-QE NC background events, a conservative uncertainty was calculated by removing all primary γ rays from the events and evaluating the difference in total selected events. The effect is relatively small since the pion-absorption events which make up the bulk of the NC non-QE background produce many secondary γ rays and so are still detected thanks to the low threshold of the analysis.

The uncertainty on secondary γ -ray production is dominated by uncertainties on the production of neutrons. It was

evaluated by comparing alternate models of neutron production and interaction from *GALOR* and *NEUT* and how they altered the observed Cherenkov light level for our simulated events, for both signal events and the pion-absorption background. Even with changes of a factor of 2 or more in the average amount of energy deposited in the detector between the different neutron interaction models, the change in the number of selected events is relatively small because we only count events and do not attempt to distinguish between events with one or multiple γ rays. Events with secondary γ rays typically have multiple γ rays and often have primary γ rays, as well, so the detection efficiency is kept high even with a significant change in the secondary γ -ray production probability.

The detector uncertainty includes contributions from uncertainties in the SK energy scale, vertex resolution, and selection efficiency. It is estimated by comparing simulation and data from the linear electron accelerator installed above SK [48]. The systematic uncertainty due to the atmospheric oscillation parameters, θ_{23} and $|\Delta m_{32}^2|$, is estimated by varying the parameters within their uncertainties from the T2K measurement of these parameters [41].

There are two final systematic uncertainties that were evaluated but have a negligible impact on the result. We evaluated the potential nonuniformity of the selection efficiency with respect to Q^2 by changing the value of the MC axial mass to distort the differential cross section. This variation changes the final calculated cross section by less than a percent. The beam-unrelated background is estimated from the out-of-time events which have a statistical error of 0.8%.

V. MEASURED CROSS SECTION

The NCQE cross section is measured by comparing the NCQE cross section as calculated in recent theoretical work [8] averaged over the unoscillated T2K flux with the observed number of events after background subtraction:

$$\langle \sigma_{\nu, \text{NCQE}}^{\text{obs}} \rangle = \frac{N^{\text{obs}} - N_{\text{bkg}}^{\text{exp}}}{N^{\text{exp}} - N_{\text{bkg}}^{\text{exp}}} \langle \sigma_{\nu, \text{NCQE}}^{\text{theory}} \rangle, \quad (1)$$

where $\langle \sigma_{\nu, \text{NCQE}}^{\text{obs}} \rangle$ is the observed flux-averaged NCQE cross section and $\langle \sigma_{\nu, \text{NCQE}}^{\text{theory}} \rangle = 2.01 \times 10^{-38} \text{ cm}^2$ is the flux-averaged cross section from Ref. [8]. The NCQE cross section includes all spectral states, since even states without primary γ -ray emission are often observed with secondary γ rays. The total number of observed events is N^{obs} (43), the total number of expected events is N^{exp} (51.0), and $N_{\text{bkg}}^{\text{exp}}$ (16.2) denotes the expected number of background events.

The obtained flux-averaged neutrino-oxygen NCQE cross section is $1.55 \times 10^{-38} \text{ cm}^2$ at a median neutrino flux energy of 630 MeV. The 68% confidence interval on the cross section is $(1.20, 2.26) \times 10^{-38} \text{ cm}^2$ and the 90% confidence interval is $(0.96, 2.78) \times 10^{-38} \text{ cm}^2$.




FIG. 6 (color online). The T2K measurement of the flux-averaged NCQE cross section, represented by a black point, compared with the calculated cross section from Ref. [8]. The dashed line shows the cross section versus neutrino energy, and the solid horizontal line shows the flux-averaged cross section. The vertical error bar on the data represents the 68% confidence interval on the measured cross section, while the horizontal error bar is placed at the central value from our data and represents 68% of the flux at each side of the median energy. The solid gray histogram shows the unoscillated T2K neutrino flux.

They include both statistical and systematic errors and were calculated by using a Monte Carlo method to account for the systematic errors that are correlated between different samples. While the underlying systematic uncertainties are symmetric and Gaussian, the confidence interval is asymmetric around the central value because some of the uncertainties, primarily the production of secondary γ rays and to a lesser extent the neutrino flux, are correlated between the background expectation and the signal expectation which are found in the numerator and denominator, respectively, of Eq. (1). Figure 6 shows our result compared with a theoretical calculation of the NCQE cross section [8]. The vertical error bar for data shows the 68% confidence interval on the data, and the horizontal error bar represents 68% of the flux at each side of the median energy. The measurement is consistent with the recent theoretical calculation at the 68% confidence level. Using the same Monte Carlo method used to calculate the confidence intervals, we reject the background-only hypothesis (NCQE cross section of 0) with a p value of 4×10^{-8} , which corresponds to a one-sided z score of 5.4σ .

VI. CONCLUSION

We have reported the first measurement of the cross section of neutrino-oxygen NCQE interactions, measured via the detection of nuclear deexcitation γ rays in the Super-Kamiokande detector by using the T2K narrow-band neutrino beam. Our measurement is consistent with

the theoretical expectation at the 68% confidence level. Because of the similar peak energies for T2K neutrinos and atmospheric neutrinos, the present work will shed light on the study of the atmospheric background events for low-energy astrophysical phenomena in neutrino experiments. In this paper, we assume that oscillations do not modify the total flux of neutrinos at SK, but oscillations to a sterile neutrino (such as those considered in [49]) could lower the overall rate of all NC events, including NCQE events.

ACKNOWLEDGMENTS

We thank the J-PARC staff for superb accelerator performance and the CERN NA61 Collaboration for

providing valuable particle production data. We acknowledge the support of MEXT, Japan; NSERC, NRC, and CFI, Canada; CEA and CNRS/IN2P3, France; DFG, Germany; INFN, Italy; National Science Centre (NCN), Poland; RSF, RFBR, and MES, Russia; MINECO and ERDF funds, Spain; SNSF and SER, Switzerland; STFC, United Kingdom; and DOE, USA. We also thank CERN for the UA1/NOMAD magnet, DESY for the HERA-B magnet mover system, NII for SINET4, and the WestGrid and SciNet consortia in Compute Canada, GridPP, United Kingdom. In addition, participation of individual researchers and institutions has been further supported by funds from ERC (FP7), EU; JSPS, Japan; Royal Society, United Kingdom; and DOE Early Career program, USA.

-
- [1] B. Bodmann *et al.* (KARMEN Collaboration), *Phys. Lett. B* **280**, 198 (1992).
- [2] B. Armbruster *et al.* (KARMEN Collaboration), *Phys. Lett. B* **423**, 15 (1998).
- [3] K. Langanke, P. Vogel, and E. Kolbe, *Phys. Rev. Lett.* **76**, 2629 (1996).
- [4] E. Kolbe, K. Langanke, S. Krewald, and F. Thielemann, *Nucl. Phys. A* **540**, 599 (1992).
- [5] J. F. Beacom and P. Vogel, *Phys. Rev. D* **58**, 053010 (1998); **58**, 093012 (1998).
- [6] S. Nussinov and R. Shrock, *Phys. Rev. Lett.* **86**, 2223 (2001).
- [7] E. Kolbe, K. Langanke, and P. Vogel, *Phys. Rev. D* **66**, 013007 (2002).
- [8] A. M. Ankowski, O. Benhar, T. Mori, R. Yamaguchi, and M. Sakuda, *Phys. Rev. Lett.* **108**, 052505 (2012).
- [9] K. Bays *et al.* (Super-Kamiokande Collaboration), *Phys. Rev. D* **85**, 052007 (2012).
- [10] K. Ueno *et al.* (Super-Kamiokande Collaboration), *Astropart. Phys.* **36**, 131 (2012).
- [11] Y. Fukuda *et al.* (Super-Kamiokande Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **501**, 418 (2003).
- [12] J. F. Beacom and M. R. Vagins, *Phys. Rev. Lett.* **93**, 171101 (2004).
- [13] K. Abe *et al.*, arXiv:1109.3262.
- [14] T. Akiri *et al.* (LBNE Collaboration), arXiv:1110.6249.
- [15] K. Abe *et al.* (T2K Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **659**, 106 (2011).
- [16] K. Abe *et al.* (T2K Collaboration), *Phys. Rev. D* **87**, 092003 (2013).
- [17] G. Battistoni *et al.*, *Nucl. Instrum. Methods Phys. Res., Sect. B* **269**, 2850 (2011).
- [18] N. Abgrall *et al.* (NA61/SHINE Collaboration), *Phys. Rev. C* **84**, 034604 (2011).
- [19] N. Abgrall (NA61/SHINE Collaboration), *Phys. Rev. C* **85**, 035210 (2012).
- [20] M. Bonesini, A. Marchionni, F. Pietropaolo, and T. Tabarelli de Fatis, *Eur. Phys. J. C* **20**, 13 (2001).
- [21] R. Brun, F. Carminati, and S. Giani, Report No. CERN-W5013, 1994.
- [22] C. Zeitnitz and T. Gabriel, *Nucl. Instrum. Methods Phys. Res., Sect. A* **349**, 106 (1994).
- [23] K. Abe *et al.* (T2K Collaboration), *Phys. Rev. D* **87**, 012001 (2013).
- [24] M. Nakahata *et al.* (KAMIOKANDE Collaboration), *J. Phys. Soc. Jpn.* **55**, 3786 (1986).
- [25] Y. Hayato, *Nucl. Phys. B, Proc. Suppl.* **112**, 171 (2002).
- [26] O. Benhar, N. Farina, H. Nakamura, M. Sakuda, and R. Seki, *Phys. Rev. D* **72**, 053005 (2005).
- [27] O. Benhar, A. Fabrocini, S. Fantoni, and I. Sick, *Nucl. Phys. A* **579**, 493 (1994).
- [28] R. Bradford, A. Bodek, H. S. Budd, and J. Arrington, *Nucl. Phys. B, Proc. Suppl.* **159**, 127 (2006).
- [29] M. Leuschner *et al.*, *Phys. Rev. C* **49**, 955 (1994).
- [30] M. Bernheim *et al.*, *Nucl. Phys. A* **375**, 381 (1982).
- [31] K. Fissum *et al.* (Jefferson Lab Hall A Collaboration), *Phys. Rev. C* **70**, 034606 (2004).
- [32] D. Rohe *et al.*, *Phys. Rev. Lett.* **93**, 182501 (2004).
- [33] F. Ajzenberg-Selove, *Nucl. Phys. A* **523**, 1 (1991).
- [34] K. Kobayashi *et al.*, arXiv:nucl-ex/0604006.
- [35] D. Rein and L. M. Sehgal, *Ann. Phys. (N.Y.)* **133**, 79 (1981).
- [36] E. Oset, H. Toki, and W. Weise, *Phys. Rep.* **83**, 281 (1982).
- [37] D. Ashery, I. Navon, G. Azuelos, H. K. Walter, H. J. Pfeiffer, and F. W. Schlepütz, *Phys. Rev. C* **23**, 2173 (1981).
- [38] M. Jones *et al.*, *Phys. Rev. C* **48**, 2800 (1993).
- [39] R. Giannelli *et al.*, *Phys. Rev. C* **61**, 054615 (2000).
- [40] H. Engelhardt, C. Lewis, and H. Ullrich, *Nucl. Phys. A* **258**, 480 (1976).
- [41] K. Abe *et al.* (T2K Collaboration), *Phys. Rev. Lett.* **111**, 211803 (2013).
- [42] K. Abe *et al.* (Super-Kamiokande Collaboration), *Phys. Rev. D* **83**, 052010 (2011).
- [43] M. Smy, in *Proceedings of the Thirtieth International Cosmic Ray Conference*, Vol. 5, edited by R. Caballero,

- J. C. D'Olivo, G. Medina-Tanco, L. Nellen, F. A. Snchez, and J. F. Valdés-Galicia (Universidad Nacional Autónoma de México, Mexico City, 2008), pp. 1279–1282.
- [44] K. Abe *et al.*, *Nucl. Instrum. Methods Phys. Res., Sect. A* **737**, 253 (2014).
- [45] J. Cravens *et al.* (Super-Kamiokande Collaboration), *Phys. Rev. D* **78**, 032002 (2008).
- [46] K. Abe *et al.* (T2K Collaboration), *Phys. Rev. Lett.* **107**, 041801 (2011).
- [47] H. Ejiri, *Phys. Rev. C* **48**, 1442 (1993).
- [48] M. Nakahata *et al.* (Super-Kamiokande Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **421**, 113 (1999).
- [49] P. Adamson *et al.* (MINOS Collaboration), *Phys. Rev. Lett.* **107**, 011802 (2011).