



**HAL**  
open science

## Asymmetric electron capture in HCI collisions with rare gas dimers

J. Matsumoto, A. Leredde, X. Fléchar, S Shiromaru, Jimmy Rangama, C L Zhou, W. Iskandar, S. Guillous, D. Hennecart, A. Méry, et al.

► **To cite this version:**

J. Matsumoto, A. Leredde, X. Fléchar, S Shiromaru, Jimmy Rangama, et al.. Asymmetric electron capture in HCI collisions with rare gas dimers. 28<sup>e</sup> Internationale Conference on Photonic, Electronic and Atomic Collisions (ICPEAC2013), Jul 2013, Lanzhou, China. pp.102010, 10.1088/1742-6596/488/10/102010 . in2p3-01597281

**HAL Id: in2p3-01597281**

**<https://hal.in2p3.fr/in2p3-01597281>**

Submitted on 8 Jun 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

**OPEN ACCESS**

## Asymmetric electron capture in HCI collisions with rare gas dimers

To cite this article: J Matsumoto *et al* 2014 *J. Phys.: Conf. Ser.* **488** 102010

View the [article online](#) for updates and enhancements.

### Related content

- [Screening effect during a collision in multiple ionization of rare gas dimers by highly charged ions](#)  
Tomoko Ohyama-Yamaguchi and Atsushi Ichimura
- [The role of the partner atom and resonant excitation energy in ICD in rare gas dimers](#)  
Patrick O'Keeffe, Enrico Ripani, Paola Bolognesi et al.
- [Multiple ionization of rare gas dimers by slow highly charged ions: screening effect during a collision](#)  
Tomoko Ohyama-Yamaguchi and Atsushi Ichimura




**The Electrochemical Society**  
Advancing solid state & electrochemical science & technology

The ECS is seeking candidates to serve as the **Founding Editor-in-Chief (EIC) of ECS Sensors Plus**, a journal in the process of being launched in 2021

The goal of ECS Sensors Plus, as a one-stop shop journal for sensors, is to advance the fundamental science and understanding of sensors and detection technologies for efficient monitoring and control of industrial processes and the environment, and improving quality of life and human health.

*Nomination submission begins: May 18, 2021*



## Asymmetric electron capture in HCI collisions with rare gas dimers

J. Matsumoto<sup>\*1</sup>, A. Leredde<sup>†</sup>, X. Flécharde<sup>†</sup>, H. Shiromaru<sup>\*</sup>, J. Rangama<sup>#</sup>, C.L. Zhou<sup>#</sup>, W. Iskandar<sup>#</sup>,  
S. Guillous<sup>#</sup>, D. Hennecart<sup>#</sup>, A. Mery<sup>#</sup>, B. Gervais<sup>#</sup>, A. Cassimi<sup>#2</sup>

<sup>\*</sup> Department of Chemistry, Tokyo Metropolitan University, Hachioji, Tokyo 192-0397, Japan

<sup>†</sup> LPCCAen, ENSICAEN/UCBN/CNRS/IN2P3, Caen, France


<sup>#</sup> CIMAP, CEA/CNRS/ENSICAEN/UCBN, BP 5133, F-14070 Caen cedex 5, France

**Synopsis** Low-energy collisions between different rare gas dimers ( $\text{Ar}_2$ ,  $\text{Ne}_2$ ) and different projectiles ( $\text{O}^{3+}$ ,  $\text{Ar}^{9+}$ ,  $\text{Xe}^{20+}$ ) show that the weight of the different fragmentation processes, Coulomb explosion and Radiative Charge Transfer, strongly depends on the projectile charge state. This result is understood in term of impact parameter from which the electrons are captured on the projectile.

In a previous paper concerning ionic fragmentation of Ar dimers induced by electron-capture collision, we have shown that, due to reduced electron mobility across the dimer, the fragment ions preserve information of the electron removal location [1, 2]. Usually, such information is lost for covalent molecules, where the electrons freely move from one atom to the other, thus building the binding covalent bond. We report on fragmentation of different rare gas dimers ( $\text{Ar}_2$  and  $\text{Ne}_2$ ) multiply ionized by collisions with projectiles of different charge state ( $\text{O}^{3+}$ ,  $\text{Ar}^{9+}$  and  $\text{Xe}^{20+}$  ions) at an energy of 15 qkeV. The experiments were performed on the ARIBE beam line at the GANIL facility (Caen, France) using a COLTRIMS setup. The fragment ions from doubly to quadruply ionized dimers and the scattered projectile ions were detected in coincidence. The different fragmentation channels, the branching ratios as well as the kinetic energy release (KER) distributions were determined. This kinematically complete experiment allows the orientation of the dimer with respect to the projectile beam to be analyzed.

For all the projectile charge states, the KER spectra for double ionization of  $\text{Ar}_2$  and  $\text{Ne}_2$  followed by ionic fragmentation always show two peaks: one assigned to Coulomb explosion (CE) and the other to Radiative Charge Transfer (RCT) processes respectively. It has been observed that the RCT process is systematically reduced as the projectile charge state increases (Fig. 1). These results are readily understandable in the view of the classical-over-barrier model. The higher charge state projectiles capture electrons from the target at larger impact parameters, thus the preference in electron captures from one site, which is associated to RCT process, fades out. The correlation between the projectile scattering angle and target orientation

was derived for the asymmetric channels. The results show that the projectile is preferentially scattered in the same direction as the fragment with the higher charge state. Namely, the projectiles capture more electrons from the nearest site and then fragmentation occurs while keeping the primary dimer orientation. This scenario is again consistent with the semi-classical picture.



**Figure 1.** Kinetic Energy Release Distribution for the ( $\text{Ar}^+ + \text{Ar}^+$ ) fragmentation channel obtained with different projectile charge states. CE: Coulomb explosion process. RCT: Radiative Charge Transfer process.

### References

- [1] J. Matsumoto *et al* 2010 *Phys. Rev. Lett.* **105** 263202
- [2] J. Matsumoto *et al* 2011 *Phys. Scr.* **T144** 014016

<sup>1</sup> E-mail: junmatsu@tmu.ac.jp

<sup>2</sup> E-mail: cassimi@ganil.fr

