



HAL
open science

Search for CP-violation in $B \rightarrow D^* \mu \nu$ at LHCb

Vlad Dedu

► **To cite this version:**

Vlad Dedu. Search for CP-violation in $B \rightarrow D^* \mu \nu$ at LHCb. GDR-InF annual workshop, Nov 2021, Paris, France. in2p3-03457522

HAL Id: in2p3-03457522

<https://hal.in2p3.fr/in2p3-03457522>

Submitted on 30 Nov 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Search for CPV in $B \rightarrow D^* \mu \bar{\nu}_\mu$ at LHCb

Vlad Dedu

Aix Marseille Univ, CNRS/IN2P3, CPPM, IPhU, Marseille, France

GDR-InF annual workshop 16.11.2021




Introduction - CPV in $b \rightarrow c\ell\nu$ decays

- Semileptonic decays are interesting probes of NP
- $(RD - RD^*)$ anomalies can be explained by different NP models in $b \rightarrow c\ell\nu$
- This project: Search for CPV in $B \rightarrow D^*\ell\nu$, first analysis of CPV in a SL decay. Closely related to the angular analysis, but focus only on CPV terms.
- No CPV in SM in SL decays \rightarrow theoretically clean probe of NP.
- Sensitive to effects that could fake CP asymmetry
- Start with μ instead of τ channel: more statistics, easier analysis (τ reconstruction is difficult). Although NP should couple more to τ due to larger mass, same NP may affect μ . Can still provide constraints for NP.

Introduction - CP violation

- Search for Charge Parity Violation in $B \rightarrow D^* \ell \nu$ ($b \rightarrow c \ell \nu$) analysis



- In SM $B^0 \rightarrow D^* \ell \bar{\nu}_\ell$ has only 1 amplitude \rightarrow **no CPV**
- NP amplitude will have different weak phase but no strong phase (QCD amplitudes are the same for SM and NP)
 \rightarrow **not enough for direct CPV**

Possible ways to obtain CPV in SL decays:

- Four-(or more)-body B decays: Triple product asymmetries
- Interference of decay amplitudes with overlapping resonances, i.e. $B \rightarrow D^{**} \mu \nu$, with $D^{**} = D_0^*, D_1, D_2^*$

Introduction - Effective Hamiltonian

- Effective field theory for $b \rightarrow c \ell \bar{\nu}$ decays

$$\mathcal{H}_{\text{eff}}^{NP} = \frac{G_F V_{cb}}{\sqrt{2}} \sum_i g_i \mathcal{O}_i + \text{h.c.},$$

$$\mathcal{O}_S = \bar{c} b \ell (1 - \gamma_5) \nu$$

$$\mathcal{O}_P = \bar{c} \gamma_5 b \ell (1 - \gamma_5) \nu$$



$$\mathcal{O}_L = \bar{c} \gamma^\mu (1 - \gamma_5) b \ell \gamma_\mu (1 - \gamma_5) \nu$$

$$\mathcal{O}_R = \bar{c} \gamma^\mu (1 + \gamma_5) b \ell \gamma_\mu (1 - \gamma_5) \nu$$


$$\mathcal{O}_T = \bar{c} \sigma^{\mu\nu} (1 - \gamma_5) b \ell \sigma_{\mu\nu} (1 - \gamma_5) \nu$$

- $SM : g_S = g_P = g_L = g_R = g_T = 0; \mathcal{H}_{\text{eff}}^{SM} \propto \mathcal{O}_L$

- Couplings g_L, g_R, g_S, g_P, g_T can be complex.




Helicity angles



- $B^0 \rightarrow D^*(\rightarrow D^0 \pi) \mu \bar{\nu}_\mu$ decay is described by 4 kinematic parameters: 3 helicity angles ($\theta_\ell, \theta_D, \chi$) and q^2

Neutrino reconstruction







- Kinematic reco: Estimate B, ν momenta from decay topology using the B line of flight between PV and b-vertex [D. Hill et al]

$$|\vec{p}_{B^0}| = \frac{(m_Y^2 + m_{B^0}^2)|\vec{p}_Y| \cos \theta_{B^0, Y} \pm E_Y \sqrt{(m_{B^0}^2 - m_Y^2)^2 - 4m_{B^0}^2 |\vec{p}_Y|^2 \sin^2 \theta_{B^0, Y}}}{2(E_Y^2 - |\vec{p}_Y|^2 \cos^2 \theta_{B^0, Y})}$$

- Run full refit (DecayTreeFit) of the decay tree including all possible info: missing ν , vertex, mass constraints
- Improve precision in reconstructing quantities of interest ($\theta_L, \theta_D, \chi, q^2$)

Angle resolutions after DTF - simulation



Triple products

$$\frac{d^4\Gamma}{dq^2 d\cos\theta_\ell d\cos\theta^* d\chi} = \frac{3}{8\pi} \frac{G_F^2 |V_{cb}|^2 (q^2 - m_\ell^2)^2 |p_{D^*}|}{2^8 \pi^3 m_B^2 q^2} \times \mathcal{B}(D^* \rightarrow D\pi) \left(N_1 + \frac{m_\ell}{\sqrt{q^2}} N_2 + \frac{m_\ell^2}{q^2} N_3 \right)$$

- Angular distribution derived from most general $\mathcal{H}_{\text{eff}}^{NP}$ [D.London et al]
- NP couplings \rightarrow CPV terms (triple products) $\propto \sin\chi$
- Same magnitude and sign for $B0$ and $\bar{B}0$

Not suppressed	Coupling	Angular Function
$\text{Im}(\mathcal{A}_\perp \mathcal{A}_0^*)$	$\text{Im}[(1 + g_L + g_R)(1 + g_L - g_R)^*]$	$-\sqrt{2} \sin 2\theta_\ell \sin 2\theta^* \sin \chi$
$\text{Im}(\mathcal{A}_\parallel \mathcal{A}_\perp^*)$	$\text{Im}[(1 + g_L - g_R)(1 + g_L + g_R)^*]$	$2 \sin^2 \theta_\ell \sin^2 \theta^* \sin 2\chi$
$\text{Im}(\mathcal{A}_{SP} \mathcal{A}_{\perp,T}^*)$	$\text{Im}(g_P g_T^*)$	$-8\sqrt{2} \sin \theta_\ell \sin 2\theta^* \sin \chi$
$\text{Im}(\mathcal{A}_0 \mathcal{A}_\parallel^*)$	$\text{Im}[(1 + g_L - g_R)(1 + g_L + g_R)^*]$	$-2\sqrt{2} \sin \theta_\ell \sin 2\theta^* \sin \chi$

- Exploit orthogonality of Fourier terms and perform Fourier transformation of scattered data to obtain coefficient
- We construct a set of 50 orthogonal terms (up to 2nd order harmonics) \rightarrow control terms

Sensitivity study - HAMMER



[HAMMER website: paper,
manual, git etc]


- "A fast and efficient means to reweight large MC samples to any desired NP, or to any description of the hadronic matrix elements"
- Set FF scheme (BLPR) and Wilson coefficients (complex) values

$$B \rightarrow (D^* \rightarrow D\pi)\ell\nu \quad \Bigg| \quad \text{ISGW2}^*, \text{BGL}^{*\dagger}, \text{CLN}^{*\dagger}, \text{BLPR}^\dagger$$

- Calculate weight to NP scenario for each event based on true 4-momenta and particle IDs


CPV terms. Sensitivity study with simulation

- Inject NP in MC with HAMMER (reweighs each event based on NP: $g_L, g_R, g_P, g_T \neq 0$)



Stat. error
 $\sim 0.1\%$

LHCb simulation
true angles



Systematic uncertainties

We would need to control any systematics that could introduce "fake CP-asymmetry" at the 0.1% level

Non-zero $\sin \chi$ terms: what does it practically mean?


ν direction is reconstructed from topology of PV and secondary vertices.

- Term $\propto \sin \chi$: up-down asymmetry $(N_{\uparrow} - N_{\downarrow})/N_{\text{tot}}$.
- ν "up" \Leftrightarrow PV "below" $D^0\pi^+\mu^-$ plane
- ν "down" \Leftrightarrow PV "above" $D^0\pi^+\mu^-$ plane

What experimental effects can introduce non-zero "PV below-above" asymmetry?

- CPV in backgrounds
- **VELO misalignment**
- **Asymmetry of tracking efficiency**

Systematic uncertainties - VELO misalignment



- Misalignments of VELO as a whole should not introduce bias in angles, but displacements of the two halves wrt each other can
- Expect T_y and R_x to show largest source of bias

Misalignment in VELO


- VELO misalignment can affect the angles and introduce bias in CP asymmetry
- $\vec{\Theta} = \Theta_i \equiv (\theta_D, \theta_\ell, \chi, q^2)$, $P(\vec{\Theta})$ is an angular term

$$A_{CP} = \frac{1}{N} \sum_{\alpha=1}^N P(\vec{\Theta}_\alpha) = \langle P(\vec{\Theta}) \rangle$$

$$\delta A_{CP} = \sum_k \rho_k \left\langle \frac{\partial P}{\partial \Theta_i}(\vec{\Theta}) \frac{\partial \Theta_i}{\partial \rho_k}(\vec{\Theta}) \right\rangle$$

- Calculate partial derivative for each event (α) by introducing small displacement in tracks i.e.:

$$\theta'_{D,\alpha} = \theta_{D,\alpha} + \frac{\partial \theta_{D,\alpha}}{\partial \Delta x} \Delta x$$



- Typical VELO alignment precision: $1 \mu\text{m}$ for $T_{x,y}$, $3 \mu\text{m}$ for T_z , $3 \mu\text{rad}$ in $R_{x,y}$, $10 \mu\text{rad}$ in R_z
- Largest source of bias is T_y , up to 0.05%
- Can use control terms to control misalignment systematics in a data-driven way

Conclusions and outlook

Conclusions:

- $B \rightarrow D^* \mu \nu$: indirect CPV from angular distribution.
- MC study for sensitivity to CPV \rightarrow a few % with stat error $\sim 0.1\%$
- Helicity angle resolution studies (10-20 % improvement with DTF)
- Systematic uncertainties - detector misalignments

Outlook:

- Estimate all systematics: CPV in backgrounds and non-uniform detector efficiencies (similar approach as for misalignment)
- Look in data
- τ channel

BACK-UP SLIDES

Fourier transform method

- $P(\theta_\ell, \theta_D, \chi) = \sum_n C_n F_n(\theta_\ell, \theta_D, \chi)$, F_n are all orthogonal
- $C_k = \int P(\theta_\ell, \theta_D, \chi) F_k(\theta_\ell, \theta_D, \chi) d\theta_\ell d\theta_D d\chi$
- $P(\theta_\ell, \theta_D, \chi) = \frac{1}{n} \sum_{i=1}^n \delta(\theta_\ell - \theta_\ell^{(i)}) \delta(\theta_D - \theta_D^{(i)}) \delta(\chi - \chi^{(i)})$
- $C_k = \frac{1}{n} \sum_{i=1}^n F_k(\theta_\ell^{(i)}, \theta_D^{(i)}, \chi^{(i)})$
- Example:
 - $F_k(\theta_\ell, \theta_D, \chi) = \sin 2\theta_\ell \sin 2\theta_D \sin \chi$
 - $C_k = \frac{1}{n} \sum_{i=1}^n (\sin 2\theta_\ell^{(i)} \sin 2\theta_D^{(i)} \sin \chi^{(i)})$